

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/102848/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Rashid, G.M.M., Durán-Peña, M.J., Rahmanpour, R., Sapsford, Devin J. and Bugg, T.D.H. 2017. Delignification and enhanced gas release from soil containing lignocellulose by treatment with bacterial lignin degraders. *Journal of Applied Microbiology* 123 (1) , pp. 159-171. 10.1111/jam.13470 file

Publishers page: <http://dx.doi.org/10.1111/jam.13470> <<http://dx.doi.org/10.1111/jam.13470>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Delignification and Enhanced Methane Release from Soil Containing Lignocellulose by**
2 **Treatment with Bacterial Lignin Degraders**

3
4 Goran M.M. Rashid^a, Maria Jesus Duran-Pena^a, Rahman Rahmanpour^a, Devin Sapsford^b, and

5 Timothy D.H. Bugg^{a*}

6 ^aDepartment of Chemistry, University of Warwick, Coventry CV4 7AL, U.K.

7 ^bSchool of Engineering, Cardiff University, Queen's Building, The Parade, Cardiff,

8 CF24 3AA, U.K.

9 Corresponding author: Prof. T.D.H. Bugg, Department of Chemistry, University of Warwick,

10 Coventry CV4 7AL, U.K. Email T.D.Bugg@warwick.ac.uk, Tel 44-2476-573018

11 Running title: Bacterial delignification in soil

12
13 **Abstract**

14 Aims: The aim of the study was to isolate bacterial lignin-degrading bacteria from municipal solid
15 waste soil, and to investigate whether they could be used to delignify lignocellulose-containing soil,
16 and enhance methane release.

17 Methods and Results: A set of 20 bacterial lignin degraders, including 11 new isolates from
18 municipal solid waste soil, were tested for delignification and phenol release in soil containing 1%
19 pine lignocellulose. A group of 7 strains were then tested for enhancement of methane gas release
20 from soil containing 1% lignocellulose in small-scale column tests. Using an aerobic pre-treatment,
21 aerobic strains such as *Pseudomonas putida* showed enhanced gas release from the treated sample,
22 but four bacterial isolates showed 5-10 fold enhancement in gas release in an *in situ* experiment
23 under microanaerobic conditions: *Agrobacterium* sp., *Lysinibacillus sphaericus*, *Comamonas*
24 *testosteroni*, and *Enterobacter* sp..

25 Conclusions: The results show that facultative anaerobic bacterial lignin degraders found in landfill
26 soil can be used for *in situ* delignification and enhanced methane release in soil containing
27 lignocellulose.

28 Significance & impact of the study: The study demonstrates the feasibility of using an *in situ*
29 bacterial treatment to enhance gas release and resource recovery from landfill soil containing
30 lignocellulosic waste.

31

32 Keywords: Delignification; lignin degradation; bacterial treatment; methane release; municipal
33 waste treatment.

34

35 **Introduction**

36 The commitment of land for municipal waste landfill, and the eventual recovery and reuse of
37 that land for other purposes, are issues of social and commercial interest, especially in densely
38 populated parts of the world. The potential recovery of valuable resources from such landfill sites,
39 especially valuable metals, is also of growing interest, and represents a new technological challenge
40 (Jones *et al.*, 2013; Tonini *et al.*, 2013). If the rate of biodegradation of the landfill contents could
41 be enhanced using biotechnology (Mali *et al.*, 2012; Ni *et al.*, 2016), then the release of methane
42 from landfill sites which provides gas for commercial or private energy generation would be
43 accelerated, and the time needed for reuse of the land should be reduced, both of which would be
44 valuable for commercial landfill operators, and for regional town planning purposes. Addition of
45 bacterial culture to a compostable municipal solid waste (MSW) bioreactor has been shown to
46 increase methane production by 25% (Mali Sandip *et al.*, 2012), and aerobic pre-treatment of MSW
47 has also been shown to improve digestibility and methane production (Ni *et al.*, 2016).

48 Lignocellulosic waste present in landfill sites is broken down slowly, since the lignin
49 fraction is recalcitrant to microbial breakdown (Sanchez, 2009). Moreover, the documented
50 pathways for microbial breakdown are oxidative, aerobic pathways (Bugg *et al.*, 2011), whereas
51 degradation in landfill sites is largely anaerobic (Yazdani *et al.*, 2010). The aromatic heteropolymer
52 lignin is highly resistant to breakdown, since it is linked together via ether C-O and C-C bonds that
53 are not susceptible to hydrolytic cleavage (Bugg *et al.*, 2011a). Microbial degradation of lignin has
54 been studied mainly in basidiomycete fungi (Sanchez, 2009; Bugg *et al.*, 2011a), which grow only
55 in aerobic environments, but in recent years there has been renewed interest in bacterial lignin
56 degradation (Ahmad *et al.*, 2010; Bugg *et al.*, 2011b), which offer potential applications in
57 biotechnology, due to the relative ease of protein expression for bacterial enzymes, and genetic
58 tools available for genetic manipulation in bacteria (Bugg *et al.*, 2011b). A number of lignin-
59 oxidising bacteria have been isolated from environmental soil samples (DeAngelis *et al.*, 2011a;
60 Taylor *et al.*, 2012), and bacterial DyP-type peroxidase enzymes have been discovered that can

61 oxidise lignin in *Rhodococcus jostii* RHA1 (Ahmad *et al.*, 2011), *Amycolatopsis* sp. 75iv2 (Brown
62 *et al.*, 2012), and *Pseudomonas fluorescens* Pf-5 (Rahmanpour & Bugg, 2015), and extracellular
63 manganese superoxide dismutase enzymes with activity for lignin oxidation have been discovered
64 in *Sphingobacterium* sp. T2 (Rashid *et al.*, 2015).

65 In this work we wished to examine the hypothesis that bacterial lignin-degrading strains or
66 lignin-degrading enzymes could be used for delignification of lignocellulosic waste in soil, and
67 hence could be used to enhance the rate of gas release from MSW-containing soil. Kumar and co-
68 workers have previously shown that the application of fungal Mn peroxidase on a small scale could
69 enhance the rate of gas release from lignin-rich waste materials (Jayasinghe *et al.*, 2011), and
70 addition of enzyme to anaerobic bioreactors also enhanced methane production (Hettiaratchi *et al.*,
71 2014), and Feng *et al.* have shown that addition of fungal Mn peroxidase to lignocellulolytic waste
72 composting enhanced carbon utilization (Feng *et al.*, 2011), therefore it seemed feasible that an
73 equivalent bacterial lignin-oxidising enzyme could be used in a similar fashion. The use of a
74 bacterial strain for delignification *in situ* in soil is a more challenging application, since lignin
75 degradation is currently believed to be an exclusively aerobic process (Bugg *et al.*, 2011a), and
76 while the topsoil layer of soil is aerobic, the micro-organisms responsible for gas production are
77 strictly anaerobic (Yazdani *et al.*, 2010). Here we report that selected bacterial isolates can be used
78 for delignification and enhancement of gas production in small-scale and lab-scale experiments.

79

80 **Materials & Methods**

81 *Growth media.*

82 Bacteria were grown in either M9 minimal media (unbuffered) or Luria-Bertani media, at 30 °C, in
83 some cases containing additives as described below.

84

85 *Isolation of lignin-degrading bacteria from MSW-containing soil*

86 Soil was collected from a municipal landfill site at Sandford Farm (Woodley/Reading, UK), and
87 was stored in a sealed container at -20 °C prior to use. Method A: 25 mg of waste sample and 10
88 mg of wheat straw were mixed, to which 5 mL of M9 salts added. The mixture was incubated at
89 30°C for 21 days with shaking, then 200 µL of enriched sample was streaked on M9 plates
90 containing 1.5% (w/v) Bacto-agar and 2% wheat straw). The plate was sprayed with nitrated pine
91 lignin (nitrated lignin was prepared as described by Ahmad *et al.*, 2010), colonies with different
92 appearance (colour, size or shape) were picked and transferred into 10 mL Luria Bertani (LB) broth
93 and grown overnight at 30°C with shaking at 180 rpm. Samples from selected colonies were taken
94 for Gram staining and 16S rRNA gene was amplified by polymerase chain reaction, and submitted
95 for DNA sequencing, and sequences analysed using the BLAST algorithm on the EBI server
96 (www.ebi.ac.uk). Oligonucleotide primers for amplification of 16S rRNA: Forward 5'-
97 AGAGTTTGATCMTGGCTCAG-3' and reverse 5'-TACGGYTACCTTGTTACGACTT-3' .

98
99 Method B: Soil samples were incubated in M9 minimal media containing 0.5% alkali Kraft lignin
100 (Sigma-Aldrich, ref 471003) for 3 days, and then plated on M9 agar plates containing 0.5% alkali
101 Kraft lignin, and colonies picked, and re-streaked on the same media. This method resulted in 2
102 further isolates, whose 16S rRNA gene was amplified by polymerase chain reaction, and submitted
103 for DNA sequencing.

104
105 *Delignification of lignocellulose (pine) using lignin-degrading bacteria*

106 Bacterial strains were grown overnight in Luria-Bertani broth at 30°C (10 mL), then cells were
107 harvested by centrifugation (5000 rpm, 10 min), and the cell pellets were washed with M9 minimal
108 media (2 mL) and re-pelleted by centrifugation, and then re-suspended in M9 minimal media (2
109 mL). 100 µL of suspended bacterial cells were inoculated in 10 mL of M9 salts containing 1g of
110 pine powder and 0.1% glucose, then the cultures were grown at 30°C in a shaking incubator (180
111 rpm) for one week. The treated lignocellulose was filtered through Whitman no. 1 filter paper,

112 washed with distilled water, and dried prior to estimation of lignin content using the Klason method
113 (see below). The supernatant was filtered through 0.2 μm syringe filter and its phenolic content
114 estimated using Folin-Ciocalteu method (see below).

115 The Klason assay for lignin determination is based on a published method (Kirk & Obst, 1988).
116 Lignocellulose samples were dried in oven at $110^{\circ}\text{C} \pm 2$ for 2 hr, of which $0.25 \text{ g} \pm 0.001$ placed in a
117 25 mL conical flask. The samples were digested with chilled concentrated H_2SO_4 ($3.75 \pm 0.02 \text{ mL}$)
118 for 2 hr at room temperature, with stirring. The hydrolysates were diluted with deionised water (140
119 mL), then refluxed for 4 hr in round bottom flasks. The residues were collected by filtration (using
120 Whatman® no.1 filter paper) and washed three times with deionised water. The residue (insoluble
121 lignin) was dried at 110°C for 1 hr and its weight was measured on a 4-figure balance, from which
122 the percentage lignin was calculated.

123 The Folin-Ciocalteu assay is adapted from a published method (Meda *et al.*, 2005). The general
124 method involved the successive addition of 80 μL of deionised water and 50 μL Folin's reagent
125 (Sigma Aldrich) to 20 μL of supernatant from bacterial treatment, or standard (*p*-hydroxybenzoic
126 acid used as a standard with concentration of 50, 100, 200, 300 and 400 $\mu\text{g/mL}$). The mixture was
127 incubated for 3 min at room temperature, then 250 μL of 20% sodium carbonate was added, and
128 samples incubated in the dark for 30 min. Absorbance was then measured at 760 nm, and total
129 phenol content was expressed as g/L based on *p*-hydroxybenzoic acid as standard. For measurement
130 of phenol release at different times, bacterial strains were inoculated and grown on M9 minimal
131 media (10 mL) containing 1% pine lignocellulose as described above for 10 days. Samples (100
132 μL) for phenol assay were taken after 2, 4, 6, 8 and 10 days of incubation.

133

134 *Enhancement of methane production by addition of lignin-degrading strains (small scale)*

135 Syringe columns were made by adding 16 g of MSW soil mixed with 1% chopped pine to a 10 mL
136 plastic syringe, to which was added 6.25 g/L sodium acetate (see Supporting Information Figure S1

137 for illustration). The top of each syringe was sealed to be gas-tight using the rubber stopper from
138 the syringe plunger, which was pierced with a plastic tube to supply fresh media, and a second
139 plastic tube to collect gas. The second tube was connected to gas collection unit, comprising an
140 upturned 5 mL syringe containing saturated NaCl solution, into which gas was bubbled. Bacterial
141 culture (100 μ L in M9 minimal media) was applied to all columns after 4 days of incubation except
142 for a control column in which 100 μ L in M9 salts was added. The volume of generated gas was
143 measured by eye using the printed volume scale at 2-day intervals for 36 days.

144

145 *Enhancement of methane production in two-step process (aerobic delignification followed by*
146 *anaerobic digestion)*

147 Bacterial strains were grown in M9 minimal media (10 mL) containing 150 mg pine powder
148 (autoclaved), supplemented with 0.1% glucose, at 30°C with shaking at 180 rpm for 7 days. The
149 grown bacterial cultures were used as starter culture for delignification experiments as described
150 below.

151 Pine powder (4.5 g) was autoclaved with 150 mL deionised water, then 150 mL of M9 salts
152 (2 x final concentration) was added, followed by 300 μ L of the starter culture of the bacterial lignin-
153 degrading strain. The resulting mixture was incubated with shaking at 180 rpm for another 7 days at
154 30°C. The solid pine residue was harvested by filtration on Whitman no. 1 filter paper and dried in
155 oven at 110°C for 2 hr, to give a “treated pine lignocellulose” sample. As a control experiment, pine
156 powder (5 g) was also de-lignified using thermochemical method by reflexing the lignocellulose
157 with 95% ethanol containing 5% HCl (0.2 M) for 4 hr. The treated pine residue was isolated by
158 filtration as described above, and dried at 110°C for 2 hr. The treated pine lignocellulose samples,
159 thermochemically treated pine lignocellulose, and a sample of pine organosolv lignin were then
160 used as substrates for methane generation via anaerobic digestion.

161 De-lignified (biologically or thermochemically) pine samples (2 g) or organosolv lignin (0.5
162 g) were mixed with 1 g of MSW soil, then deionised water (5 mL) was added to a plastic syringe as
163 described above, sealed and connected to gas collection unit. Gas generation was monitored in 2
164 day intervals for 40 days. Samples (200 μ L) for metabolite analysis by LC/MS analysis were
165 collected at 7 day intervals, at which point 200 μ L of deionised water was added to each column.

166

167 *Large scale methane production*

168 Commercially available softwood bark chips (J. Arthur Bowers, Wyevale Garden Centres,
169 50 g) were mixed with organic peat-free compost (New Horizon, Wyevale Garden Centres, 450 g)
170 then placed into an apparatus constructed from a polypropylene 2 L plastic bottle (see Supporting
171 Information Figure S2 for illustration) containing a gas-tight inlet at the top, connected to a gas
172 measuring unit, and an outlet on the bottom to collect liquid samples for metabolite analysis.
173 Deionised water was added (approximately 1 L) until all materials were submerged. A bacterial
174 starter culture (5 mL overnight culture grown in Luria-Bertani media) was added to the column,
175 then the column was sealed and connected to a gas collection unit and kept at room temperature.
176 Methane production was monitored for 35 days. Samples for metabolite analysis were taken every 7
177 days up to 4 weeks. Control experiments were also prepared as follows: 1) experiment in which
178 wood chips was replaced with compost; 2) experiment without bacterial inoculum; 3) experiment
179 containing additional bacterial inoculum from anaerobic digester (5 mL).

180 Gas samples for GC analysis were taken after 7, 14, and 21 days after inoculation. The
181 percentage of the generated methane in the gas samples was determined using an Agilent 7890B gas
182 chromatograph equipped with FID detector. Samples from residue (compost and wood chips) were
183 also taken after 6 weeks of inoculation for determination of its lignin content, using Klason assay
184 (see above), which were compared to the corresponding controls (untreated compost and wood
185 chips).

186

187 *Metabolite analysis via LC-MS*

188 Samples (0.2-5 mL) for LC/MS analysis were extracted with ethyl acetate (1-5 mL) after
189 acidification to pH 1 with 1 M HCl, then dried (Na₂SO₄), and evaporated under reduced pressure.
190 Samples were then re-dissolved in 300 µL of 1:1 MeOH: H₂O. Aliquots (50 µL) were injected onto
191 a Phenomenex Luna C₁₈ reverse phase column (5 µm, 100 Å, 50 x 4.6 mm) on an Agilent 1200 and
192 Bruker HCT Ultra mass spectrometer, at a flow rate of 0.5 mL/min, monitoring at 310 and 270 nm.
193 The solvents were water 0.1% formic acid as solvent A and MeOH 0.1% formic acid as solvent B.
194 The gradient was 5-30% B from 0-30 min; 30-40% from 30-35 min, 40-70% from 35-40 min, 70-
195 100% from 40-45 min, 100% solvent B continued from 45-57 min and followed by 100-5% solvent
196 B for 3 min.

197

198 *Incubation of β-aryl ether lignin model compound with anaerobic consortium*

199 Guaiacylglycerol-β-guaiacyl ether (Tokyo Chemical Industry UK Ltd., 1 mM) was added to
200 30 mL M9 salts to which 1 mL of anaerobic culture from municipal solid waste was added, then the
201 headspace gas replaced with nitrogen. Samples (200 µL) were taken at 0, 2, 8 and 16 days after
202 incubation (at room temperature).

203

204 **Results**

205

206 **Isolation of lignin-degrading bacteria from MSW-containing soil**

207 We first examined whether lignin-degrading bacteria could be isolated from MSW-containing
208 soil, and if so, whether they are similar classes of bacteria to those isolated previously from
209 woodland soil (Taylor *et al.*, 2012). Two screening methods were used: firstly, a method previously
210 published by our group, involving the use of a nitrated lignin assay as a screen for lignin-degrading

211 bacteria on agar plates (Ahmad *et al.*, 2010; Taylor *et al.*, 2012); and secondly, growth of soil
212 samples on minimal media containing Kraft lignin.

213 For the first method, samples of MSW-containing soil were incubated for 14 days in minimal
214 M9 media containing 1% pine lignocellulose, in order to enhance the population of lignin-
215 degrading bacteria. Samples were then plated out on M9 minimal media agar plates containing 1%
216 pine lignocellulose for 3 days, then sprayed with a solution of nitrated lignin, and incubated
217 overnight at 30 °C. Colonies showing yellow coloration were picked and re-streaked in order to
218 obtain single bacterial isolates. This method resulted in 12 isolates, whose 16S rRNA gene was
219 amplified by polymerase chain reaction, and submitted for DNA sequencing. For the second
220 method, soil samples were incubated in M9 minimal media containing 0.5% Kraft lignin for 3 days,
221 and then plated on M9 agar plates containing 0.5% Kraft lignin, and colonies picked. This method
222 resulted in 2 further isolates, whose 16S rRNA gene was amplified by polymerase chain reaction,
223 and submitted for DNA sequencing.

224 By comparison with database 16S rRNA sequences, the identity of the isolates was
225 established, as shown in Table 1. Three of the isolates were found to be in the *Ochrobactrum* class
226 of α -proteobacteria. Our group has previously reported the isolation of two *Ochrobactrum* isolates
227 with activity for lignin oxidation from woodland soil (Taylor *et al.*, 2012). A further *Agrobacterium*
228 sp. isolate is also a member of the α -proteobacteria; there is a previous report of an *Agrobacterium*
229 isolate able to degrade Kraft lignin (Deschamps *et al.*, 1980). A *Comamonas testosteroni* isolate is a
230 member of the β -proteobacteria; there is a recent report of a *Comamonas* isolate able to degrade
231 Kraft lignin (Chen *et al.*, 2012). Two *Enterobacter* isolates are members of the γ -proteobacteria;
232 there are reports of an *Enterobacter lignolyticus* strain isolated from tropical soils amended with
233 Kraft lignin that is able to degrade lignin under anaerobic conditions (DeAngelis *et al.*, 2011a;
234 DeAngelis *et al.*, 2011b). One *Pseudomonas* isolate was found, which is also a member of the γ -
235 proteobacteria; we and others have previously reported activity of *Pseudomonas putida* for lignin
236 degradation (Ahmad *et al.*, 2010; Salvachua *et al.*, 2015). One *Microbacterium* isolate was found,

237 which is a member of the actinobacteria; our group has previously reported the isolation of three
238 *Microbacterium* strains active for lignin breakdown from woodland soil (Taylor *et al.*, 2012). Two
239 isolates were found to members of the Firmicutes phylum. The *Lysinibacillus* isolate is from the
240 bacilli class: although our group has found previously that *Bacillus subtilis* has no activity for lignin
241 oxidation (Ahmad *et al.*, 2010), there are reports of *Bacillus* and *Paenibacillus* isolates with activity
242 for Kraft lignin oxidation (Chandra *et al.*, 2008), and we have also identified here a *Paenibacillus*
243 isolate able to grow on minimal media containing Kraft lignin.

244

245 Table 1. Bacterial lignin-degrading strains isolated from municipal landfill soil via either nitrated
246 lignin spray assay (method A) or growth on minimal media containing Kraft lignin (method B).

247

248 **Activity of bacterial strains for delignification**

249 The collection of new bacterial isolates were tested for delignification activity alongside
250 four further lignin-degrading bacterial strains (two further *Microbacterium* strains, a *Rhodococcus*
251 *erythropolis* strain, and *Sphingobacterium* sp. T2) isolated from woodland soil (Taylor *et al.*, 2012),
252 and *Rhodococcus jostii* RHA1 and *Pseudomonas putida* mt-2 known to have lignin degradation
253 activity (Ahmad *et al.*, 2010), and *Pseudomonas fluorescens* Pf-5 from which a lignin-oxidising
254 Dyp1B peroxidase has been identified (Rahmanpour & Bugg, 2015). The assay involved incubation
255 of a bacterial starter culture with 1 g of chopped pine lignocellulose for 7 days. The treated solid
256 residue and untreated lignocellulose was then analysed for percentage lignin content using the
257 Klason assay (Kirk & Obst, 1988). Samples of the treated aqueous supernatant after 4 days and 7
258 days were also analysed for total phenol content using the Folin-Ciocalteu colorimetric assay
259 (Meda *et al.*, 2005). The results are shown in Table 2.

260 The results obtained from Klason assay of the treated material showed that 4 strains gave
261 20-25% delignification after 7 days (*P. putida*, *P. fluorescens*, *E. cloacae*, and *L. sphaericus*), as

262 shown in Table 2, with another 4 strains giving 15-20% delignification (*O. pectoris*, *Agrobacterium*
263 sp., *E. ludwigii*, and *Microbacterium* sp.).

264 Using the Folin-Ciocalteu assay, 10 strains gave >20% increase in phenol release after 4 or
265 7 days, with highest phenol release observed with *C. testosteroni* (74% increase) and *R.*
266 *erythropolis* (63% increase). Significant differences between phenol release was observed between
267 4- and 7-day time-points, with some strains showing maximum phenol release after 7 days, whereas
268 for other strains maximum phenol release was observed after 4 days, decreasing after 7 days. This is
269 ascribed to the high phenol degradation activity of some strains, which would start to degrade low
270 molecular weight phenols as their concentration builds up.

271

272 Table 2. Delignification and phenol release of bacterial lignin-degrading isolates, using milled pine
273 lignocellulose as substrate.

274


275 **Activity of recombinant bacterial lignin-oxidising enzymes for delignification**

276 The activity of purified recombinant bacterial lignin-oxidising enzymes for delignification and
277 phenol release from lignocellulose was also tested, comparing *P. fluorescens* Dyp1B (Rahmanpour
278 & Bugg, 2015) with *Sphingobacterium* sp. T2 manganese superoxide dismutase (Rashid *et al.*,
279 2015), and also comparing with commercially available lignin peroxidase from *Phanerochaete*
280 *chryso sporium* (Sigma-Aldrich). Doses of 0.2-1.0 mg enzyme/g lignocellulose were incubated with
281 1.0 g pine lignocellulose for 1 hr, and the residual solid assayed for lignin content using the Klason
282 method (Kirk & Obst, 1988), and the aqueous sample tested for total phenol content using the
283 Folin-Ciocalteu colorimetric assay (Meda *et al.*, 2005), as shown in Figure 1.

284 Using the Klason assay, dose-dependent delignification was observed for each enzyme, with
285 highest activities at 1 mg/g dose. At this dose 26% delignification was observed for *P. fluorescens*
286 Dyp1B, 31% for *Sphingobacterium* sp. T2 MnSOD, and 31% for *P. chryso sporium* lignin
287 peroxidase. These levels of delignification are comparable to the 20-25% decreases in lignin

288 content for bacterial strain treatment after 7 days (see Table 2), but were achieved *in vitro* in a 1 hr
289 treatment.

290 Using the Folin-Ciocalteu assay, dose-dependent increases in phenol release were observed for
291 *Sphingobacterium* sp. T2 MnSOD, up to 30% increase for the 1 mg/g dose. Only very small
292 changes were observed upon treatment with *P. fluorescens* Dyp1B, with a 10% decrease in phenol
293 content at low dose, and 8% increase at high dose, compared with the untreated lignocellulose
294 control. This behaviour might be due to the tendency of lignin-oxidising peroxidases to catalyse
295 repolymerisation as well as depolymerisation of lignin fragments (Rahmanpour *et al.*, 2017). For
296 fungal lignin peroxidase, a decrease in phenol content was also observed at low dose, perhaps for
297 the same reason, but at higher dose a dose-dependent increase in phenol release was observed, with
298 >2-fold phenol release at 1 mg/g dose.




299
300

301 Figure 1. Delignification and phenol release by *Sphingobacterium* sp. T2 MnSOD1, *P. fluorescens*
302 Dyp1B and *P. chrysosporium* lignin peroxidase.

303

304 **Enhancement of gas release from soil containing lignocellulose**


305 We then examined whether addition of a bacterial lignin degrader to soil containing
306 lignocellulose could enhance gas release. The experimental design (see Figure 2A) involved
307 samples of soil collected from a municipal solid waste (MSW) site, mixed with 1% (w/w) chopped
308 pine lignocellulose, packed in 2.5, 5, or 10 mL plastic syringes, to which sodium acetate (6.25 g/L)
309 buffer, bacteria or enzyme was added. The syringe was then sealed, and released gas was collected
310 via plastic tubing, and the volume of gas measured. Liquid run-off was collected from the bottom of
311 the syringe for analysis, and additional buffer added via airtight needle at the top of the syringe.



312
313 Figure 2. Small-scale testing of gas release from endogenous microbial population in MSW soil. A.
314 Schematic diagram of experimental set-up. B. Gas generation from MSW soil in the presence and
315 absence of 6.25 g/L NaOAc, in 2.5, 5.0, and 10 mL syringes.
316

317 First, the generation of gas from the microbial population in the MSW soil was tested
318 without addition of exogeneous bacteria. In order to stimulate methanogenic bacteria in MSW, 6.25
319 g/L sodium acetate was added to 2.5, 5, and 10 mL syringes containing MSW, and the experiment
320 left at room temperature (20-25 °C) for up to 20 days. The data in Figure 2B show that, after a lag
321 phase of 7 days, time-dependent gas production was observed in the samples supplemented with
322 sodium acetate buffer, with optimum gas production observed in the 10 mL sample.

323 The addition of exogeneous bacteria, grown on Luria-Bertani media and then suspended in
324 M9 salts (100 µL), was then tested using 10 mL MSW containing 1% (w/w) chopped pine
325 lignocellulose, but without addition of sodium acetate, over 35 days. The 6 bacterial strains showing
326 highest levels of delignification and/or phenol release (see Table 2) were tested in this experiment,
327 namely *P. putida*, *Ochrobactrum* sp., *Agrobacterium* sp., *L. sphaericus*, *C. testosteroni*, and
328 *Paenibacillus* sp.. As shown in Figure 3, after a lag phase of 20 days, enhanced gas production was
329 observed initially with *Agrobacterium* sp. and *L. sphaericus*, giving 10-fold enhancement of gas
330 production compared with a control lacking exogenous bacteria. After 25-30 days, enhancement of
331 gas production was also observed to a lesser extent with *C. testosteroni* and *Paenibacillus* sp.,
332 giving 4-fold enhancement of gas production compared to the control lacking exogenous bacteria.
333 No significant enhancement of gas production under these conditions was observed using *P. putida*
334 or *Ochrobactrum* sp.




335

336 Figure 3. Enhancement of gas production *in situ* from 10 mL MSW soil containing 1% chopped
 337 pine lignocellulose by addition of bacterial lignin degraders

338

339 An alternative two-stage treatment scenario was also tested, whereby samples of pine
 340 lignocellulose were pre-treated aerobically with six bacterial strains in minimal media for 7 days,
 341 and then the sample centrifuged, mixed with 10 mL MSW, and incubated as above without sodium
 342 acetate. In order to compare biological vs. thermochemical pretreatment, another sample was
 343 delignified thermochemically (ethanol organosolv method). Anaerobic digestion of pre-treated
 344 (biologically and thermochemically treated) pine samples resulted in generation of up to 3-fold
 345 more gas (see Figure 4) compared to untreated pine and ethanosolv lignin from pine, over a 40-day
 346 experiment. In this scenario, optimum enhancement was observed with *P. putida* and *Paenibacillus*
 347 (3-fold enhancement), followed by *C. testosteroni* and *L. sphaericus* (2.6-fold enhancement), then
 348 *Ochrobactrum* sp. and *Agrobacterium* sp. (1.4-1.7 fold enhancement), which were similar to the
 349 thermochemically treated pine (1.4-fold enhancement, compared with untreated pine). Of particular
 350 note was that *Pseudomonas putida*, which showed the highest activity in the 2-stage treatment,


351 showed no activity in the *in situ* experiment above, whereas *Agrobacterium* sp. and *L. sphaericus*
352 showed activity in both treatment scenarios



353
354 Figure 4. Enhancement of gas production from pine lignocellulose pre-treated aerobically with
355 lignin-degrading strains, followed by anaerobic digestion with MSW soil.

356
357 The first *in situ* treatment method was then tested on a larger 0.5 kg scale, using
358 commercially available organic compost in place of MSW soil, and supplementing with 10% (w/w)
359 commercial softwood bark chips in place of chopped pine lignocellulose. Test experiments using
360 *Agrobacterium* sp. confirmed that 3-fold enhanced gas generation was observed over a 30-day
361 experiment containing organic compost supplemented with softwood bark chips, compared with
362 experiments lacking softwood bark chips, or lacking additional bacteria, as shown in Figure 5A.
363 Addition of a sample of anaerobic digester extract to the organic compost was found to make no
364 significant difference to the gas yield (see Figure 5A), indicating that there is an efficient microbial
365 population for anaerobic gas production present in commercial organic compost.

366
367



368

369 Figure 5. Enhancement of gas release from 0.5 kg organic compost supplemented with 10% (w/v)
 370 softwood bark chips, to which was added bacterial culture.

371

372 Using these optimised conditions, the four bacterial isolates that showed activity in small-
 373 scale trials (*Agrobacterium* sp., *Lysinibacillus sphaericus*, *Comamonas testosteroni*, *Paenibacillus*

374 *sp.*) were tested on a 0.5 kg scale experiment over 30 days. Enhanced gas production was observed
375 with all four isolates, as shown in Figure 5B, showing 4-5 fold enhancement of gas production,
376 compared with a control experiment lacking bacteria. Greatest enhancement of gas release was
377 observed with *Lysinibacillus sphaericus* at 10-30 days. In this experiment, samples of gas at 7 day
378 intervals were analysed by gas chromatography, revealing that the methane content was 15% after 7
379 days, 14% after 14 days, and 42% after 21 days.


380

381 **Incubation of β -aryl ether lignin model compound with anaerobic consortium.**

382 Our observation that lignin degradation occurs under microanaerobic conditions is
383 surprising, since the known lignin degradation pathways are oxidative and aerobic (Bugg *et al.*,
384 2011a). Since the strains that are most effective under these conditions are facultative anaerobes, it
385 is possible that these organisms might use reductive or non-redox pathways to break down lignin. In
386 order to probe the molecular basis of the biotransformation of lignin under anaerobic conditions, a
387 sample of anaerobic consortium collected from the 0.5 kg scale experiment described above
388 (supplemented with *Agrobacterium sp.*) was incubated with β -aryl ether lignin model compound
389 guaiacylglycerol- β -guaiacyl ether in M9 minimal media under anaerobic conditions for 16 days.
390 Analysis of supernatant samples by LC/MS showed that a peak at 27.5 min corresponding to the
391 lignin model compound (MNa^+ 343) was reduced in size after 2 days, and consumed after 8 days
392 (see Supporting Information Figure S3).

393 Extracted ion chromatographic analysis of the LC-MS data showed the formation of new
394 compounds of reduced molecular mass: a new species at m/z 284.9 was observed after 2 days
395 (Supporting Information Figure S4); while several new species were observed after 8 days at m/z
396 329, 270.9, 258.9, 274.9, 244.9 and 315 (Supporting Information Figure S5). The products at m/z
397 329 and 315 indicate two successive demethylation reactions occurring on compound **1**, consistent
398 with the formation of compounds **2** and **3**, shown in Figure 6. The new product observed at m/z 275
399 is consistent with loss of formaldehyde via C-C fragmentation of the β,γ -bond to form compound **4**,

400 which could be rationalised by oxidation of the α -hydroxyl group to a ketone, followed by retro-
 401 aldol cleavage. Demethylation of **4** followed by reduction of the α -keto group to the alcohol would
 402 give compound **5**, consistent with the observed peak at m/z 285. The observed species at m/z 245 is
 403 consistent with dehydroxylation of compound **4** in the *para* position, and demethylation, to form
 404 compound **6**. Although aromatic dehydroxylation is unusual, it is preceded in the bacterial
 405 anaerobic degradation of phenol via dehydroxylation of 4-hydroxybenzoyl-CoA (Glöckler *et al.*,
 406 1989; Boll *et al.*, 2014). Hence the observed metabolites are consistent with demethylation, β,γ -
 407 fragmentation, and *para*-dehydroxylation reactions, as shown in Figure 6.



408
 409 Figure 6. Proposed degradation route of β -aryl ether lignin model compound by anaerobic
 410 consortium from MSW.

411

412 Discussion

413 We have identified 11 new bacterial lignin-degrading isolates, 7 using a nitrated lignin spray
 414 assay method previously published (Taylor *et al.*, 2012), and 4 that are able to grow on Kraft lignin
 415 as a sole carbon source. Of these strains, three *Ochrobactrum* isolates and one *Microbacterium*

416 isolate are of similar type to those isolated from woodland soil, while *Pseudomonas* strains are
417 known to have activity for lignin degradation (Ahmad *et al.*, 2010). The majority of bacterial lignin-
418 degrading strains isolated previously are in the actinobacteria or α - or γ -proteobacteria phyla (Bugg
419 *et al.*, 2011b; Tian *et al.*, 2014; Wang *et al.*, 2016). We have also isolated several new facultative
420 anaerobic lignin-degrading strains in this study: an *Agrobacterium* sp. isolate from the α -
421 proteobacteria phylum, consistent with two previous reports of lignin-degrading *Agrobacterium*
422 strains (Deschamps *et al.*, 1980, Si *et al.*, 2015); two *Enterobacter* isolates from the γ -
423 proteobacteria phylum, consistent with reports of a lignin-degrading *Enterobacter lignolyticus*
424 strain (DeAngelis *et al.*, 2011a; DeAngelis *et al.*, 2011b); a *Comamonas testosteroni* strain from the
425 β -proteobacteria, consistent with a lignin-degrading *Comamonas* isolate (Chen *et al.*, 2012); and
426 *Paenibacillus* and *Lysinibacillus sphaericus* strains in the Firmicutes phylum, consistent with
427 *Paenibacillus* strains reported to degrade industrial lignins (Chandra *et al.*, 2008; Mathews *et al.*,
428 2016). We note that the bacteria that we isolated previously from woodland soil were all aerobic
429 bacteria (Taylor *et al.*, 2012), whereas some of the bacteria isolated here are facultative anaerobes,
430 which might reflect the sampling of topsoil in the earlier study vs. subsoil in this study.

431 We have tested 20 lignin-degrading bacteria for delignification of pine lignocellulose, and
432 have observed 15-24% delignification via Klason assay after a 7-day treatment with 8 of the strains
433 (see Table 2). Phenol release was also observed with >10 strains, indicative of lignin breakdown,
434 but there is not a clear correlation between the data for phenol release and delignification in Table
435 2. We suggest that this is because some of these strains are also very efficient aromatic degraders
436 (Taylor *et al.*, 2012), hence they would rapidly consume phenolic compounds that are produced
437 from lignin breakdown, therefore a lack of phenol release is not necessarily indicative of a lack of
438 lignin breakdown. Treatment with white-rot fungi such as *Phanerochaete chrysosporium* is known
439 to improve biomass digestibility (Akin *et al.*, 1993), but bacteria such as *Streptomyces griseorubens*
440 have also been shown to improve enzymatic saccharification yields (Saritha *et al.*, 2013), and there

441 is interest in the use of microbial treatment for delignification in the context of cellulosic bioethanol
442 production (Moreno *et al.*, 2015).

443 We have also tested recombinant lignin-oxidising enzymes for delignification activity, and
444 we have observed a dose-dependent reduction in Klason lignin content using *P. fluorescens* Dyp1B
445 (Rahmanpour & Bugg, 2015) and *Sphingobacterium* sp. T2 MnSOD1 (Rashid *et al.*, 2015), as
446 shown in Figure 1. Similar levels of delignification were observed using fungal *P. chrysosporium*
447 lignin peroxidase (31.1% delignification at 1 mg/g lignocellulose), although the fungal enzyme
448 showed higher levels of phenol release (Figure 1). The loadings of enzyme required to see this level
449 of delignification (1.0 mg/g lignocellulose) are fairly high, but are comparable with doses of 0.1-
450 0.15 mg/g used in previous treatments with fungal MnP and LiP enzymes (Jayasinghe *et al.*, 2011;
451 Hettiaratchi *et al.*, 2014).

452 We have then tested whether using a lignin-degrading bacterium as an additive, enhanced
453 gas generation can be achieved during anaerobic lignocellulose breakdown. We have examined two
454 different scenarios for carrying out a bacterial treatment. Carrying out microbial treatment under
455 aerobic conditions, and then using the treated lignocellulose for anaerobic digestion, aerobic
456 degraders such as *Pseudomonas putida* show highest activity (see Figure 4), but for an *in situ*
457 bacterial treatment under microanaerobic conditions, facultative anaerobes such as *Agrobacterium*
458 sp. and *Lysinibacillus sphaericus* show highest activity (see Figures 3 and 6). For a treatment of
459 landfill soil in the environment, an *in situ* treatment would have the advantage of not needed to
460 disturb the landfill soil, hence the ability of the bacteria used in Figure 6 to operate under
461 microanaerobic conditions could be very useful.

462 It is surprising that lignin degradation occurs under microanaerobic conditions, since the
463 known lignin degradation pathways are oxidative and aerobic (Bugg *et al.*, 2011a), hence we have
464 studied the degradation of a β -aryl ether lignin model compound by an anaerobic consortium. The
465 observed metabolites are consistent with demethylation, β,γ -fragmentation, and *para*-
466 dehydroxylation reactions taking place (see Figure 6). Anoxic oxidation of alkyl sidechains is

467 preceded in anaerobic aromatic-degrading bacteria, via either flavocytochrome c hydroxylases or
468 molybdenum-dependent hydroxylases (Boll *et al.*, 2014), hence such enzymes might be responsible
469 for the demethylation reactions observed here, or a non-redox demethylation reaction could be
470 taking place, similar to the tetrahydrofolate-dependent demethylase LigM from *Sphingobium* SYK-
471 6 (Rosini *et al.*, 2016). It seems plausible that the 1-carbon unit released via anaerobic
472 demethylation may ultimately be converted to methane gas by the anaerobic consortium.
473 Interestingly, the majority of bacteria known to carry out anaerobic aromatic degradation are in the
474 α -proteobacteria and β -proteobacteria (Boll *et al.*, 2014), in which phyla the *Agrobacterium* and
475 *Comamonas testosteroni* strains identified here respectively belong. A strain of *Dysgonomonas* sp.
476 from the Bacteroides phylum has recently been reported to carry out degradation of Kraft lignin
477 under anaerobic conditions (Duan *et al.*, 2016), though the biochemical pathways used by this
478 organism are not known. The ability to delignify lignocellulose *in situ* offers possible applications
479 for treatment of landfill waste, and for delignification of biomass or lignocellulosic waste generated
480 from agriculture, pulp/paper manufacture, or industries utilising lignocellulosic feedstocks.

481

482 **Acknowledgements.** This work was supported by NERC research grant NE/L013983/1, as part of
483 the Resource Recovery from Waste initiative. The authors would like to thank Dr Talib Mahdi
484 (Cardiff School of Engineering, Cardiff University) and VertaseFLI Limited (Bristol, BS20 6PN,
485 UK) for collection of soil samples, Dr Orkun Soyer (School of Life Sciences, University of
486 Warwick) for the gift of a sample of anaerobic digester extract, Dr Hendrik Schäfer (School of Life
487 Sciences, University of Warwick) for assistance with gas analysis via gas chromatography, and
488 Andrew Rankin (University of Warwick) for preliminary work in the development of
489 delignification methods.

490 **Conflict of Interest.** The authors declare no conflict of interest in the execution and submission of
491 this work.

492

493 **References**

494 Ahmad, M., Taylor, C. R., Pink, D., Burton, K., Eastwood, D., Bending, G. R. and Bugg, T. D. H.
495 (2010) Development of novel assays for lignin degradation: comparative analysis of bacterial and
496 fungal lignin degraders. *Mol. Biosystems* **6**, 815-821.

497 Ahmad, M., Roberts, J.N., Hardiman, E.M., Singh, R., Eltis, L.D. and Bugg, T.D.H. (2011)
498 Identification of DypB from *Rhodococcus jostii* RHA1 as a lignin peroxidase. *Biochemistry* **50**,
499 5096-5107.

500 Akin, D.E., Sethuraman, A., Morrison, W.H., Martin, S.A. and Eriksson, K.E. (1993) Microbial
501 delignification with white rot fungi improves forage digestibility. *Appl. Environ. Microbiol.* **59**,
502 4274-4282.

503 Boll, M., Löffler, C., Morris, B.E.L. and Kung, J.W. (2014) Anaerobic degradation of homocyclic
504 compounds by arylcarboxyl-coenzyme A esters: organisms, strategies, and key enzymes. *Environ.*
505 *Microbiol.* **16**, 612-627.

506 Brown, M.E., Barros, T. and Chang, M.C.Y. (2012) Identification and characterization of a
507 multifunctional dye peroxidase from a lignin-reactive bacterium. *ACS Chem. Biol.* **7**, 2074-2081.

508 Bugg, T.D.H., Ahmad, M., Hardiman, E.M. and Rahmanpour, R. (2011a) Pathways for degradation
509 of lignin in bacteria and fungi. *Nat. Prod. Reports* **28**, 1883-1896.

510 Bugg, T.D.H., Ahmad, M., Hardiman, E.M. and Singh, R. (2011b) The emerging role for bacteria
511 in lignin degradation and bio-product formation. *Curr. Opin. Biotech.* **22**, 394-400.

512 Chandra, R., Singh, S., Krishna Reddy, M.M., Patel, D.K., Purohit, A.J. and Kapley, A. (2008)
513 Isolation and characterisation bacterial strains *Paenibacillus* sp. and *Bacillus* sp. for kraft lignin
514 decolorization from pulp paper mill waste. *J. Gen. Appl. Microbiol.* **54**, 399-407.

515 Chen, Y.H., Chai, L.Y., Zhu, Y.H., Yang, Z.H., Zheng, Y. and Zhang, H. (2012) Biodegradation of
516 kraft lignin by a bacterial strain *Comamonas* sp. B-9 isolation from eroded bamboo slips. *J. Appl.*
517 *Microbiol.* **112**, 900-906.

518 DeAngelis, K.M., Allgaier, M., Chavarria, Y., Fortney, J.L., Hugenholtz, P., Simmons, B., Sublette,
519 K., Sliver, W.L. and Hazen, T.C. (2011a) Characterization of trapped lignin-degrading microbes in
520 tropical forest soil. *PLoS ONE* **6**, e19306.

521 DeAngelis, K.M., D’Haeseleer, P., Chivian, D., Fortney, J.L., Khudyakov, J., Simmons, B., Woo,
522 H., Arkin, A.P., Davenport, K.W., Goodwin, L., Chen, A., Ivanova, N., Kyrpides, N.C.,
523 Mavromatis, K., Woyke, T. and Hazen, T.C. (2011b) Complete genome sequence of *Enterobacter*
524 *lignolyticus* SCF1. *Stand. Genomic Sci.* **5**, 69-85.

525 Deschamps, A.M., Mahoudeau, G. and Lebault, J.M. (1980) Fast degradation of Kraft lignin by
526 bacteria. *Eur. J. Appl. Microbiol. Biotechnol.* **9**, 45-51.

527 Duan, J., Liang, J., Wang, Y., Du, W. and Wang, D. (2016) Kraft lignin degradation by
528 *Dysgonomonas* sp. WJDL-Y1, a new anaerobic bacterial strain isolated from sludge of a pulp and
529 paper mill. *J. Microbiol. Biotechnol.* **26**, 1765-1773.

530 Feng, C-L., Zeng, G-M., Huang, D-L., Hu, S., Zhao, M-H., Lai, C., Huang, C., Wei, Z. and Li, N-J.
531 (2011) Effect of lignolytic enzymes on lignin degradation and carbon utilization during
532 lignocellulosic waste composting. *Process Biochem.* **46**, 1515-1520.

533 Glöckler, R., Tschsch, A. and Fuchs, G. (1989) Reductive dehydroxylation of 4-hydroxybenzoyl-
534 CoA to benzoyl-CoA in a denitrifying, phenol-degrading *Pseudomonas* species. *FEBS Lett.* **251**,
535 237-240.

536 Hettiaratchi, J.P.A., Jayasinghe, P.A., Bartholameuz, E.M. and Kumar, S. (2014) Waste degradation
537 and gas production with enzymatic enhancement in anaerobic and aerobic landfill bioreactors.
538 *Biores. Technol.* **159**, 433-436.

539 Jayasinghe, P.A., Hettiaratchi, J.P.A., Mehrotra, A.K. and Kumar, S. (2011) Effect of enzyme
540 additions on methane production and lignin degradation of landfilled sample of municipal solid
541 waste. *Biores. Technol.* **102**, 4633-4637.

542 Jones, P.T., Geysen, D., Tielemans, Y., Van Passel, S., Potikes, Y., Blanpain, B., Quaghebeur, M.
543 and Hoekstra, N. (2013) Enhanced landfill mining in view of multiple resource recovery: a critical
544 review. *J. Cleaner Production* **55**, 45-55.

545 Kirk, T.K. and Obst, J.R. (1988) Lignin determination. *Methods Enzymol.* **161**, 87–101.

546 Mali S.T., Khare K.C. and Biradar A.H. (2012) Enhancement of methane production and bio-
547 stabilisation of municipal solid waste in anaerobic bioreactor landfill. *Biores. Technol.*, **110**, 10-17.

548 Mathews, S.L., Grunden, A.M., and Pawlak, J. (2016) Degradation of lignocellulose and lignin by
549 *Paenibacillus glucanolyticus*. *Int. Biodeter, Biodegr.* **110**, 79-86.

550 Meda, A., Lamien, C.E., Romito, M., Millogo, J. and Nacoulma, O.G. (2005) Determination of the
551 total phenolic, flavonoid and proline contents in Burkina Fasan honey, as well as their radical
552 scavenging activity. *Food Chemistry* **91**, 571–577.

553 Moreno, A.D., Ibarra, D., Alvira, P., Tomas-Pejo, E. and Ballesteros, M. (2015) A review of
554 biological delignification and detoxification methods for lignocellulosic bioethanol production.
555 *Crit. Rev. Biotechnol.* **35**, 342-354.

556 Ni, Z., Liu, J., Girotto, F., Cossu, R. and Qi, G. (2016) Targeted modification of organic
557 components of municipal solid waste by short-term pre-aeration and its enhancement on anaerobic
558 degradation in simulated landfill bioreactors. *Biores. Technol.* **216**, 250-259.

559 Rahmanpour R. and Bugg, T.D.H. (2015) Characterisation of Dyp-type peroxidases from
560 *Pseudomonas fluorescens* Pf-5: oxidation of Mn(II) and polymeric lignin by Dyp1B. *Arch.*
561 *Biochem. Biophys.* **574**, 93-98.

562 Rahmanpour, R., King, L.D.W. and Bugg, T.D.H. (2017) Identification of an extracellular bacterial
563 flavoenzyme that can prevent re-polymerisation of lignin fragments. *Biochem. Biophys. Res.*
564 *Commun.* **482**, 57-61.

565 Rashid, G.M.M., Taylor, C.R., Liu, Y., Zhang, X., Rea, D., Fülöp, V. and Bugg, T.D.H. (2015)
566 Identification of manganese superoxide dismutase from *Sphingobacterium* sp. T2 as a novel
567 bacterial enzyme for lignin oxidation. *ACS Chem. Biol.* **10**, 2286-2294.

568 Rosini, E., D-Arrigo, P. and Pollegioni, L. (2016) Demethylation of vanillic acid by recombinant
569 LigM in a one-pot cofactor regeneration system. *Catalysis Sci. Technol.* **6**, 7729-7737.

570 Salvachua, D., Karp, E.M., Nimlos, C.T., Vardon, D.R. and Beckham, G.T. (2015) Towards lignin
571 consolidated bioprocessing: simultaneous lignin depolymerization and product generation by
572 bacteria. *Green Chemistry* **17**, 4951-4967.

573 Sanchez, C. (2009) Lignocellulosic residues: biodegradation and bioconversion by fungi.
574 *Biotechnol. Adv.* **27**, 185-194.

575 Saritha, M., Arora, A., Singh, S. and Nain, L. (2013) *Streptomyces griseorubens* mediated
576 delignification of paddy straw for improved enzymatic saccharification yields. *Biores. Technol.*
577 **135**, 12-17.

578 Si, W., Wu, Z-W., Wang, L-L., Yang, M-M. and Zhao, X. (2015) Enzymological characterization
579 of Atm, the first laccase from *Agrobacterium* sp. S5-1, with the ability to enhance *in vitro*
580 digestibility of maize straw. *PlosOne* **10**, e0128204.

581 Taylor, C.R., Hardiman, E.M., Ahmad, M., Sainsbury, P.D., Norris, P.R. and Bugg, T.D.H. (2012)
582 Isolation of bacterial strains able to metabolize lignin from screening of environmental samples. *J.*
583 *Appl. Microbiol.* **113**, 521-530.

584 Tian, J-H., Purcher, A-M., Bouchez, T., Gelhaye, E. and Peu, P. (2014) Occurrence of lignin
585 degradation genotypes and phenotypes among prokaryotes. *Appl. Microbiol. Biotechnol.* **98**, 9527-
586 9544.

587 Tonini, D., Martinez-Sanchez, V. and Astrup, T.F. (2013) Material resources, energy, and nutrient
588 recovery from waste: are waste refineries the solution for the future? *Environ. Sci. Technol.* **47**,
589 8962-8969.

590 Wang, L., Nie, Y., Tang, Y-Q., Song, X-M., Cao, K., Sun, L-Z., Wang, Z-J., Wu, X-L. (2016)
591 Diverse bacteria with lignin degrading potentials isolated from two ranks of coal. *Frontiers*
592 *Microbiol.* **7**, 1428.

593 Yazdani, R., Mostafid, M.E., Han, B., Imhoff, P.T., Chiu, P., Augenstein, D., Kayhanian, M. and
594 Tchobanoglous, G. (2010) Quantifying factors limiting aerobic degradation during aerobic
595 bioreactor landfilling. *Environ. Sci. Technol.* **44**, 6215-6220.

596

597

598 Table 1. Bacterial lignin-degrading strains isolated from municipal landfill soil via either nitrated
 599 lignin spray assay (method A) or growth on minimal media containing Kraft lignin (method B).
 600 Strain identification by sequence alignment of 16S rRNA gene sequence to Genbank sequence
 601 database.

Isolation method	Growth temp (°C)	Highest identity sequence match from 16S rRNA sequence	GenBank accession	Sequence identity	Bacterial phylum
A	30	<i>Pseudomonas</i> sp.	HM 219617	98%	γ -Proteobacteria
A	30	<i>Microbacterium oxydans</i>	JF 730219	99%	Actinobacteria
A	30	<i>Ochrobactrum pituitosum</i>	NR 115043	99%	α -Proteobacteria
A	30	<i>Comamonas testosteroni</i>	KJ 806363	96%	β -Proteobacteria
A	30	<i>Enterobacter ludwigii</i>	GQ 284566	99%	γ -Proteobacteria
A	30	<i>Enterobacter cloacae</i>	KF 017288	98%	γ -Proteobacteria
B	30	<i>Ochrobactrum</i> sp.	KF 737375	99%	α -Proteobacteria
B	30	<i>Ochrobactrum pecoris</i>	NR 117053	99%	α -Proteobacteria
B	30	<i>Agrobacterium</i> sp.	JX 872342	99%	α -Proteobacteria
B	30	<i>Paenibacillus</i> sp.	FR849917.1	99%	Firmicutes
A	45	<i>Lysinibacillus sphaericus</i>	HQ 259956	99%	Firmicutes

602

603

604 Table 2. Delignification and phenol release of bacterial lignin-degrading isolates, using milled pine lignocellulose as substrate. Lignin content
 605 measured by Klason assay (Kirk & Obst, 1988), and phenol release measured by Folin-Ciocalteau assay (Meda et al., 2005), as described in Materials
 606 and Methods section. NT, Not tested.

Bacterial strain	Phylum	Strain reference	% lignin decrease in 7 days	% phenol increase in 4 days	% phenol increase in 7 days
<i>Ochrobactrum sp.</i>	α -Proteobacteria	This study	12.9	26.8	6.7
<i>Ochrobactrum pectoris</i>	α -Proteobacteria	This study	17.4	NT	18.0
<i>Ochrobactrum pituitosum</i>	α -Proteobacteria	This study	5.6	NT	18.5
<i>Agrobacterium sp.</i>	α -Proteobacteria	This study	16.1	22.8	22.7
<i>Comamonas testosteroni</i>	β -Proteobacteria	This study	9.8	17.6	74.6
<i>Pseudomonas putida mt-2</i>	γ -Proteobacteria	Ahmad et al, 2010	21.2	18.8	3.0
<i>Pseudomonas fluorescens Pf-5</i>	γ -Proteobacteria	Rahmanpour & Bugg, 2015	20.6	NT	0.6
<i>Enterobacter ludwigii</i>	γ -Proteobacteria	This study	17.3	NT	20.0
<i>Enterobacter cloacae</i>	γ -Proteobacteria	This study	22.7	9.7	14.1
<i>Sphingobacterium sp. T2</i>	Bacteroides	Taylor et al, 2012	8.5	13.2	43.0
<i>Microbacterium phyllosphaerae</i>	Actinobacteria	Taylor et al, 2012	7.8	16.3	31.3
<i>Microbacterium sp.</i>	Actinobacteria	Taylor et al, 2012	17.6	NT	31.3
<i>Microbacterium oxydans</i>	Actinobacteria	This study	4.8	NT	1.6
<i>Rhodococcus jostii RHA1</i>	Actinobacteria	Ahmad et al, 2010	9.5	NT	38.7
<i>Rhodococcus erythropolis</i>	Actinobacteria	Taylor et al, 2012	7.3	39.1	63.7
<i>Paenibacillus sp.</i>	Firmicutes	This study	1.0	21.0	1.8
<i>Lysinibacillus sphaericus</i>	Firmicutes	This study	24.0	19.1	10.0

607

--	--	--	--	--	--

608 **Figure Legends.**

609 Figure 1. Delignification (panels A-C) and phenol release (panels D-F) by *Sphingobacterium* sp. T2
610 MnSOD1 (panels A,D), *P. fluorescens* Dyp1B (panels B,E) and *P. chrysosporium* lignin peroxidase
611 (panels C,F), at ratios of 0.2-1.0 mg enzyme/g lignocellulose, using milled pine lignocellulose as
612 substrate. Lignin content measured by Klason assay, and phenol release measured by Folin-
613 Ciocalteau assay, as described in Materials and Methods section.

614
615 Figure 2. Small-scale testing of gas release from endogenous microbial population in MSW soil. A.
616 Schematic diagram of experimental set-up. B. Gas generation from MSW soil in the presence of
617 6.25 g/L NaOAc, in 2.5 mL (purple crosses), 5.0 mL (blue diamonds), and 10 mL (red squares)
618 syringes; control experiment in absence of NaOAc (green triangles).

619
620 Figure 3. Enhancement of gas production *in situ* from 10 mL MSW soil containing 1% chopped
621 pine lignocellulose by addition of bacterial lignin degraders. A (green squares), *Ochrobactrum* sp.;
622 B (orange diamonds), *Lysinibacillus sphaericus*; C (dark blue circles), *Comamonas testosteroni*; D
623 (red crosses), *Paenibacillus* sp.; E (purple diamonds), *Pseudomonas putida*; F (yellow/blue circles),
624 *Ochrobactrum* sp.; G (blue circles), control (no bacteria added).

625
626 Figure 4. Enhancement of gas production from pine lignocellulose pre-treated aerobically with
627 lignin-degrading strains, followed by anaerobic digestion with MSW soil. A (purple crosses), pine
628 treated with *Agrobacterium* sp.; B (orange circles), pine treated with *Paenibacillus* sp.; C (red dots),
629 pine treated with *Comamonas testosteroni*; D (pale blue crosses), pine treated with *Lysinibacillus*
630 *sphaericus*; E (light green triangles), pine treated with *Agrobacterium* sp.; F (green lines), pine
631 treated with *Ochrobactrum* sp.; G (dark green triangles), pine delignified by thermochemical
632 treatment; H (blue diamonds), untreated pine incubated with MSW soil; J (red squares), organosolv
633 lignin incubated with MSW soil.

634

635 Figure 5. Enhancement of gas release from 0.5 kg organic compost supplemented with 10% (w/v)
636 softwood bark chips, supplemented with bacterial culture. A. Testing of gas production with
637 *Agrobacterium sp.* (1, green triangles), with added anaerobic digest extract (2, purple crosses),
638 versus control incubations lacking wood chips (3, red squares) and compost only (4, blue
639 diamonds). B. Testing of *Lysinibacillus sphaericus* (5, green triangles), *Comamonas testosteroni* (6,
640 purple crosses), *Paenibacillus sp.* (7, red squares), or *Agrobacterium sp.* (8, blue circles) over 30
641 days, versus compost only control (9, blue crosses).

642

643 Figure 6. Proposed degradation route of β -aryl ether lignin model compound by anaerobic
644 consortium from MSW.

645 **Supporting Information.**

646 Figure S1-S2. Apparatus for small-scale (S1) and 500 mL scale (S2) methane generation
647 experiments

648 Figure S3-S5. LC-MS data for analysis of incubation of β -aryl ether lignin model compound with
649 anaerobic consortium. S3, Disappearance of m/z 343 species; S2, selected ion chromatograms after
650 2 days incubation; S3, selected ion chromatograms after 8 days incubation.

651