

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/103554/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Brennan, Patrick J., Cheng, Tan-Yun, Pellicci, Daniel G., Watts, Gerald F. M., Veerapen, Natacha, Young, David C., Rossjohn, Jamie, Besra, Gurdyal S., Godfrey, Dale I., Brenner, Michael B. and Moody, D. Branch 2017. Structural determination of lipid antigens captured at the CD1d-T-cell receptor interface. Proceedings of the National Academy of Sciences of the United States of America 114 (31) , pp. 8348-8353. 10.1073/pnas.1705882114 file

Publishers page: <http://dx.doi.org/10.1073/pnas.1705882114>  
<<http://dx.doi.org/10.1073/pnas.1705882114>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# Structural determination of lipid antigens captured at the CD1d-T cell receptor interface

Patrick J. Brennan<sup>a,1,3</sup>, Tan-Yun Cheng<sup>a,1</sup>, Daniel G. Pellicci<sup>b,c,1</sup>, Gerald F. Watts<sup>a</sup>, Natacha Veerapen<sup>d</sup>, David C. Young<sup>a</sup>, Jamie Rossjohn<sup>e,f,g</sup>, Gurdyal S. Besra<sup>d</sup>, Dale I. Godfrey<sup>b,c,2</sup>, Michael B. Brenner<sup>a,2,3</sup>, D. Branch Moody<sup>a,2,3</sup>

<sup>a</sup>Department of Medicine, Division of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA. <sup>b</sup>Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Parkville, Australia. <sup>c</sup>ARC Centre of Excellence in Advanced Molecular Imaging, University of Melbourne, Parkville, Australia. <sup>d</sup>School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK. <sup>e</sup>Infection and Immunity Program and Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute, Monash University, Clayton, Victoria 3800, Australia <sup>f</sup>Institute of Infection and Immunity, Cardiff University School of Medicine, Heath Park, Cardiff CF14 4XN, UK <sup>g</sup>ARC Centre of Excellence in Advanced Molecular Imaging, Monash University, Clayton, Australia. <sup>1</sup>These authors contributed equally to this work. <sup>2</sup>These authors jointly directed this work. <sup>3</sup>To whom correspondence should be addressed.

Submitted to Proceedings of the National Academy of Sciences of the United States of America

**Glycolipid antigens recognized by  $\alpha\beta$  T cell receptors (TCRs) drive the activation of invariant natural killer T (iNKT) cells, a specialized subset of innate T lymphocytes. Glycolipids with  $\alpha$ -linked anomeric carbohydrates have been identified as potent microbial lipid antigens for iNKT cells, and their unusual  $\alpha$ -anomeric linkage has been thought to define a 'foreign' lipid antigen motif. Yet mammals use endogenous lipids to select iNKT cells, and there is compelling evidence for iNKT cell responses in various types of sterile inflammation. The nature of endogenous or environmental lipid antigens encountered by iNKT cells is not well defined. Here, we sought to identify lipid antigens in cow's milk, a prominent part of the human diet. We developed a method to directly capture lipid antigens within CD1d-lipid-TCR complexes, while excluding CD1d-bound to non-antigenic lipids, followed by direct biochemical analysis of the lipid antigens trapped at the TCR-CD1d interface. The specific antigens captured by this 'TCR trap' method were identified as  $\alpha$ -linked monohexosylceramides by mass spectrometry fragmentation patterns that distinguished  $\alpha$ - from  $\beta$ -anomeric monohexosylceramides. These data provide direct biochemical evidence for  $\alpha$ -linked lipid antigens from a common dietary source.**

NKT | iNKT | Lipid antigen | CD1d

## Introduction

Adaptive mammalian T lymphocytes are well known for their use of combinatorial diversity to generate a broad repertoire of antigen receptors, a process that evolved to recognize newly-encountered peptide epitopes from pathogens. However, certain subsets of human T cells, such as invariant natural killer T (iNKT) cells, mucosal-associated invariant T (MAIT) cells, and germline-encoded mycolyl-reactive (GEM) T cells reliably generate T cell receptors (TCRs) that are conserved among genetically diverse humans, and in some cases, across species (1). Recent data indicate that T cells expressing such invariant TCRs comprise a significant fraction of all human T cells, estimated in the range of 10 to 20 percent (2). Further, these conserved T cell subsets are not restricted by major histocompatibility (MHC)-encoded antigen presenting molecules, but instead can recognize different chemical classes of antigens presented by non-polymorphic antigen presenting molecules including CD1, MR1 or HLA-E (3). One of these conserved T cell populations, iNKT cells, recognizes specific glycolipids presented by CD1d molecules. Following activation, iNKT cells respond rapidly with vigorous cytokine production that can profoundly shape immune responses (4).

Although a glycolipid antigen for iNKT cells was identified from a marine sponge more than 20 years ago as part of a chemical library screen (5), the origins and chemical identities of physiologically-relevant antigens that drive selection and activation of iNKT cells remain controversial. Natural antigenic glycol-

ipids have been detected from various foreign sources, including non-pathogenic  $\alpha$ -proteobacteria (6-8), spirochetes (9), and other microbes such as *Streptococcus pneumoniae* (10), and *Bacteroides fragilis* (11, 12). These known microbial lipid antigens for iNKT cells share a key biochemical feature, namely an  $\alpha$ -anomeric linkage of one hexose sugar to the lipid backbone. Mammals abundantly produce  $\beta$ -linked hexosylceramides, glucosylceramide and galactosylceramide, but were not previously thought to produce  $\alpha$ -linked hexosylceramides. Therefore, a major principle of antigen recognition by iNKT cells has been that self-glycolipids are  $\beta$ -linked and not strongly recognized, while pathogens produce lipids that are recognized as foreign based on their  $\alpha$ -anomeric linkage (5). However, pathogen-derived lipids presumably do not explain the TCR- and CD1d-dependent activation of iNKT cells during positive selection in the thymus, or in settings such as viral infection, cancer, or sterile inflammation (4). Endogenous lipids have long been proposed to activate iNKT cells in these settings.


It is possible that the longstanding view that mammalian monohexosylceramides are always  $\beta$ -linked is incorrect. Kain and colleagues showed that antibodies raised against  $\alpha$ -linked glycosphingolipids could stain mammalian tissues (13). We recently fractionated mammalian milk-derived lipids, finding that antigenic substances were enriched in the hexosylceramide-containing fraction, and that antigenicity was not abrogated by  $\beta$ -specific glycosidases or chromatographic depletion of  $\beta$ -

## Significance

**Invariant natural killer T (iNKT) cells are activated quickly and play a key role in the control of many microbial infections via their ability to rapidly secrete cytokines and chemokines that enhance many immune responses. Microbial glycolipid antigens that activate iNKT cells have been identified as  $\alpha$ -anomeric linked glycolipids. However, the nature of the endogenous lipid antigens that are important contributors to the biology of iNKT cells has been unclear. In this study, antigenic lipids from cow's milk were isolated using a T cell receptor trap method, and their stereochemical structures were determined to be hexosylceramides with  $\alpha$ -linked hexose headgroups, thus identifying dietary lipid antigens for iNKT cells.**

## Reserved for Publication Footnotes

137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204



**Fig. 1.** iNKT cell activation by milk hexosylceramides. (A) A cow's milk glucosylceramide-fraction was digested with recombinant lysosomal glucosidase to reduce  $\beta$ -glucosylceramide. The remaining material detected at the monohexosylceramide retention time was purified by preparative TLC and further resolved to upper and lower fractions. (B) TLC fractions were tested for activity by co-culture with the DN32 iNKT cell hybridoma and CD1d-transfected RAW-264.7 cells. (C) CD1d monomers loaded with the indicated lipids were assembled into tetramers that were used to stain BW58 cells transfected with an iNKT cell TCR or with a diverse NKT cell TCR that does not recognize  $\alpha$ -galactosylceramide. Numbers indicate geometric mean fluorescence values. Data are representative of two independent experiments performed with separate starting material.


glucosylceramides (14). Both studies point to the possibility that  $\alpha$ -linked lipids are produced in mammalian tissues. However, if they exist, they are likely scarce compounds and they have not been isolated in sufficient yield to confirm  $\alpha$ -anomeric structures through direct biochemical analysis. Given the importance of the 'alpha-linked self-lipid hypothesis' to iNKT cell biology, we sought to identify mammal-derived antigenic lipids.

Here, we used soluble iNKT TCR and soluble CD1d to capture the subset of lipids from cow's milk that form stable TCR-lipid-CD1d complex through direct analysis of TCR-lipid-CD1d complexes. Combining this TCR 'trapping' of CD1d-lipid antigen complexes with nanoelectrospray ionization (ESI) mass spectrometry (MS) and liquid chromatography-mass spectrometry (LC-MS), we could detect monohexosylceramides from cow's milk as lipid antigens. Using MS-based methods, we determined that the captured lipid antigens were  $\alpha$ -linked monohexosylceramides. This study identifies the biochemical structures of natural cow's milk-derived lipid antigens as  $\alpha$ -anomeric hexosylceramides, supporting the hypothesis that  $\alpha$ -linked lipids are iNKT cell antigens even in the absence of pathogens.

**Results**

**Pre-purification of milk lipid antigens for iNKT cells.** We previously showed that the glucosylceramide-containing fraction of cow's milk contains antigenic lipids for iNKT cells (14, 15). To further purify and identify the milk-derived lipid antigens, we digested a milk hexosylceramide lipid fraction with  $\beta$ -glucosidase to destroy the  $\beta$ -linked glucosylceramides. Thin layer chromatography (TLC) showed that this treatment markedly reduced the total mass of glycolipids co-migrating with a glucosylceramide


205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272




**Fig. 2.** TCR trap assay. (A) Synthetic  $\alpha$ -glucosylceramide was loaded in recombinant CD1d molecules, mixed with recombinant iNKT cell TCRs (2C12), and run over a size-exclusion chromatography column. (B) Lipids were eluted from the early fraction excluded from the column and analyzed by linear ion trap MS. A dominant ion corresponding to the  $\alpha$ -glucosylceramide standard was recovered and few other ions were detected. (C) The activity-enriched milk monohexosylceramide fraction was loaded in CD1d and treated as in (Fig. 2A). (D) The early fraction, CD1d alone, or TCR alone, were resolved on an SDS-PAGE gel and total proteins stained, yielding bands corresponding to TCR, CD1d, and  $\beta$ 2M. Data are representative of two independent experiments performed with separate starting material.

standard, yet did not reduce the potency of iNKT cell activation (Fig. 1A). This result could be explained if the starting material was predominantly composed of non-antigenic  $\beta$ -linked glucosylceramides, and antigenic  $\alpha$ -linked hexosylceramides were present in milk that survived the enzyme treatment. To address this possibility, we used a TLC separation method that resolved monohexosylceramide  $\alpha$ - and  $\beta$ - anomers (15), yielding two bands. Only the lower band, which co-migrated with an  $\alpha$ -glucosylceramide standard, stimulated an iNKT cell response (Fig. 1B), suggesting that  $\alpha$ -linked hexosylceramides could be present. However, this interpretation is contrary to the prevailing view that mammalian products contain only  $\beta$ -linked hexosylceramides. The lower band that activated iNKT cells (Fig. 1A,B) which we refer to as the "activity-enriched" fraction, represented a very small fraction of total milk hexosylceramides, which provided an obstacle to formal biochemical identification of the antigen. Thus, we considered new approaches to antigen discovery.

**Milk lipids mediate CD1d-TCR binding.** Nearly all prior approaches to antigen discovery, including that described above (Fig. 1A), have relied on a correlation between two assays that separately measure T cell activation and biochemical content. The final antigen identification relies on the premise that the most readily detectable molecules are also the T cell stimulants. By definition, antigenic lipids ligate TCRs to antigen-presenting molecules with higher affinity than non-antigenic lipids, so we reasoned that a recombinant iNKT TCR with relatively high affinity for CD1d- $\alpha$ -galactosylceramide complexes, known as 2C12 (16), might selectively trap those CD1d complexes carrying milk-derived antigens, while leaving behind CD1d molecules that were bound to non-antigenic lipids. Such a TCR trap approach was recently used to identify CD1a autoantigens (17). We hypothesized that this approach might be feasible for CD1d and milk-derived antigens, since partially-enriched milk fractions could mediate the physical binding of CD1d tetramers to cells expressing an iNKT cell Va14Ja18 V $\beta$ 8.2J $\beta$ 2.1 TCR, but not to cells expressing a sulfatide-reactive diverse NKT cell Va1Ja26 V $\beta$ 16J $\beta$ 2.1 TCR (18, 19). Staining of iNKT TCR-expressing cells with CD1d tetramers



**Fig. 3.** Linear ion trap MS analysis of CD1-lipid-TCR complexes. (A) Lipids extracted from the TCR trap assay fractions indicated Fig. 2C were analyzed by linear ion trap MS. Prominent ions from the late fraction were identified as (B) phosphatidylcholine, (C) phosphatidylethanolamine, (D) sphingomyelin, and (E) type 1 monohexosylceramide. (F) All major ions in the early fraction were an alkane series, and collision of 6 of the most abundant ions generated type 2 monohexosylceramide patterns. (G) Lipids were extracted from the early fraction, quantified, and co-cultured with the DN32 iNKT cell hybridoma and CD1d-transfected RAW-264.7 cells. Experiments were performed on lipids extracted two independent TCR trap assays.




**Fig. 4.** HPLC-MS analysis with Q-ToF MS detection. Lipids extracted from TCR trap assay fractions were analyzed by Q-ToF HPLC-MS experiments to provide high mass accuracy confirmation of the structures deduced in Fig. 3, measure lipid binding to TCR and to measure retention time in comparison with standards. HPLC tracings are shown for selected targeted ions based on predicted masses for the structures identified in Fig. 3. Experiments were performed on lipids extracted two independent TCR trap assays.

loaded with activity-enriched milk lipids was stronger than staining with unloaded CD1d tetramers, but weaker than tetramers generated with known potent lipid antigens  $\alpha$ -galactosylceramide and  $\alpha$ -glucosylceramide (Fig. 1C). Thus, cow's milk contained compounds capable of mediating CD1d-TCR binding.

**TCR Trap Assay.** To develop a cell-free assay to detect lipid antigens, we tested the TCR trap system by loading recombinant soluble CD1d with synthetic  $\alpha$ -glucosylceramide, and then mixing this antigen-loaded CD1d with recombinant soluble iNKT cell TCR monomers. The subset of CD1d proteins carrying  $\alpha$ -glucosylceramide was expected to bind TCR heterodimers leading to the formation of a higher molecular weight ternary complex, which could be purified by size exclusion gel chromatography. A mixture of  $\alpha$ -glucosylceramide-treated CD1d and TCRs showed a large early fraction that eluted prior to separate CD1d and TCR monomers (Fig. 2A), consistent with a larger protein complex. After eluting lipids from this fraction using a mixture

of chloroform and methanol, positive mode nano-electrospray MS identified a prominent ion at  $m/z$  832.7, corresponding to  $\alpha$ -glucosylceramide ( $[M+Na]^+$ ,  $M=C_{48}H_{91}NO_8$ ), as well as its isotopes. The lack of other ions present above background levels indicated that lipids from culture media or those potentially carried by the outer surface of CD1d or TCRs did not substantially confound the analysis (Fig. 2B). Thus,  $\alpha$ -glucosylceramide performed its expected role in ligating CD1d to TCR, and direct analysis of CD1d-lipid-TCR complexes was sufficient to identify the ligating molecule without interference by other substances adherent to the lipid-protein complex.

Next, to identify unknown lipids, we used this TCR trap assay with CD1d molecules that had been loaded with activity-enriched cow's milk monohexosylceramides (Fig. 1A,B). A small but detectable peak was observed in the early size exclusion fractions and this showed no overlap with CD1d or TCR monomer fractions (Fig. 2C). Coomassie staining of the early fraction revealed



**Fig. 5.** Linear ion trap MS distinguishes  $\alpha$ - from  $\beta$ -hexosylceramides. (A) Lipid standards were synthesized based on the structures of the type 2 monohexosylceramides and tested in co-culture with the DN32 iNKT cell hybridoma and CD1d-transfected RAW-264.7 cells. (B) MS-MS of sodiated adducts demonstrated characteristic fragmentation patterns for  $\alpha$ - and  $\beta$ -glucosylceramide synthetic standards when studied over a range of collision energy settings. (C) Linear ion trap MS-MS was performed on the cow's milk unenriched, activity-enriched, and TCR trap early fraction lipids. Three experiments were performed for (A) and (B). Experiments in (C) were performed on lipids extracted two independent TCR trap assays.

three proteins matching the apparent molecular weights of CD1d,  $\beta$ 2m, and TCR with band similar intensity, consistent with 1:1:1 stoichiometry (Fig. 2D). The peak height of the protein trace of the early eluting fraction made with milk lipids (Fig. 2C) was substantially lower than that of the early fraction made with CD1d carrying synthetic  $\alpha$ -glucosylceramide (Fig. 2A), suggesting that TCRs bound to a small fraction of the total pool of CD1d-lipid complexes present. However, since this peak was observed at the same elution time as the synthetic  $\alpha$ -glucosylceramide standard-loaded CD1d, it suggested that the milk lipid-derived ternary complex carried a high-affinity antigen.

Nanoelectrospray analysis of all eluted milk lipids in the late fraction containing CD1d and TCR monomers found numerous ions, and the mass intervals among the most abundant ions were diverse, suggesting the presence of structurally diverse lipids from many classes (Fig. 3A, upper). In contrast, the early fraction, which contained CD1d-TCR complexes, showed fewer eluted ions (Fig. 3A, lower). The lower peak height from the size exclusion tracing and the less complex MS profiles suggested that the iNKT cell TCR selected small subset of CD1d-milk lipid complexes from among all CD1d-lipid complexes present in the starting material. In fact, nearly all nanoelectrospray MS signals derived from the early-eluting CD1d-TCR complexes could be accounted for by a single alkane series comprised of lipids of  $m/z$  710.7, apparent chain-length analogs ( $m/z$  724.7, 738.7, 752.6, 794.7, 808.8, 822.8, 836.8), and the isotopes of these lipids.

**TCR ligands identified using nanoelectrospray MS.** Next we identified the lipid ligands trapped-in or excluded-from CD1d-TCR complexes using shotgun nanoelectrospray with multi-stage ion trap MS. This method provides high sensitivity, broad detection, and structural information from collision-induced dissociation (CID), but offers relatively low mass accuracy. Considering lipids eluted from the late fraction containing CD1d and TCR monomers, CID-MS of selected ions at  $m/z$  738.4, 753.5, and 808.5 identified phosphatidylethanolamine, sphingomyelin, and phosphatidylcholine (Fig. 3B-D). In addition, fragments from the ion at  $m/z$  722.6 corresponded a hexosylceramide whose two fatty acid chains had a combined lipid tail length of C34 with one unsaturation (Fig. 3E). This 'type 1' monohexosylceramide collision pattern was characterized by a dehydration (loss of 18 mu,  $m/z$  704.5), loss of a hexosyl unit (162 mu,  $m/z$  560.5), and loss of both fragments (loss of 180 mu,  $m/z$  542.4). In contrast, ions from early fraction containing CD1d-TCR complexes did not show detectable signals for phosphatidylethanolamine, sphingomyelin, or phosphatidylcholine. Instead, complexes in the early fraction released matched expected masses of a 'type 2' hexosylceramide, which unlike type 1 patterns, lacked a dehydration ion but did show a loss of 162 mu. In fact, the abundant ions in the early

fraction were all accounted for by single alkane series corresponding to an monohexosylceramide with an overall lipid length ranging between C33 ( $m/z$  710.7) and C42 ( $m/z$  836.8). Separate analysis of six members of this alkane series demonstrated highly reproducible 'type 2' collision pattern (Fig. 3F).

The detailed profiles of these released lipids suggested several general conclusions. First, monomeric CD1d proteins were bound to phospholipids, sphingomyelin, type 1 hexosylceramide, as well as other lipids that were not specifically identified (Fig. 3A). Second, TCR-CD1d complex formation largely or completely excluded CD1d bound to phosphatidylcholine ( $m/z$  808.6 with the  $m/z$  749.3 product), sphingomyelin ( $m/z$  753.6 with  $m/z$  694.4 product), phosphatidylethanolamine ( $m/z$  738.5 with  $m/z$  695.3 product), and the type 1 hexosylceramide with a monounsaturated lipid tail ( $m/z$  722.6 with  $m/z$  542.4 fragment). Third, lipids eluted from the TCR-CD1d complexes contained type 2 monohexosylceramides ranging from C33-42 in chain length (Fig. 3A). Further, the characteristic patterns of loss in CID analyses by type 1 (18, 162, and 180 mu) and type 2 (162 mu) hexosylceramides implicated a differing conformation of these ceramide-based lipids. This interpretation is consistent with the TCR discriminating structural classes of lipids including glycosylceramides, and possibly  $\alpha$ - versus  $\beta$ -linked hexosylceramides.

The molar concentrations of hexosylceramides eluted from the early and late complexes were determined using HPLC-MS in comparison to standards, and this material tested for their ability to elicit cytokine production from an iNKT cell hybridoma, DN32. The late fraction did not have detectable activity, while the molar activity observed in the early fraction was similar to that of a synthetic antigen for iNKT cells known as  $\alpha$ -galactosylceramide, KRN7000 (Fig. 3G). Thus, our approach was successful as a TCR-based affinity trap for selecting antigenic glycolipids that ligate TCR to CD1d out of a complex mixture of natural lipids. The equipotency of type 2 hexosylceramides and  $\alpha$ -galactosylceramide was consistent with the type 2 hexosylceramides being an  $\alpha$ -linked lipid. Although only trace amounts of the type 2 monohexosylceramide could be recovered from ternary complexes of CD1d-lipid-TCR, 1D-proton nuclear magnetic resonance spectroscopy (NMR) carried out on a 900 MHz instrument identified a doublet with a chemical shift of 4.85 ppm, consistent with the anomeric proton resonance of  $\alpha$ -monohexosylceramide (Fig. S1) (15).

**Identification of ligands at high mass resolution.** To further biochemically characterize the lipids that promoted the formation of TCR-lipid-CD1d complexes, we separately analyzed eluents from milk lipid-treated protein monomers present in the late fraction, untreated CD1d, and untreated TCR (Fig. 4). Here, we took advantage of the higher mass accuracy of a quadrupole time-of-flight (Q-ToF) mass detector coupled with chromatographic

separation in normal-phase HPLC-MS to confirm the identity of the key ions. Using this method, the major ions previously identified by nanoelectrospray MS were confirmed and assigned based on high mass accuracy and HPLC retention time, compared with standards as phosphatidylcholine (38.1 min,  $m/z$  786.601), phosphatidylethanolamine (20.5 min,  $m/z$  716.523), type 1 hexosylceramide (4.1 min,  $m/z$  700.578), and type 2 hexosylceramide (4.1 min,  $m/z$  702.590, 716.604 and 786.682) (Figure 4). No substantial false positive signals were derived from lipids released from the TCR, indicating that the lipids detected were carried by CD1d. As seen with the nanospray method (Fig. 3), CD1d alone and mixed CD1d and TCR monomers bound phosphatidylcholine, phosphatidylethanolamine, and type 1 monohexosylceramides (Fig. 4 and S2), but these were excluded from CD1d-TCR complexes. Conversely, type 2 hexosylceramides eluted with low or undetectable signals from milk lipid treated monomers, but were highly enriched in CD1d-TCR complexes. Thus, CD1d bound many lipids, but the iNKT TCRs selectively captured CD1d bound to type 2 hexosylceramides (Fig. S3).

#### Structural identification of type 1 and 2 hexosylceramides.

Despite their strikingly-different patterns of binding within, or being excluded from, CD1d-TCR complexes, normal-phase HPLC-MS showed that type 1 and 2 hexosylceramides had equivalent retention time and the same underlying molecular composition, except for variations in lipid tail length and unsaturation. CID-MS experiments using the ion trap architecture, however showed different fragmentation patterns for type 1 and type 2 hexosylceramides (Fig. 3). We next asked whether anomeric variants could be distinguished by CID-MS. First, we produced synthetic standards differing only in the stereochemistry of their hexose attachment. Both glucosyl and galactosyl variants were synthesized, with dihydrosphingosine (d18:0) as their lipid backbones based on the molecular composition of type 2 hexosylceramides identified in the milk-derived early fraction. These synthetic standards had a similar retention time by normal-phase HPLC to the early-eluting milk lipids (Fig. S4). The synthetic standards were tested for their ability to activate the DN32 iNKT cell hybridoma after loading in CD1d-expressing RAW264.7 cells. For both glucosyl and galactosyl synthetic standards, activity was observed for the  $\alpha$ -linked variants, but not for the  $\beta$ -linked variants (Fig. 5A).

Positive mode ion trap CID-MS of sodiated adducts of these synthetic hexosylceramides was carried out under conditions that matched those used for milk-derived compounds (Fig. 3), and over a range of collision energy settings (Fig. 5B). Collision of the  $\alpha$ -glucosylceramide synthetic standard demonstrated almost exclusively a primary loss of 162 mu, matching the pattern for type 2 milk-derived hexosylceramides. In contrast,  $\beta$ -glucosyl standards matched the fragmentation pattern seen for type 1 hexosylceramides from milk, showing a loss of 18 mu at a similar frequency as loss of 162 mu. Although both patterns involved the loss of 162 mu as a hexosyl unit, analysis over a range of collision energy demonstrated that  $\beta$ -glucosylceramide also lost 18 mu and 180 mu, whereas  $\alpha$ -glucosylceramide showed neither of these fragmentations at a substantial frequency. Parallel analyses of  $\alpha$ - and  $\beta$ -galactosylceramide standards showed similar characteristic  $\alpha$ - and  $\beta$ - collision patterns (Fig. S5). Matched synthetic  $\alpha$ - and  $\beta$ -glucosylceramide and galactosylceramide standards based on the type 1 hexosylceramide lipid backbone structure showed similar characteristic  $\alpha$ - and  $\beta$ - collision patterns, demonstrating that the anomeric linkage discrimination by this technique was not limited to dihydrosphingosine-based glycosphingolipids (Fig. S6). Ion trap CID-MS of  $\alpha$ -galactosylceramide with a phytosphingosine base also fragmented with the pattern matching type 2 hexosylceramides. We concluded from these experiments that linear ion trap MS could sensitively distinguish  $\alpha$ - from  $\beta$ -hexosylceramides under the conditions measured for a range of lipids with differing tail lengths and unsaturations.

We compared the collision patterns of the major hexosylceramide ions from cow's milk lipids. Unfractionated cow's milk glucosylceramides demonstrated a fragmentation pattern characteristic of  $\beta$ -hexosylceramide standards (Fig. 5C), consistent with the main lipid in the starting material being  $\beta$ -glucosylceramide. The activity-enriched cow's milk glucosylceramide showed a  $\beta$ -hexosylceramide collision pattern, but with an increased frequency of a type 2 pattern characterized by the loss of the sugar headgroup (162 mu). The antigenic lipids captured in the TCR trap assay showed the fragmentation pattern typical of  $\alpha$ -hexosylceramides. We concluded from these experiments that the TCR trap captured type 2  $\alpha$ -linked hexosylceramides from cow's milk. To address the possibility that these type 2 hexosylceramides were derived from microbes present in cow's milk, we purified the monohexosylceramide fraction from calf thymus, a mammalian lipid source previously shown activate iNKT cells (15). Several ions with the  $m/z$  seen in cow's milk were also present in calf thymus, including ions corresponding to type 2 hexosylceramides  $m/z$  724.7 and 808.8 (Fig. S7). Linear ion trap CID-MS of untreated calf thymus hexosylceramides showed a characteristic  $\beta$ -hexosylceramide pattern. After activity enrichment of thymus hexosylceramides by treatment with  $\beta$ -glucosidase and TLC fractionation as in Fig. 1, we observed a collision pattern consistent with the presence of  $\alpha$ -hexosylceramides, based on a substantial increase in the abundance of the ion representing a primary loss of the sugar headgroup.

#### Discussion

Uncovering the specific antigens recognized by iNKT cells is central to understanding their roles in health and disease. Although lipid antigens for iNKT cells have been identified in the microbial world (6-10), experimental evidence has demonstrated important roles for endogenous and environmental lipid antigens. Several candidate endogenous lipid antigens have been proposed (13, 14, 20-22), though technical limitations in the ability to detect these antigens has hampered our ability to determine their contextual roles. Here, we provide direct biochemical evidence for  $\alpha$ -hexosylceramide derived from a mammalian source, cow's milk.

There is increasing evidence that iNKT cells encounter lipid antigens at mucosal surfaces. CD1-dependent immunomodulatory lipids have been identified in gut microbes (11, 12), pollen (23), house dust (24), and in our work, cow's milk. We have previously demonstrated that cow's milk contains substantially more lipid antigen activity than human or mouse milk. Although native triglycerides are removed from infant formula during processing, bovine polar lipids, including lipid antigens for iNKT cells, are retained in the final product (15). Studies in germ-free mice suggest that the iNKT cell compartment can be shaped by CD1d-lipid signals in the enteric tract (11, 25), and thus early-life exposure to cow's milk-based infant formula might similarly shape human mucosal iNKT cells in early life. In adult mice, orally-delivered antigen activates iNKT cells in the gut (26), suggesting that exposure to dietary lipid antigens in adult life may also have consequences. Interestingly, it is unclear whether cows have iNKT cells (27), as their CD1d does not present the long-chain  $\alpha$ -galactosylceramide commonly used as a tool in mice and humans (28). CD1d can present short-chain  $\alpha$ -galactosylceramides (28, 29), and it is possible that an increased presence of  $\alpha$ -hexosylceramides in their milk and tissues could lead to thymic deletion of iNKT cells, as has been observed in mice in the presence of exogenous  $\alpha$ -galactosylceramide (30).

Although the bulk solids in milk derive overwhelmingly from mammary glands, we cannot formally rule out that colonizing microbes synthesized, or played a role in the synthesis of the  $\alpha$ -hexosylceramides present in cow's milk. Several features of our data argue against possible contamination of milk by microbial products. We have previously observed iNKT cell-

681 stimulating activity in the monohexosylceramide fractions of  
 682 the milks, sera, and internal organs of multiple animal species  
 683 (14, 15). The *N*-acyl chain usage of the  $\alpha$ -hexosylceramides  
 684 that we identified in this report mirror that of the bulk  $\beta$ -  
 685 glucosylceramides in cow's milk, suggesting that these lipids are  
 686 of mammalian origin. We also identified type 2 hexosylceramides  
 687 from calf thymus with the same *m/z* as those in cow's milk.  
 688 These thymic hexosylceramides, like those found in cow's milk,  
 689 shifted from a  $\beta$ -hexosylceramide pattern toward a characteris-  
 690 tic  $\alpha$ -hexosylceramide collision pattern after activity enrichment.  
 691 Currently, the level of  $\alpha$ -hexosylceramide identified from cow's  
 692 milk, a relatively abundant and potent natural source of lipid  
 693 antigen, is near the limit of detection for the TCR trap assay  
 694 system. However, we are continuing to improve the system so  
 695 that lipids can be captured and identified from smaller amounts  
 696 of source material.

697 The stereochemistry of organic molecules can have significant  
 698 biological consequences. Determination of stereochemistry has  
 699 largely been accomplished by NMR. However, for rare com-  
 700 pounds such as endogenous iNKT cell lipid antigens, purification  
 701 of sufficient material for complete, multi-dimensional NMR anal-  
 702 ysis is currently not possible. As shown in this report and others  
 703 (31, 32), CID-MS can also be used to determine stereochemical  
 704 features. We tested  $\alpha$ - and  $\beta$ -linked synthetic standards with both  
 705 sphingosine and dihydrospingosine bases, and found consistent  
 706 characteristic  $\alpha$ - and  $\beta$ - fragmentation patterns, showing that this  
 707 detection method was not limited to certain lipid backbones.  
 708 Interestingly, these collision patterns were only observed with  
 709 selected positive mode adducts, including sodium and rubidium.  
 710 We hypothesize that for the  $\beta$ -hexosylceramide structure, positive  
 711 adducts with a sufficiently large valence, such as sodium and  
 712 rubidium, can coordinate elements on both sides of the glycoside  
 713 bond, stabilizing the glycoside bond during collision, thus reduc-  
 714 ing the propensity for cleavage of the sugar headgroup. For  $\alpha$ -  
 715 hexosylceramide, this headgroup stabilization may be sterically

hindered, thus allowing for fragmentation across the glycoside  
 bond at a lower energy.

749 Identification of CD1d-presented lipid antigens has largely  
 750 relied on indirect evidence such as extrapolation from cells de-  
 751 ficient in lipid synthesis genes (22, 33), or fractionation of crude  
 752 lipid sources with activity tracking within these fractions (14,  
 753 20). These approaches have been useful, but do not directly  
 754 demonstrate recognition by the TCR, and rely on correlations  
 755 between two or more assays. Here, we used a direct method for  
 756 antigen discovery, the TCR trap assay. This approach that uses  
 757 ternary complex formation both as the preparative method to  
 758 recover lipids of interest and the key assay to demonstrate the  
 759 role of these lipids in ligating CD1d and TCR. The TCR trap  
 760 assay has also been used successfully to identify CD1a ligands  
 761 (17), and could be used to identify ligands for other CD1 isoforms,  
 762 or even MR1. This approach will allow tracking of rare lipids  
 763 from virtually any source, including from complex mixtures, or  
 764 limited starting materials, which represent common challenges  
 765 when starting with mammalian tissues and cells.

#### 766 Methods and materials

767 Initial lipid activity enrichment was performed as previously  
 768 described (15). Mass spectrometry analyses were performed on  
 769 a Thermo-Fisher LXQ ion trap mass spectrometer or an Agilent  
 770 6520 Accurate-Mass Q-ToF with HPLC. For details of experimen-  
 771 tal conditions and analysis, see SI Materials and Methods.

772 **Acknowledgements.** We thank C. Costello, C. Heiss, and P. Azadi for helpful  
 773 discussions and J. Buter for assistance with figure production. P.J.B. was  
 774 supported by the NIH (AI102945) and generous support from the Violin and  
 775 Karol families, D.B.M. (AR048632, AI111224, and the Bill and Melinda Gates  
 776 Foundation), M.B.B. NIH (AI113046, AI063428), G.S.B. J. Badrick, the Royal  
 777 Society, and the Medical Research Council, DIG and DGP were supported  
 778 by the National Health and Medical Research Council of Australia ((NHMRC)  
 779 1013667, 1113293, 1020770, 1117766 and 1054431) and the Australian Re-  
 780 search Council (ARC; CE140100011), JR by the Australian Research Council  
 781 (FL160100049, CE140100011).

1. Godfrey DI, Uldrich AP, McCluskey J, Rossjohn J, & Moody DB (2015) The burgeoning family of unconventional T cells. *Nat Immunol* 16(11):1114-1123.
2. Young MH & Gapin L (2011) Group 1 CD1-restricted T cells take center stage. *Eur J Immunol*.
3. Van Rhijn I, Godfrey DI, Rossjohn J, & Moody DB (2015) Lipid and small-molecule display by CD1 and MR1. *Nat Rev Immunol* 15(10):643-654.
4. Brennan PJ, Brigl M, & Brenner MB (2013) Invariant natural killer T cells: an innate activation scheme linked to diverse effector functions. *Nat Rev Immunol* 13(2):101-117.
5. Kawano T, et al. (1997) CD1d-restricted and TCR-mediated activation of valpha14 NKT cells by glycosylceramides. *Science* 278(5343):1626-1629.
6. Kinjo Y, et al. (2005) Recognition of bacterial glycosphingolipids by natural killer T cells. *Nature* 434(7032):520-525.
7. Mattner J, et al. (2005) Exogenous and endogenous glycolipid antigens activate NKT cells during microbial infections. *Nature* 434(7032):525-529.
8. Sriram V, Du W, Gervay-Hague J, & Brutkiewicz RR (2005) Cell wall glycosphingolipids of *Sphingomonas paucimobilis* are CD1d-specific ligands for NKT cells. *Eur J Immunol* 35(6):1692-1701.
9. Kinjo Y, et al. (2006) Natural killer T cells recognize diacylglycerol antigens from pathogenic bacteria. *Nat Immunol* 7(9):978-986.
10. Kinjo Y, et al. (2011) Invariant natural killer T cells recognize glycolipids from pathogenic Gram-positive bacteria. *Nat Immunol* 12(10):966-974.
11. An D, et al. (2014) Sphingolipids from a symbiotic microbe regulate homeostasis of host intestinal natural killer T cells. *Cell* 156(1-2):123-133.
12. Wieland Brown LC, et al. (2013) Production of alpha-galactosylceramide by a prominent member of the human gut microbiota. *PLoS Biol* 11(7):e1001610.
13. Kain L, et al. (2014) The identification of the endogenous ligands of natural killer T cells reveals the presence of mammalian alpha-linked glycosylceramides. *Immunity* 41(4):543-554.
14. Brennan PJ, et al. (2011) Invariant natural killer T cells recognize lipid self antigen induced by microbial danger signals. *Nat Immunol* 12(12):1202-1211.
15. Brennan PJ, et al. (2014) Activation of iNKT cells by a distinct constituent of the endogenous glycosylceramide fraction. *Proc Natl Acad Sci U S A* 111(37):13433-13438.
16. Burdin N, et al. (1998) Selective ability of mouse CD1 to present glycolipids: alpha-galactosylceramide specifically stimulates V alpha 14+ NK T lymphocytes. *J Immunol* 161(7):3271-3281.
17. Birkinshaw RW, et al. (2015) alpha beta T cell antigen receptor recognition of CD1a presenting self lipid ligands. *Nat Immunol* 16(3):258-266.
18. Cardell S, et al. (1995) CD1-restricted CD4+ T cells in major histocompatibility complex class II-deficient mice. *J Exp Med* 182(4):993-1004.
19. Patel O, et al. (2012) Recognition of CD1d-sulfatide mediated by a type II natural killer T cell antigen receptor. *Nat Immunol* 13(9):857-863.
20. Facciotti F, et al. (2012) Peroxisome-derived lipids are self antigens that stimulate invariant natural killer T cells in the thymus. *Nat Immunol* 13(5):474-480.
21. Fox LM, et al. (2009) Recognition of lyso-phospholipids by human natural killer T lymphocytes. *PLoS Biol* 7(10):e1000228.
22. Zhou D, et al. (2004) Lysosomal glycosphingolipid recognition by NKT cells. *Science* 306(5702):1786-1789.
23. Agea E, et al. (2005) Human CD1-restricted T cell recognition of lipids from pollens. *J Exp Med* 202(2):295-308.
24. Wingender G, et al. (2011) Invariant NKT cells are required for airway inflammation induced by environmental antigens. *J Exp Med* 208(6):1151-1162.
25. Olszak T, et al. (2012) Microbial exposure during early life has persistent effects on natural killer T cell function. *Science* 336(6080):489-493.
26. Lee YJ, et al. (2015) Tissue-Specific Distribution of iNKT Cells Impacts Their Cytokine Response. *Immunity* 43(3):566-578.
27. Van Rhijn I, et al. (2006) The bovine CD1 family contains group 1 CD1 proteins, but no functional CD1d. *J Immunol* 176(8):4888-4893.
28. Wang J, et al. (2012) Crystal structures of bovine CD1d reveal altered alphaGalCer presentation and a restricted A' pocket unable to bind long-chain glycolipids. *PLoS One* 7(10):e47989.
29. Nguyen TK, et al. (2013) The bovine CD1D gene has an unusual gene structure and is expressed but cannot present alpha-galactosylceramide with a C26 fatty acid. *Int Immunol* 25(2):91-98.
30. Pellicci DG, et al. (2003) Intrathymic NKT cell development is blocked by the presence of alpha-galactosylceramide. *Eur J Immunol* 33(7):1816-1823.
31. Kanie O, et al. (2009) Analysis of behavior of sodiated sugar hemiacetals under low-energy collision-induced dissociation conditions and application to investigating mutarotation and mechanism of a glycosidase. *Proc Jpn Acad Ser B Phys Biol Sci* 85(6):204-215.
32. Moody DB, et al. (2000) CD1c-mediated T-cell recognition of isoprenoid glycolipids in *Mycobacterium tuberculosis* infection. *Nature* 404(6780):884-888.
33. Darmono A, et al. (2010) Lysosomal alpha-galactosidase controls the generation of self lipid antigens for natural killer T cells. *Immunity* 33(2):216-228.
34. Jervis PJ, et al. (2010) Synthesis and biological activity of alpha-glucosyl C24:0 and C20:2 ceramides. *Bioorg Med Chem Lett* 20(12):3475-3478.
35. Uldrich AP, et al. (2013) CD1d-lipid antigen recognition by the gammadelta TCR. *Nat Immunol* 14(11):1137-1145.
36. Matsuda JL, et al. (2000) Tracking the response of natural killer T cells to a glycolipid antigen using CD1d tetramers. *J Exp Med* 192(5):741-754.
37. Pellicci DG, et al. (2009) Differential recognition of CD1d-alpha-galactosyl ceramide by the V beta 8.2 and V beta 7 semi-invariant NKT T cell receptors. *Immunity* 31(1):47-59.

817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884

38. Lantz O & Bendelac A (1994) An invariant T cell receptor alpha chain is used by a unique subset of major histocompatibility complex class I-specific CD4+ and CD4-8- T cells in mice and humans. *J Exp Med* 180(3):1097-1106.

39. Muindi K, et al. (2010) Activation state and intracellular trafficking contribute to the repertoire of endogenous glycosphingolipids presented by CD1d [corrected]. *Proc Natl Acad Sci U S A* 107(7):3052-3057.

40. Bligh EG & Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Canadian journal of biochemistry and physiology* 37(8):911-917.

41. Layre E, et al. (2011) A comparative lipidomics platform for chemotaxonomic analysis of *Mycobacterium tuberculosis*. *Chem Biol* 18(12):1537-1549.

885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952

# Submission PDF