

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/108334/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Lechner, G., Li, D., Queffélec, H. and Rodriguez-Piazza, L. 2018. Approximation numbers of weighted composition operators. *Journal of Functional Analysis* 274 (7) , pp. 1928-1958. 10.1016/j.jfa.2018.01.010 file

Publishers page: <https://doi.org/10.1016/j.jfa.2018.01.010> <<https://doi.org/10.1016/j.jfa.2018.01.010>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Approximation numbers of weighted composition operators

G. Lechner*, D. Li†, H. Queffélec‡, L. Rodríguez-Piazza§

December 24, 2017

Abstract

We study the approximation numbers of weighted composition operators $f \mapsto w \cdot (f \circ \varphi)$ on the Hardy space H^2 on the unit disc. For general classes of such operators, upper and lower bounds on their approximation numbers are derived. For the special class of weighted lens map composition operators with specific weights, we show how much the weight w can improve the decay rate of the approximation numbers, and give sharp upper and lower bounds. These examples are motivated from applications to the analysis of relative commutants of special inclusions of von Neumann algebras appearing in quantum field theory (Borchers triples).

1 Introduction

In the study of composition operators $C_\varphi : f \mapsto f \circ \varphi$ acting on a Hilbert space H of analytic functions (on the unit disk \mathbb{D}), one is typically interested in understanding how function-theoretic properties of φ are related to operator-theoretic properties of C_φ . Basic properties such as boundedness or compactness of C_φ are by now well characterized in terms of φ in many cases [35, 16]. More recently, also the membership of C_φ in various smaller ideals I of bounded operators on H (such as the p -Schatten class),

*Cardiff University, School of Mathematics, LechnerG@cardiff.ac.uk

†Université d'Artois, Laboratoire de Mathématiques de Lens (LML), daniel.li@euler.univ-artois.fr

‡Université Lille Nord de France, Herve.Queffelec@univ-lille1.fr

§Universidad de Sevilla, piazza@us.es

and more precisely the behavior of the approximation numbers $a_n(C_\varphi)$ of C_φ , was studied in depth in several papers (see e.g. [27, 25, 26, 28]).

If $\mathcal{M}(H)$ denotes the space of multipliers of H (those $w \in H$ such that $wf \in H$ for each $f \in H$), we can twist a composition operator C_φ , assumed to map H to itself, by composing it on the left with the operator M_w of multiplication by $w \in \mathcal{M}(H)$. We then get a so-called *weighted composition operator* $T = M_w C_\varphi$ (see e.g. [24] or [12, 21]).

A careful distinction must be made between the multipliers of H , denoted $\mathcal{M}(H)$, and those of $C_\varphi(H)$, denoted $\mathcal{M}(H, \varphi)$, namely those functions $w \in H$ such that $wf \in H$ for each f belonging to the range $C_\varphi(H)$, not necessarily to the whole of H . For example, if $H = H^2$ is the Hardy space, then $\mathcal{M}(H) = H^\infty$ consists of all bounded analytic functions on \mathbb{D} . It can be proved that (see [2, 14] and [19], respectively):

$$\begin{aligned}\mathcal{M}(H^2, \varphi) = H^\infty &\Leftrightarrow \varphi \text{ is a finite Blaschke product,} \\ \mathcal{M}(H^2, \varphi) = H^2 &\Leftrightarrow \|\varphi\|_\infty < 1.\end{aligned}$$

In this paper, we study approximation numbers of weighted composition operators in the case of the Hardy space $H = H^2$ of the disc, and a weight $w \in \mathcal{M}(H) \subset \mathcal{M}(H, \varphi)$. Then, since we are dealing with ideals, twisting with the bounded operator M_w can but reinforce the membership in I , and improve the rate of decay of approximation numbers.

There are (at least) three motivations for considering weighted composition operators: First, they form a natural and non-trivial generalization of composition operators on $H^2(\mathbb{D})$. In this context, it is natural to ask how much faster the approximation numbers $a_n(M_w C_\varphi)$ can decay in comparison to the $a_n(C_\varphi)$. For example, can $M_w C_\varphi$ be compact when C_φ is non-compact, or can the $a_n(M_w C_\varphi)$ decay quite fast when the $a_n(C_\varphi)$ decay rather slowly? We will address these questions in the body of the text¹.

As a second motivation, suppose C_φ^G is a composition operator on a Hardy space $H^2(G)$ over a simply connected region properly contained in \mathbb{C} . Then a choice of Riemann map $\tau : \mathbb{D} \rightarrow G$ induces a unitary between $H^2(\mathbb{D})$ and $H^2(G)$ [17], and we can equivalently formulate C_φ^G as an operator on $H^2(\mathbb{D})$. This operator on $H^2(\mathbb{D})$, however, turns out to be a *weighted* composition operator $M_{w_\tau} C_{\varphi_\tau}^{\mathbb{D}}$ in general (see [36] and Section 3). Thus composition operators on domains other than \mathbb{D} automatically produce weighted composition operators on $H^2(\mathbb{D})$.

¹Another application of weighted composition operators to the study of composition operators on spaces of several complex variables can be found in [29].

A third motivation for studying weighted composition operators comes from applications in a completely different field, namely inclusions of von Neumann algebras, used in mathematical physics to model quantum field theories [20]. For \mathcal{N} a von Neumann algebra with a cyclic and separating vector Ω on a Hilbert space \mathcal{H} , we will consider the Hilbert space \mathcal{D} obtained by closing the domain of the modular operator of (\mathcal{N}, Ω) [38] in its graph norm. If \mathcal{N} carries additional structure (a Borchers triple), this setting is related to complex analysis because an irreducible component of \mathcal{D} can be naturally identified with a Hardy space $H^2(\mathbb{S})$ on a strip region $\mathbb{S} \subset \mathbb{C}$, bounded by two lines parallel to \mathbb{R} (see Section 3).

In applications in mathematical physics, one is interested in specific inclusions $\tilde{\mathcal{N}} \subset \mathcal{N}$ and their relative commutants, the size of which can be controlled if a map built from the modular operator has sufficiently quickly decaying approximation numbers [8]. On the level of the irreducible component giving rise to the Hardy space $H^2(\mathbb{S})$, this condition translates to a weighted restriction operator $R_w : H^2(\mathbb{S}) \rightarrow L^2(\mathbb{R})$, $f \mapsto (w \cdot f)|_{\mathbb{R}}$, where the real line \mathbb{R} lies in the interior of the strip, and $w \in H^\infty(\mathbb{S})$ is an inner function on \mathbb{S} obtained from the inclusion $\tilde{\mathcal{N}} \subset \mathcal{N}$. For the application in physics, sharp upper bounds on the approximation numbers of R_w are desirable [1].

Mapping the strip \mathbb{S} to the disc, R_w can be formulated as a Carleson embedding operator (the definition of which we recall in Section 2.1). These operators are often used in estimating approximation numbers of composition operators [25, 26]. In turn, we find from the strip picture that the embedding operator can be estimated from above by special weighted composition operators on the disc, namely those whose symbol is a lens map $\varphi = \varphi_\lambda$, $0 < \lambda < 1$ (see Section 4 for the definition). This closes the connection to composition operators on $H^2(\mathbb{D})$, where the C_{φ_λ} are among the best studied examples [25].

Given these motivations, this article is organized as follows. In Section 2, we introduce our notation and setup, and study weighted composition operators with general symbols φ and weights w on the disc. After deriving a simple upper bound, we give an example how a weight can turn a non-compact composition operator into a compact one. Regarding lower bounds, we show that the worst possible behavior of the $a_n(C_\varphi)$ (exponential if $\|\varphi\|_\infty < 1$, subexponential if $\|\varphi\|_\infty = 1$) is the same for the weighted operators $M_w C_\varphi$.

In Section 3, we explain the links between modular theory of von Neumann algebras, Hardy spaces on strips, and weighted restriction operators. In that section we also show how weighted lens map composition operators

appear. Section 3 can be read independently of the other parts of the article.

Finally, we consider in Section 4 the specific case of weighted lens map composition operators $M_w C_{\varphi_\lambda}$ as our primary example. In the case without weight, the approximation numbers of C_{φ_λ} are known to decay like $e^{-c\sqrt{n}}$ [26]. A natural question in this context is how close the decay rate of the $a_n(M_w C_{\varphi_\lambda})$ can come to exponential decay e^{-cn} (the optimal one in the context of composition operators). For the weights motivated by the considerations in Section 3, we show that $a_n(M_w C_{\varphi_\lambda})$ decays like $e^{-c\frac{n}{\log n}}$.

2 Weighted composition operators on \mathbb{D}

2.1 Preliminaries

We begin by recalling a few operator-theoretic and function-theoretic facts. The approximation numbers $a_n(T) = a_n$ of an operator $T : H \rightarrow H$ (with H a Hilbert space) are defined by

$$a_n = \inf_{\text{rank } R < n} \|T - R\|,$$

and T is compact if and only if $\lim_{n \rightarrow \infty} a_n(T) = 0$. According to a result of Allahverdiev [11, p. 155], $a_n = s_n$, the n -th singular number of T . We have the following alternative definition (a variant of Kolmogorov numbers) of $a_n(T)$ [28], in which B_H denotes the closed unit ball of H and $d(g, A)$ the distance of g to $A \subset H$:

$$(2.1) \quad a_n(T) = \inf_{\dim E < n} \left[\sup_{f \in B_H} d(Tf, TE) \right].$$

The definition of $a_n(T)$ also makes sense for $T : X \rightarrow Y$ an operator between Banach spaces (see Theorem 2.8 to come).

Coming back to the hilbertian setting, two other useful alternative definitions (respectively in terms of *Bernstein* and *Gelfand* numbers) are, denoting by S_E the unit sphere of a subspace E of H (see [11, Chapter 2], or [27]):

$$(2.2) \quad a_n(T) = \sup_{\dim E = n} \left[\inf_{f \in S_E} \|Tf\| \right],$$

$$(2.3) \quad a_n(T) = \inf_{\text{codim } E < n} \|T|_E\| = \inf_{\text{codim } E < n} \left[\sup_{f \in S_E} \|Tf\| \right].$$

The following parameters $0 \leq \beta^-(T) \leq \beta^+(T) \leq 1$ were used in [28]:

$$\beta^+(T) = \limsup_{n \rightarrow \infty} [a_n(T)]^{1/n}, \quad \beta^-(T) = \liminf_{n \rightarrow \infty} [a_n(T)]^{1/n}.$$

When the limit exists, we denote it by $\beta(T)$. It is proved in [28] that this is the case for T a composition operator on the Hardy, Bergman, or Dirichlet space. Observe that $\beta^-(T) = 1$ signifies a subexponential decay for $a_n(T)$, namely $a_n(T) \geq e^{-n\varepsilon_n}$ where $\varepsilon_n > 0$ and $\varepsilon_n \rightarrow 0$.

Let now \mathbb{D} be the open unit disk of the complex plane and H^2 the usual Hardy space of \mathbb{D} . Recall [35, p. 12] that the norm of $f(z) = \sum_{n=0}^{\infty} f_n z^n \in H^2$ is defined by $\|f\|_2^2 = \sum_{n=0}^{\infty} |f_n|^2$, or alternatively by

$$(2.4) \quad \|f\|_2^2 = \int_{\mathbb{T}} |f^*(u)|^2 dm(u)$$

where m denotes the Haar measure of the unit circle \mathbb{T} and $f^*(u)$ is the (m -almost everywhere existing by Fatou's theorem) radial limit $\lim_{r \rightarrow 1^-} f(ru)$, often again denoted $f(u)$.

The space of multipliers of H^2 is isometrically isomorphic to the space H^∞ of functions analytic and bounded on \mathbb{D} [18]. This means that any function $w \in H^\infty$ defines a bounded multiplication operator $M_w : H^2 \rightarrow H^2$ by the formula $M_w(f) = wf$ and that

$$\|M_w\| := \sup_{\|f\|_2 \leq 1} \|wf\|_2 = \|w\|_\infty := \sup_{z \in \mathbb{D}} |w(z)|.$$

If φ is a *non-constant* and analytic self-map of \mathbb{D} (often called a symbol), the associated composition operator $C_\varphi : H^2 \rightarrow H^2$ is defined by

$$C_\varphi(f) = f \circ \varphi.$$

The fact that C_φ boundedly maps H^2 to itself for any symbol φ is the well-known subordination principle of Littlewood ([16, p. 29], [35, p. 16]).

Next, a positive and bounded measure μ on \mathbb{D} is called a Carleson measure (for H^2) if the identity map $R_\mu, R_\mu(f) = f$ maps $H = H^2$ to $L^2(\mu)$, that is if there exists a constant C such that:

$$\int_{\mathbb{D}} |f(z)|^2 d\mu(z) \leq C \|f\|_2^2 \quad \forall f \in H^2.$$

The best constant C is called the Carleson-norm of μ and is denoted $\|\mu\|_C$. That is $\|\mu\|_C = \|R_\mu\|^2$. Let us set

$$\rho_\mu(h) := \sup_{\xi \in \mathbb{T}} \mu[S(\xi, h)]$$

where, for $\xi \in \mathbb{T} = \partial\mathbb{D}$, $S(\xi, h)$ is the Carleson box

$$S(\xi, h) = \{z \in \mathbb{D} : |z - \xi| \leq h\}.$$

With those notations, the Carleson embedding theorem [16, p. 37] gives a geometric characterization of Carleson measures:

Theorem 2.1. *Let μ be a positive and bounded measure on \mathbb{D} . Then, μ is a Carleson measure if and only if, for some constant K ,*

$$\rho_\mu(h) \leq Kh \quad \forall h \in]0, 1].$$

In this case, $\|\mu\|_C \leq aK$ where $a > 0$ is an absolute constant.

Let again $\varphi^*(u) = \lim_{r \rightarrow 1^-} \varphi(ru)$. Littlewood's subordination principle implies that $m_\varphi = \varphi^*(m)$, the image under φ^* of this Haar measure, is a Carleson measure for H^2 , and we will write ρ_φ instead of ρ_{m_φ} . We will also write $\|\cdot\|$ instead of $\|\cdot\|_2$ when there is no ambiguity.

The main subject of this article are *weighted composition operators* on H^2 , defined in terms of a weight w (typically $w \in H^\infty$) and a symbol φ , according to $T_{w,\varphi} := M_w C_\varphi$. Often we will denote this operator by T for short.

Depending on φ , the operator C_φ can be compact or not. As we already said, passing from C_φ to $M_w C_\varphi$ can only improve this compactness, or the behavior of singular numbers, thanks to the ideal property of both notions. It is the purpose of this paper to investigate the question more closely.

2.2 A simple general upper bound

Throughout this section and the rest of this paper, we use the notation $A \lesssim B$ (resp. $A \gtrsim B$) to indicate that $A \leq \lambda B$ (resp. $A \geq \lambda B$) where λ is a uniform positive constant, "uniform" being clear from the context. Let φ be a symbol continuous on $\overline{\mathbb{D}}$, fixing 1, and $w_0 \in H^\infty$ a weight. We set $\gamma(t) = \varphi(e^{it})$. We assume that $\varphi(\mathbb{D})$ has no other contact points than 1 with \mathbb{T} and more precisely that:

$$(2.5) \quad |t| \leq \pi \Rightarrow 1 - |\gamma(t)| \geq \omega(|t|),$$

where $\omega : [0, \pi] \rightarrow \mathbb{R}^+$ is an increasing function with $\omega(0) = 0$. We also set

$$(2.6) \quad \delta_{w_0}(h) = \sup_{|t| \leq \omega^{-1}(h)} |w_0(\gamma(t))|.$$

We then have an upper bound for a special class of weights related to φ :

Theorem 2.2. *Let φ be a symbol satisfying (2.5), w_0 a weight, $w = w_0 \circ \varphi$, $T = M_w C_\varphi$, and $a_n = a_n(T)$. Then:*

$$(2.7) \quad a_n \lesssim \inf_{0 < h < 1} \left[e^{-nh} + \delta_{w_0}(h) \right] =: \rho_n.$$

Proof. The proof is close to that of [26, Thm. 5.1]. Let $f = z^n g \in z^n H^2 =: E$, a subspace of H^2 of codimension $n < n + 1$. Assume that $\|f\| = 1$, so that $\|g\| = 1$. We see that, given $0 < h < 1$:

$$\begin{aligned} \|T(f)\|^2 &= \int_{\mathbb{T}} |w_0(\varphi(u))|^2 |\varphi(u)|^{2n} |g(\varphi(u))|^2 dm(u) \\ &= \int_{\mathbb{D}} |w_0(z)|^2 |z|^{2n} |g(z)|^2 dm_\varphi(z) \\ &= \int_{(1-h)\overline{\mathbb{D}}} |w_0(z)|^2 |z|^{2n} |g(z)|^2 dm_\varphi(z) + \int_{\mathbb{D}} |z|^{2n} |g(z)|^2 d\mu_h(z) \\ &\lesssim (1-h)^{2n} + \|\mu_h\|_C \lesssim e^{-2nh} + \|\mu_h\|_C, \end{aligned}$$

where μ_h denotes the restriction (trace) of the measure $|w_0|^2 dm_\varphi$ to the annulus $A_h = \{z : 1-h < |z| < 1\}$. It remains to estimate $\|\mu_h\|_C$, which we do through Carleson's embedding theorem.

Since μ_h is carried by A_h , we can consider only boxes $S = S(\xi, r)$ with $0 < r \leq h$. Then

$$\begin{aligned} I_r &:= \int 1_S(u) d\mu_h(u) = \int 1_S(u) |w_0(u)|^2 dm_\varphi(u) \\ &= \int_{S \cap \varphi(\partial\mathbb{D})} |w_0(u)|^2 dm_\varphi(u) \leq \left(\sup_{S \cap \varphi(\partial\mathbb{D})} |w_0(u)|^2 \right) m_\varphi(S) \end{aligned}$$

since m_φ is carried by $\varphi(\partial\mathbb{D})$. Now, if $u = \varphi(e^{it}) \in S$ and $|t| \leq \pi$, then

$$\omega(|t|) \leq 1 - |\gamma(t)| \leq |\xi - \gamma(t)| \leq r,$$

that is $|t| \leq \omega^{-1}(r)$. Therefore, $I_r \leq \delta_{w_0}^2(r) \rho_\varphi(r) \lesssim r \delta_{w_0}^2(h)$, and by Carleson's theorem:

$$\|\mu_h\|_C \lesssim \sup_{0 < r < h} \frac{I_r}{r} \lesssim \delta_{w_0}^2(h),$$

giving $a_{n+1} \lesssim \rho_n$ in view of the alternative definition (2.3) of approximation numbers, and ending the proof after a change of $n + 1$ to n . \square

2.3 From non-compactness to compactness

It is known that twisting a non-compact composition operator C_φ with a multiplication operator M_w can result in $M_w C_\varphi$ being compact (see, for example, [19]). We now give a first application of Theorem 2.2, in which this effect is demonstrated in terms of explicit estimates on approximation numbers, which seems to be new.

Note that the compactness result of part *ii*) of the following theorem also follows by applying [19, Thm. 2.8] to the specified symbol φ and weight w .

Theorem 2.3. *Let $\varphi(z) = \frac{1+z}{2}$ and $w(z) = (1-z)^\alpha$, $\alpha > 0$. Then:*

- i) C_φ is non-compact and indeed $\|C_\varphi\|_e = \|C_\varphi\| = \sqrt{2}$.*
- ii) $T = M_w C_\varphi$ is compact and its approximation numbers verify*

$$a_n(T) \lesssim \left(\frac{\log n}{n}\right)^{\alpha/2}.$$

In particular, a weighted composition operator $T = M_w C_\varphi$ can be compact while its “compositional symbol” φ has no fixed point inside \mathbb{D} .

Proof. Recall that $\|C_\varphi\|_e =: \lim_{n \rightarrow \infty} a_n(C_\varphi)$ is the essential norm of C_φ . The first item *i*) is well-known ([34], see also [13]). For upper bounds, we may use Theorem 2.2 since $w = w_0 \circ \varphi$ where $w_0(z) = 2^\alpha (1-z)^\alpha$. Next, we observe that, for $|t| \leq \pi$: $1 - |\gamma(t)| = 1 - \cos(t/2) = 2 \sin^2(t/4) \geq \delta t^2$, so that, up to absolute constants, we are allowed to take $\omega(h) = h^2$ and $\omega^{-1}(h) = \sqrt{h}$ in Theorem 2.2. Since $|w_0(\gamma(t))| \lesssim |1 - \gamma(t)|^\alpha \leq |t|^\alpha$, this implies that $\delta_{w_0}(h) \lesssim h^{\alpha/2}$, and subsequently that

$$a_n(T) \lesssim \inf_{0 < h < 1} \left[e^{-nh} + \delta_{w_0}(h) \right] \lesssim \inf_{0 < h < 1} \left[e^{-nh} + h^{\alpha/2} \right] \lesssim \left(\frac{\log n}{n}\right)^{\alpha/2}$$

by taking $h = C \frac{\log n}{n}$ where C is a large numerical constant. This gives the claimed upper bound.

Finally, the fixed point of φ is 1 and $1 \notin \mathbb{D}$. □

Remark: The simple estimates given here are not sharp (Theorem 4.1 will give sharper results) and are just intended to show that multiplication by w can improve the decay of approximation numbers. In particular, the logarithmic factor can be dropped in the example of Thm. 2.3, and whereas the estimate in Theorem 2.3 *ii*) give membership in the Hilbert-Schmidt class for $\alpha > 1$, one actually has the following stronger statement.

Proposition 2.4. *The following are equivalent in the previous example:*

i) $M_w C_\varphi$ is Hilbert-Schmidt.

ii) $\alpha > 1/2$.

Proof. If $T = M_w C_\varphi$ and if (e_n) is the canonical basis of H^2 , we find that

$$\sum_{n=0}^{\infty} \|T(e_n)\|^2 = 2 \int_{\mathbb{T}} \frac{|w(e^{i\theta})|^2}{1 - \cos \theta} d\theta \approx \int_0^\pi \theta^{2\alpha-2} d\theta$$

and the latter integral is finite iff $\alpha > 1/2$. Alternatively, we could use $\|T(e_n)\| \approx n^{-\alpha/2-1/4}$. \square

2.4 Maximal general possible decay

It was proved in [26] that singular numbers of composition operators never have a superexponential decay. The same holds for weighted composition operators.

Theorem 2.5. *Let $T = M_w C_\varphi$ be a weighted composition operator. Then $\beta^-(T) > 0$, that is, there exist positive constants δ and ρ such that for any integer $n \geq 1$:*

$$a_n(T) \geq \delta \rho^n.$$

Proof. We first recall that the interpolation constant I_z of a (finite or not) sequence $z = (z_j)$ of distinct points of \mathbb{D} is the smallest constant K such that, for any bounded sequence $c = (c_j)$, one can find $h \in H^\infty$ such that

$$h(z_j) = c_j \quad \forall j \quad \text{and} \quad \|h\|_\infty \leq K \sup_j |c_j|.$$

The connection between interpolation constants and reproducing kernels is given by the well-known two-sided inequality [31, p. 302-303], valid for all scalars λ_j :

$$(2.8) \quad I_z^{-2} \sum_j |\lambda_j|^2 \|K_{z_j}\|^2 \leq \left\| \sum_j \lambda_j K_{z_j} \right\|^2 \leq I_z^2 \sum_j |\lambda_j|^2 \|K_{z_j}\|^2.$$

We shall now rely on the following lemma:

Lemma 2.6. *Let $u = (u_j)_{1 \leq j \leq n}$ be a sequence of length n of points of \mathbb{D} and $v = (v_j) = (\varphi(u_j))_{1 \leq j \leq n}$. We assume that the v_j 's (and hence the u_j 's) are distinct. Let I_v be the interpolation constant of v . Then if $T = M_w C_\varphi$:*

$$a_n(T) \gtrsim \left(\inf_{1 \leq j \leq n} |w(u_j)| \right) \times \left(\inf_{1 \leq j \leq n} \sqrt{1 - |u_j|^2} \right) \times I_v^{-2}.$$

Indeed, we use the ‘‘model space’’ E generated by the reproducing kernels K_{u_1}, \dots, K_{u_n} of H^2 as well as the mapping equation for weighted composition operators

$$(2.9) \quad T^*(K_a) = \overline{w(a)} K_{\varphi(a)},$$

a well-known and readily verified fact since for all $g \in H^2$:

$$\begin{aligned} \langle g, T^*(K_a) \rangle &= \langle Tg, K_a \rangle = \langle w(g \circ \varphi), K_a \rangle \\ &= w(a) g(\varphi(a)) = w(a) \langle g, K_{\varphi(a)} \rangle = \langle g, \overline{w(a)} K_{\varphi(a)} \rangle. \end{aligned}$$

Now, (2.9) and estimates analog to those of [27] prove the lemma. We provide some details. Let $f = \sum_{j=1}^n \lambda_j K_{u_j} \in S_E$, the unit sphere of E . So that, using (2.8) for the finite sequence $u = (u_j)_{1 \leq j \leq n}$:

$$(2.10) \quad 1 \leq I_u^2 \sum_{j=1}^n |\lambda_j|^2 \|K_{u_j}\|^2.$$

Similarly, since $T^*(f) = \sum_{j=1}^n \lambda_j \overline{w(u_j)} K_{v_j}$ where $v_j = \varphi(u_j)$, there holds:

$$(2.11) \quad \|T^*(f)\|^2 \geq I_v^{-2} \sum_{j=1}^n |\lambda_j|^2 |w(u_j)|^2 \|K_{v_j}\|^2.$$

Now, $I_u \leq I_v$ (if $h \in H^\infty$ interpolates c at v , $h \circ \varphi$ interpolates c at u), and clearly $\|K_{v_j}\|^2 \geq (1 - |u_j|^2) \|K_{u_j}\|^2$ for all j . Lemma 2.6 ensues via (2.2). \square

Finally, let $U \subset \mathbb{D}$ be a compact disk on which w does not vanish, let $\delta = \inf_U |w| > 0$ and V be a closed disk with positive radius contained in $\varphi(U)$. Let (v_j) be a sequence of n equidistributed points on ∂V and (u_j) a sequence of length n in U such that $\varphi(u_j) = v_j$. We know (see [18], p. 284) that $I_v \leq C^n$ where C only depends on V . We set $\eta^2 = \text{dist}(U, \partial \mathbb{D}) > 0$. Lemma 2.6 then gives us

$$a_n(T) \gtrsim \delta \times \eta \times C^{-2n},$$

which ends the proof of Thm. 2.5. \square

2.5 Maximal special possible decay

Theorem 2.5 proved that one never has superexponential decay for the approximation numbers $a_n(M_w C_\varphi)$. The following result indicates that, when $\|\varphi\|_\infty = 1$, then whatever the non-zero weight w , the numbers $a_n(M_w C_\varphi)$ indeed have at most subexponential decay.

Theorem 2.7. *Suppose that the symbol φ satisfies $\|\varphi\|_\infty = 1$. Let now $T = M_w C_\varphi$ where $w \in \mathcal{M}(H^2, \varphi)$, assumed to be compact. Then $\beta(T) = 1$, i.e. there exists a sequence (ε_n) of positive numbers with limit 0 such that*

$$a_n(C_\varphi) \geq e^{-n\varepsilon_n}.$$

Proof. We borrow from [28] some results on the Green capacity $\text{Cap}(X)$ of Borel subsets X of \mathbb{D} .

- $X \subset Y \Rightarrow \text{Cap}(X) \leq \text{Cap}(Y)$.
- $X_j \uparrow X \Rightarrow \text{Cap}(X_j) \uparrow \text{Cap}(X)$.
- $\text{Cap}(X) = \text{Cap}(\partial X) = \text{Cap}(\overline{X})$ when X is connected and $\overline{X} \subset \mathbb{D}$.
- When the X_j 's are connected, $\text{Cap}(X_j) \rightarrow \infty$ if $\text{diam } X_j \rightarrow 1$, where diam denotes the diameter of X_j for the pseudo-hyperbolic distance ρ in \mathbb{D} :

$$\rho(a, b) = \left| \frac{a - b}{1 - \overline{a}b} \right|.$$

We also set $\Gamma(X) = \exp(-1/\text{Cap}(X))$. With these notations, we will now prove the following extension of the main result of [28] to weighted composition operators, which implies Theorem 2.7.

Theorem 2.8. *Let $T = M_w C_\varphi$ where φ is an arbitrary symbol, and let $w \in \mathcal{M}(H^2, \varphi)$. Then, $\beta(T)$ exists and moreover*

$$\beta(T) = \Gamma(\varphi(\mathbb{D})).$$

In particular, $\|\varphi\|_\infty = 1 \Rightarrow \beta(T) = 1$.

For the upper bound we can assume (see the proof of the lower bound), that $\|\varphi\|_\infty < 1$, so that $C_\varphi : H^2 \rightarrow H^\infty$. Since $w \in \mathcal{M}(H^2, \varphi)$, we have $w \in H^2$ and $M_w : H^\infty \rightarrow H^2$ is bounded. By the ideal property of approximation numbers, we obtain

$$(2.12) \quad a_n(T) \leq \|M_w : H^\infty \rightarrow H^2\| \times a_n(C_\varphi : H^2 \rightarrow H^\infty).$$

But for each r with $\|\varphi\|_\infty < r < 1$, we can write $\varphi = \beta_r \circ \gamma_r$ where $\beta_r(z) = rz$ and $\gamma_r(z) = \frac{\varphi(z)}{r}$, so that $C_\varphi = C_{\gamma_r} C_{\beta_r}$, with $C_{\beta_r} : H^2 \rightarrow H^\infty$ and $C_{\gamma_r} : H^\infty \rightarrow H^\infty$, and (using again the ideal property of approximation numbers) $a_n(C_\varphi : H^2 \rightarrow H^\infty) \leq \|C_{\beta_r}\| \times a_n(C_{\gamma_r} : H^\infty \rightarrow H^\infty)$. Now, using a result of Widom (see [28], Theorem 3.6), we can assert that

$$(2.13) \quad \limsup_{n \rightarrow \infty} [a_n(C_{\gamma_r} : H^\infty \rightarrow H^\infty)]^{1/n} \leq \Gamma(\gamma_r(\mathbb{D})) = \Gamma\left(\frac{\varphi(\mathbb{D})}{r}\right).$$

This implies:

$$\limsup_{n \rightarrow \infty} [a_n(C_\varphi : H^2 \rightarrow H^\infty)]^{1/n} \leq \Gamma\left(\frac{\varphi(\mathbb{D})}{r}\right).$$

Letting r tend to 1^- gives $\limsup_{n \rightarrow \infty} [a_n(C_\varphi : H^2 \rightarrow H^\infty)]^{1/n} \leq \Gamma(\varphi(\mathbb{D}))$. Inserting this estimate in (2.12) finally gives

$$\limsup_{n \rightarrow \infty} [a_n(T)]^{1/n} \leq \Gamma(\varphi(\mathbb{D}))$$

or else $\beta^+(T) \leq \Gamma(\varphi(\mathbb{D}))$.

For the lower bound, we will make use of a second result of H. Widom (see [28]), in which K is a compact subset of \mathbb{D} of positive capacity, and $\|\cdot\|_{\mathcal{C}(K)}$ denotes the sup-norm on the space of continuous functions on K :

If E is a subspace of H^2 with $\dim E < n$, there exists $f \in B_{H^\infty}$, the unit ball of H^∞ , such that (a denoting a positive absolute constant)

$$(2.14) \quad \|f - h\|_{\mathcal{C}(K)} \geq a[\Gamma(K)]^n \text{ for all } h \in E.$$

Now, since w is not identically 0, we can find a sequence (r_j) with $r_j \uparrow 1$ such that $|z| = r_j \Rightarrow w(z) \neq 0$, implying

$$\delta_j := \inf_{|z|=r_j} |w(z)| > 0.$$

Set $K_j = \varphi(r_j\mathbb{T})$ and let $E \subset H^2$ with $\dim E < n$. By (2.14), we can find $f \in B_{H^\infty} \subset B_{H^2}$ such that, for all $h \in E$:

$$(2.15) \quad \|f - h\|_{\mathcal{C}(K_j)} \geq a[\Gamma(K_j)]^n.$$

It ensues that

$$\begin{aligned} \|w(f \circ \varphi - h \circ \varphi)\|_{\mathcal{C}(r_j\mathbb{T})} &\geq \left(\inf_{r_j\mathbb{T}} |w|\right) \left(\sup_{r_j\mathbb{T}} |f \circ \varphi - h \circ \varphi|\right) \\ &\geq \delta_j \|f - h\|_{\mathcal{C}(K_j)} \geq a\delta_j [\Gamma(K_j)]^n. \end{aligned}$$

Moreover, we obviously have the inequality $\|g\|_{\mathcal{C}(r_j\mathbb{T})} \leq L_j \|g\|_2$ for all functions $g \in H^2$, where the positive constant L_j only depends on j . This implies

$$a\delta_j [\Gamma(K_j)]^n \leq \|w(f \circ \varphi - h \circ \varphi)\|_{\mathcal{C}(r_j\mathbb{T})} \leq L_j \|w(f \circ \varphi - h \circ \varphi)\|_2 = L_j \|Tf - Th\|_2$$

so that, for some positive constant L'_j depending only on j :

$$d(Tf, TE) \geq L'_j [\Gamma(K_j)]^n.$$

Since this holds for every subspace E of H^2 with $\dim E < n$, we derive from (2.1) that $a_n(T) \geq L'_j [\Gamma(K_j)]^n$. Taking n th-roots and passing to \liminf as $n \rightarrow \infty$, we get

$$(2.16) \quad \beta^-(T) \geq \Gamma(K_j).$$

To finish, we set

$$\omega_j = r_j \mathbb{D}, \quad K'_j = \varphi(\overline{\omega_j}) = \overline{\varphi(\omega_j)} \supset K_j.$$

Clearly, $\text{Cap}(K'_j) \geq \text{Cap}(K_j)$. However, $\partial\varphi(\omega_j) \subset \varphi(\partial\omega_j) = K_j$ since $\varphi(\omega_j)$ is open. We hence get, using the reminded results on the capacity of connected sets:

$$\text{Cap}(K'_j) = \text{Cap}[\varphi(\omega_j)] = \text{Cap}[\partial\varphi(\omega_j)] \leq \text{Cap}(K_j).$$

So that $\text{Cap}(K'_j) = \text{Cap}(K_j)$ and that, using (2.16):

$$(2.17) \quad \beta^-(T) \geq \Gamma(K'_j).$$

Letting $j \rightarrow \infty$, we obtain, since $K'_j \uparrow \varphi(\mathbb{D})$: $\beta^-(T) \geq \Gamma(\varphi(\mathbb{D}))$ and hence $\beta(T) = \Gamma(\varphi(\mathbb{D}))$.

Now, suppose that $\|\varphi\|_\infty = 1$. By conformal invariance of the parameters involved ($\beta(C_\varphi)$, $\text{Cap}(\varphi(\mathbb{D}))$, $\text{diam}(\varphi(\mathbb{D}))$), we can assume that $\varphi(0) = 0$, without loss of generality. In that case, $\text{diam}(K'_j) \rightarrow 1$ and subsequently $\Gamma(K'_j) \rightarrow 1$. So that $\Gamma(\varphi(\mathbb{D})) = 1$, and finally $\beta(T) = 1$. \square

Remark. In the preceding, we limited ourselves to the case of the Hardy space, but several results, for example Theorems 2.5 and 2.7, hold true for other Hilbert spaces of analytic functions on the disk. Indeed, they hold true [28] if the ambient norm in H is defined by

$$\|f\|^2 = |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 \omega(z) \frac{dA(z)}{\pi}$$

where A is the area measure on \mathbb{C} and ω a radial weight on \mathbb{D} , integrable on $(0, 1)$. This framework includes for example the Hardy space ($\omega(r) = 1 - r$), the Bergman space ($\omega(r) = (1 - r)^2$), or the Dirichlet space \mathcal{D} ($\omega(r) = 1$), even though the multiplier space $\mathcal{M}(\mathcal{D})$ is not H^∞ . To prove these results, we can adapt the methods of [28]. We skip the details.

3 Modular theory and Hardy spaces on strips

3.1 From von Neumann algebras to Hardy spaces

We now begin our discussion about how Hardy spaces and composition operators arise in the context of (specific) inclusions of von Neumann algebras and their real standard subspaces.

The following definition is motivated by algebraic quantum field theory [20] (in two dimensions), where one builds models by assigning von Neumann algebras to regions (subsets) in \mathbb{R}^2 such that a number of geometric properties (inclusions of subsets, causal separation w.r.t. the Minkowski inner product, symmetries) are carried into corresponding algebraic properties (inclusions of algebras, commutants, group actions).

A region of particular interest is the *wedge* $W := \{x \in \mathbb{R}^2 : \pm x_\pm > 0\}$ (here $x = (x_+, x_-)$ is the parameterization of $x \in \mathbb{R}^2$ in light cone coordinates) [6, 5, 10]. The following definition, taken from [10], lists the essential properties of a von Neumann algebra \mathcal{N} (together with the space-time translations U and a vacuum vector Ω) that are required for \mathcal{N} to resemble the localization region W in a physically reasonable manner.

Definition 3.1. *A Borchers triple consists of a von Neumann algebra \mathcal{N} , acting on some Hilbert space \mathcal{H} , a cyclic and separating unit vector $\Omega \in \mathcal{H}$, and a unitary strongly continuous representation U of \mathbb{R}^2 such that*

- i) U has positive energy. That is, writing $U(x) = e^{ix_+P_+}e^{ix_-P_-}$, the two generators are positive, $P_\pm > 0$.*
- ii) $U(x)\Omega = \Omega$ for all $x \in \mathbb{R}^2$.*
- iii) $U(x)\mathcal{N}U(x)^{-1} \subset \mathcal{N}$ for $x_+ > 0, x_- < 0$.*

We will here be mostly interested with certain modular data derived from a Borchers triple, and now recall the relevant notions (see [30] for details and proofs of the claims made here).

Given a Borchers triple, we consider the closed real subspace

$$(3.1) \quad K := \{N\Omega : N = N^* \in \mathcal{N}\}^{\|\cdot\|} \subset \mathcal{H}$$

which is *standard* in the sense that $K + iK$ is dense in \mathcal{H} , and $K \cap iK = \{0\}$. To any such standard subspace one can associate a Tomita operator, that is the antilinear involution defined as

$$(3.2) \quad S : K + iK \rightarrow K + iK, \quad k_1 + ik_2 \mapsto k_1 - ik_2.$$

This operator is densely defined, closed, and typically unbounded. Its polar decomposition $S = J\Delta^{1/2}$ gives rise to a positive non-singular linear operator $\Delta^{1/2}$ with domain $\text{dom } \Delta^{1/2} = K + iK$ (the modular operator) and an anti unitary involution J (the modular conjugation), satisfying $J\Delta^{it} = \Delta^{it}J$, $t \in \mathbb{R}$, and $J\Omega = \Omega$, $\Delta^{1/2}\Omega = \Omega$.

The Hardy space (on a strip) that will arise later will be associated with the domain of the modular operator $\Delta^{1/2}$. As a first step towards this link, let us equip the dense subspace $\text{dom } \Delta^{1/2} \subset \mathcal{H}$ with the graph norm, defined by the scalar product

$$(3.3) \quad \langle \psi, \varphi \rangle_{\Delta} := \langle \psi, \varphi \rangle + \langle \Delta^{1/2}\psi, \Delta^{1/2}\varphi \rangle = \langle \psi, (1 + \Delta)\varphi \rangle.$$

Since $\Delta^{1/2}$ is a closed operator, $\text{dom } \Delta^{1/2}$ is closed in the graph norm [33] $\|\cdot\|_{\Delta} = \langle \cdot, \cdot \rangle_{\Delta}^{1/2}$. That is, $(\text{dom } \Delta^{1/2}, \langle \cdot, \cdot \rangle_{\Delta})$ is a complex Hilbert space. When considering $\text{dom } \Delta^{1/2}$ with this scalar product, we refer to it as \mathcal{D} .

The following proposition gathers basic information on the modular data from this point of view.

Proposition 3.2. *Let K be a closed real standard subspace and \mathcal{D} the complex Hilbert space defined as $\text{dom } \Delta^{1/2}$ with its graph scalar product (3.3).*

- i) On \mathcal{D} , the Tomita operator S is an anti unitary involution.*
- ii) On \mathcal{D} , the modular unitaries Δ^{it} , $t \in \mathbb{R}$, are still unitary.*
- iii) For $0 \leq \mu \leq \frac{1}{2}$, the modular operator Δ^{μ} is a linear bounded operator of norm at most one when viewed as a map from \mathcal{D} to \mathcal{H} .*
- iv) The modular conjugation J is an antiunitary operator $\mathcal{D}_{\Delta} \rightarrow \mathcal{D}_{\Delta^{-1}}$.*

Proof. *i)* Let $\psi, \varphi \in \mathcal{D}$. Then, since $S^*S = \Delta$ and $\Delta^{1/2}S = \Delta^{1/2}J\Delta^{1/2} = J$,

$$\begin{aligned} \langle S\psi, S\varphi \rangle_{\Delta} &= \langle S\psi, S\varphi \rangle + \langle \Delta^{1/2}S\psi, \Delta^{1/2}S\varphi \rangle = \overline{\langle \psi, \Delta\varphi \rangle} + \langle J\psi, J\varphi \rangle \\ &= \overline{\langle \Delta^{1/2}\psi, \Delta^{1/2}\varphi \rangle} + \langle \psi, \varphi \rangle = \overline{\langle \psi, \varphi \rangle_{\Delta}}. \end{aligned}$$

This shows that S is anti unitary on \mathcal{D} .

ii) This is clear because Δ^{it} commutes with $\Delta^{1/2}$.

iii) For $\psi \in \mathcal{D}$, the \mathcal{H} -valued function $z \mapsto \Delta^{-iz}\psi$ is holomorphic on the strip $0 < \text{Im}(z) < \frac{1}{2}$, continuous on its closure, and norm-constant in the real direction, $\|\Delta^{-i(x+iy)}\psi\| = \|\Delta^y\psi\|$. Thus the three lines theorem [15] implies, $0 \leq y \leq \frac{1}{2}$,

$$\|\Delta^y\psi\|^{1/2} \leq \|\psi\|^{1/2-y} \cdot \|\Delta^{1/2}\psi\|^y,$$

from which we read off

$$\|\Delta^y\psi\| \leq \|\psi\|^{1-2y} \cdot \|\Delta^{1/2}\psi\|^{2y} \leq \sqrt{\|\psi\|^2 + \|\Delta^{1/2}\psi\|^2} = \|\psi\|_{\Delta}.$$

iv) This follows from, $\psi, \varphi \in \text{dom}(\Delta^{1/2})$,

$$\begin{aligned} \langle J\psi, J\varphi \rangle_{\Delta} &= \langle J\psi, J\varphi \rangle + \langle \Delta^{1/2}J\psi, \Delta^{1/2}J\varphi \rangle \\ &= \overline{\langle \psi, \varphi \rangle} + \overline{\langle \Delta^{-1/2}\psi, \Delta^{-1/2}\varphi \rangle} = \overline{\langle J\psi, J\varphi \rangle_{\Delta^{-1}}}. \end{aligned}$$

□

To draw the connection to Hardy spaces, we recall a theorem of Borchers [6] on the commutation relation of the modular unitaries Δ^{it} , $t \in \mathbb{R}$, and the translation unitaries $U(x)$, $x \in \mathbb{R}^2$: If (\mathcal{N}, U, Ω) is a Borchers triple, then there holds

$$(3.4) \quad \Delta^{it}U(x)\Delta^{-it} = U(\Lambda(t)x), \quad JU(x)J = U(-x), \quad x \in \mathbb{R}^2,$$

where $(\Lambda(t)x)_{\pm} = e^{\mp 2\pi t}x_{\pm}$.

The commutation relations (3.4) imply that in the presence of a Borchers triple, the operators $U(x)$, Δ^{it} , and J generate a (anti-)unitary strongly continuous representation of the proper Poincaré group $\mathcal{P}_+ = \text{SO}(1, 1) \rtimes \mathbb{R}^2$. We will denote this extended representation by the same letter U .

As the basic building blocks, we are interested in the irreducible sub-representations of U . We call a representation degenerate if there exists a non-zero vector Ψ which is invariant under $U(x)$ for all $x \in \mathbb{R}^2$, i.e. $\Psi \in \ker P_+ \cap \ker P_-$. In physical models, this vector can usually only be a multiple of Ω , so that we may restrict to non-degenerate representations of \mathcal{P}_+ .

The non-degenerate, irreducible, unitary, strongly continuous positive energy representations of \mathcal{P}_+ can be classified up to unitary equivalence according to the joint spectrum of the generators P_{\pm} , denoted $\text{Sp}(U|_{\mathbb{R}^2})$. There are equivalence classes of three types:

- m) $\text{Sp}(U|_{\mathbb{R}^2}) = \{p \in \mathbb{R}^2 : p_{\pm} > 0, p_+ \cdot p_- = m^2\}$ for some $m > 0$,
- $0+$) $\text{Sp}(U|_{\mathbb{R}^2}) = \{p \in \mathbb{R}^2 : p_+ \geq 0, p_- = 0\}$,
- $0-$) $\text{Sp}(U|_{\mathbb{R}^2}) = \{p \in \mathbb{R}^2 : p_- \geq 0, p_+ = 0\}$.

The parameter m has the physical interpretation of a mass. We therefore refer to representations of type $m > 0$ as “massive”, and to representations of type $0\pm$ as “massless”. For a concise notation, we will adopt the convention to label objects by a single label m , which can either take positive values, referring to type m), or the two special values $m = 0\pm$, referring to type $0\pm$).

The irreducible representation U_m of type m can be conveniently realized on the Hilbert space $\mathcal{H} = L^2(\mathbb{R}, d\theta)$. In fact, we have for all types the same modular unitaries and conjugation [23]

$$(3.5) \quad \mathcal{H} = L^2(\mathbb{R}, d\theta), \quad (\Delta^{it}\psi)(\theta) = \psi(\theta - 2\pi t), \quad (J\psi)(\theta) = \overline{\psi(\theta)}.$$

The translation operators are multiplication operators depending on the type, namely

$$(3.6) \quad (U_m(x)\psi)(\theta) = w_{m,x}(\theta) \cdot \psi(\theta),$$

$$(3.7) \quad w_{0\pm,x}(\theta) = e^{ix_{\pm} e^{\pm\theta}}, \quad w_{m,x}(\theta) = e^{im(x_+ e^{\theta} + x_- e^{-\theta})}.$$

The functions $w_{m,x}$ will later serve as the weights of our weighted composition operators.

Considering the representation (3.5), it becomes clear that the domain $\mathcal{D} \subset L^2(\mathbb{R})$ of $\Delta^{1/2}$ consists of functions that have an analytic continuation to the strip region $\mathbb{S}_{0,\pi}$, a special case of the more general strip

$$(3.8) \quad \mathbb{S}_{a,b} := \{\zeta \in \mathbb{C} : a < \text{Im} \zeta < b\}, \quad a < b,$$

and satisfy certain bounds on this strip. Before we make this precise, let us recall some properties of functions analytic in a strip, and corresponding function spaces.

Given $f \in \text{Hol}(\mathbb{S}_{a,b})$ (the holomorphic functions $\mathbb{S}_{a,b} \rightarrow \mathbb{C}$), we write f_{λ} , $a < \lambda < b$, for its restriction to the line $\mathbb{R} + i\lambda$. Denoting the usual norm of $L^2(\mathbb{R})$ by $\|\cdot\|_2$, we consider the norm

$$(3.9) \quad \|f\|_{\mathbb{S}_{a,b}} := \sup_{a < \lambda < b} \|f_{\lambda}\|_2 \in [0, +\infty]$$

and set

$$(3.10) \quad H_B^2(\mathbb{S}_{a,b}) := \{f \in \text{Hol}(\mathbb{S}_{a,b}) : \|f\|_{\mathbb{S}_{a,b}} < \infty\}.$$

We recall the following facts [37]:

- i) $(H_B^2(\mathbb{S}_{a,b}), \|\cdot\|_{\mathbb{S}_{a,b}})$ is a Banach space.
- ii) Any $f \in H_B^2(\mathbb{S}_{a,b})$ has L^2 -boundary values on the two boundaries $\mathbb{R} + ia$ and $\mathbb{R} + ib$, i.e. $f_{a+\varepsilon}$ and $f_{b-\varepsilon}$ converge in $L^2(\mathbb{R})$ as $\varepsilon \searrow 0$. By a slight abuse of notation, we will denote these boundary values as $f_a, f_b \in L^2(\mathbb{R})$.
- iii) As an expression of the maximum principle, the function $(a, b) \ni \lambda \mapsto \|f_\lambda\|_2$ is logarithmically convex for $f \in H_B^2(\mathbb{S})$. In particular, $\|f\| = \max\{\|f_a\|_2, \|f_b\|_2\}$.

We will refer to $H_B^2(\mathbb{S}_{a,b})$ as *Hardy Banach space* to distinguish it from a Hardy Hilbert space on $\mathbb{S}_{a,b}$ to be introduced next.

Indeed, as the strip is an unbounded region, there exist two different types of Hardy Hilbert spaces for this domain: The conformally invariant Hardy space, defined in terms of harmonic majorants, and the not conformally invariant Hardy space, defined in terms of L^2 -integrals over a sequence of Jordan curves tending to the boundary of the strip [17, Ch. 10].

For our purposes, only the latter space will be relevant. It will be convenient to characterize it in terms of a Riemann map $\tau : \mathbb{D} \rightarrow \mathbb{S}$. (To lighten our notation, we write \mathbb{S} instead of $\mathbb{S}_{a,b}$ when the boundaries of the strip are arbitrary.) Namely, we define

$$(3.11) \quad H^2(\mathbb{S}) := \{f \in \text{Hol}(\mathbb{S}) : \sqrt{\tau'} \cdot (f \circ \tau) \in H^2(\mathbb{D})\}.$$

This is a Hilbert space with scalar product

$$(3.12) \quad \langle f, g \rangle_{\mathbb{S}} := \langle \sqrt{\tau'} \cdot (f \circ \tau), \sqrt{\tau'} \cdot (g \circ \tau) \rangle_{\mathbb{D}},$$

and $(H^2(\mathbb{S}), \langle \cdot, \cdot \rangle_{\mathbb{D}})$ depends on the choice of τ only up to changing the norm $\|f\|_{\mathbb{S}} := \langle f, f \rangle_{\mathbb{S}}^{1/2}$ to an equivalent Hilbert norm. We may therefore fix τ , and make the choice

$$(3.13) \quad \tau : \mathbb{D} \rightarrow \mathbb{S}_{a,b}, \quad \tau(z) := \frac{2(b-a)}{\pi} \operatorname{arctanh}(z) + \frac{i}{2}(a+b).$$

This is a biholomorphic mapping $\tau : \mathbb{D} \rightarrow \mathbb{S}_{a,b}$, and elementary calculations show that it has inverse and derivative, $\zeta \in \mathbb{S}_{a,b}, z \in \mathbb{D}$,

$$(3.14) \quad \tau^{-1}(\zeta) = \tanh\left(\frac{\pi}{2(b-a)}\zeta - \frac{i\pi}{4}\frac{b+a}{b-a}\right), \quad \tau'(z) = \frac{2(b-a)}{\pi} \frac{1}{1-z^2},$$

$$(3.15) \quad (\tau^{-1})'(\zeta) = \frac{\pi}{2(b-a)} \frac{1}{\cosh^2\left(\frac{\pi}{2(b-a)}\zeta - \frac{i\pi}{4}\frac{b+a}{b-a}\right)}.$$

Proposition 3.3. *i) $H_B^2(\mathbb{S})$ and $H^2(\mathbb{S})$ coincide as linear spaces.*

ii) The two norms $\|\cdot\|$ and $\|\cdot\|_{\mathbb{S}_{a,b}}$ are equivalent: For any $f \in \text{Hol}(\mathbb{S})$, there holds

$$(3.16) \quad \frac{1}{\sqrt{2\pi}} \|f\| \leq \|f\|_{\mathbb{S}} \leq \frac{1}{\sqrt{\pi}} \|f\|.$$

iii) The scalar product of $H^2(\mathbb{S}_{a,b})$ can be written as

$$(3.17) \quad \langle f, g \rangle_{\mathbb{S}_{a,b}} = \frac{1}{2\pi} \left(\langle f_a, g_a \rangle_2 + \langle f_b, g_b \rangle_2 \right), \quad f, g \in H^2(\mathbb{S}_{a,b}).$$

Proof. We first work on the special strip \mathbb{S} given by $a = -1$, $b = 1$, and introduce for $f \in \text{Hol}(\mathbb{S})$ the notation $s(f) := \frac{1}{2} \sup_{0 \leq y < 1} (\|f_y\|_2^2 + \|f_{-y}\|_2^2) \in [0, +\infty]$.

It was shown in [3, Thm. 2.1 & 2.2] that for $f \in \text{Hol}(\mathbb{S})$, one has $f \circ \tau \in H^2(\mathbb{D})$ if and only if $s(w \cdot f) < \infty$, and in this case, $\|f \circ \tau\|_{\mathbb{D}}^2 = s(w \cdot f)$, with the weight $w(\zeta) = (2 \cosh \frac{\pi\zeta}{4})^{-1}$.

The condition that some $f \in \text{Hol}(\mathbb{S})$ lies in $H^2(\mathbb{S})$, i.e. that $\sqrt{\tau'}(f \circ \tau) = (f/\sqrt{(\tau^{-1})'}) \circ \tau \in H^2(\mathbb{D})$, is therefore equivalent to $s(w/\sqrt{(\tau^{-1})'} \cdot f) < \infty$. But in view of (3.15), $w(\zeta)/\sqrt{(\tau^{-1})'(\zeta)} = \pi^{-1/2}$. We thus have that $f \in \text{Hol}(\mathbb{S})$ lies in $H^2(\mathbb{S})$ if and only if $s(f) < \infty$, and in this case,

$$(3.18) \quad \|f\|_{\mathbb{S}}^2 = \|\sqrt{\tau'}(f \circ \tau)\|_{\mathbb{D}}^2 = \frac{1}{\pi} s(f) = \frac{1}{2\pi} \sup_{0 \leq y < 1} (\|f_y\|_2^2 + \|f_{-y}\|_2^2).$$

As the supremum on the right hand side clearly lies between $\|f\|^2$ and $2\|f\|^2$, the claimed equivalence of norms in *ii)* follows. This also implies *i)*.

To establish *iii)*, we use that $(-1, 1) \ni y \mapsto \|f_y\|_2$ is logarithmically convex for $f \in H_B^2(\mathbb{S})$. Thus $y \mapsto \|f_y\|_2^2 + \|f_{-y}\|_2^2$ is convex, which implies that the supremum in (3.18) is taken for the boundary values at $y = 1$, i.e.

$$\|f\|_{\mathbb{S}}^2 = \frac{1}{2\pi} (\|f_1\|_2^2 + \|f_{-1}\|_2^2) = \frac{1}{2\pi} \left(\langle f_1, f_1 \rangle_2 + \langle f_{-1}, f_{-1} \rangle_2 \right).$$

This implies *iii)*.

It remains to proceed from $\mathbb{S}_{-1,1}$ to a general strip $\mathbb{S}_{a,b}$ by means of the variable transformation $\varphi : \mathbb{S}_{-1,1} \rightarrow \mathbb{S}_{a,b}$, $\varphi(\zeta) := \frac{b-a}{2}\zeta + \frac{i}{2}(a+b)$. But by elementary substitutions, one finds that $f \mapsto f \circ \varphi$ is a bijection $H_B^2(\mathbb{S}_{a,b}) \rightarrow H_B^2(\mathbb{S}_{-1,1})$, with $\|(f \circ \varphi)_y\|_2^2 = \frac{2}{b-a} \|f_{(b-a)y/2}\|_2^2$. Since also $\|f \circ \varphi\|_{\mathbb{S}_{-1,1}}^2 = \frac{2}{b-a} \|f\|_{\mathbb{S}_{a,b}}^2$, the properties *i)–iii)* follow from the special case $a = -1$, $b = 1$. \square

Given the action of the unitaries Δ^{it} (3.5) in the irreducible representations U_m arising from the modular data of our Borchers triple, the scalar product (3.17) of $H^2(0, \pi)$ is strongly reminiscent of the graph scalar product (3.3). To make this match exact, we will use a rescaled version of the graph scalar product, namely

$$(3.19) \quad \langle \psi, \varphi \rangle_{\Delta} := \frac{1}{2\pi} \left(\langle \psi, \varphi \rangle + \langle \Delta^{1/2} \psi, \Delta^{1/2} \varphi \rangle \right).$$

Clearly, Proposition 3.2 still holds with this equivalent scalar product. Moreover, we have the following concrete realization of \mathcal{D} .

Proposition 3.4. *Consider the modular data (3.5), and denote by \mathcal{D} the complex Hilbert space $\text{dom } \Delta^{1/2}$ with scalar product (3.19).*

- i) $\mathcal{D} = H^2(\mathbb{S}_{0,\pi})$ as complex Hilbert spaces.*
- ii) The Tomita operator S acts on $H^2(\mathbb{S}_{0,\pi})$ by “crossing symmetry”, i.e.*

$$(3.20) \quad (S\psi)(\zeta) = \overline{\psi(i\pi + \bar{\zeta})}, \quad \zeta \in \mathbb{S}_{0,\pi}.$$

Proof. It was shown in [23, Lemma A.1] that the real standard subspace $K = \ker(1 - J\Delta^{1/2})$ is given by

$$K = \{ \psi \in H_B^2(\mathbb{S}_{0,\pi}) : \overline{\psi_{\pi}(\theta)} = \psi_0(\theta) \text{ a.e.} \}.$$

In view of the analyticity properties of ψ , this implies that K consists exactly of those functions $\psi \in H_B^2(\mathbb{S}_{0,\pi})$ that satisfy

$$(3.21) \quad \overline{\psi(i\pi + \bar{\zeta})} = \psi(\zeta), \quad \zeta \in \mathbb{S}_{0,\pi}.$$

Clearly any $f \in H_B^2(\mathbb{S}_{0,\pi})$ can be written as $f = \psi + i\varphi$ with $\psi, \varphi \in K$, so that we see $H_B^2(\mathbb{S}_{0,\pi}) = K + iK$. But as S is an antilinear involution, with domain $\mathcal{D} = K + iK$, and $H_B^2(\mathbb{S}_{0,\pi}) = H^2(\mathbb{S}_{0,\pi})$, it follows that $\mathcal{D} = H^2(\mathbb{S}_{0,\pi})$ as linear spaces. Also the graph scalar product (3.19) coincides with the scalar product of $H^2(\mathbb{S}_{0,\pi})$ by Prop. 3.3 *iii*). This shows *i*).

The Tomita operator S is uniquely fixed by being an antilinear involution and $Sk = k$ for all $k \in K$. But in view of the characterization (3.21) of K , it is clear that the antilinear involution defined in (3.20) leaves K pointwise invariant. This shows *ii*). \square

Having established the connection between modular data and Hardy spaces, we now explain how composition operators appear in this setting.

Any Borchers triple defines a quantum field theory on \mathbb{R}^2 [10], which makes this concept interesting in the context of constructing models. However, the quantum field theories arising from Borchers triples might be pathological in the sense of containing no strictly local observables, a situation that arises when the inclusions $U(x)\mathcal{N}U(x)^{-1} \subset \mathcal{N}$, $x \in W$, have trivial relative commutants. These pathological situations can however be ruled out [9, 22] when the so-called *modular nuclearity condition* [8, 7] holds. This condition requires that the maps

$$(3.22) \quad \Xi_{x,\mu} : \mathcal{N} \rightarrow \mathcal{H}, \quad \Xi_{x,\mu}N := \Delta^\mu U(x)N\Omega, \quad x \in W, \quad 0 < \mu < \frac{1}{2},$$

are *nuclear*² as linear maps between the two Banach spaces $(\mathcal{N}, \|\cdot\|_{\mathcal{B}(\mathcal{H})})$ and \mathcal{H} . Whereas $\Xi_{x,\mu}$ is always bounded by modular theory, it is in general not compact, so that nuclearity of (3.22) is a non-trivial requirement.

To investigate the approximation numbers of $\Xi_{x,\mu}$, we split this map as

$$(3.23) \quad \Xi_{x,\mu} : \mathcal{N} \xrightarrow{Y} \mathcal{D} \xrightarrow{U(x)} \mathcal{D} \xrightarrow{\Delta^\mu} \mathcal{H},$$

where the first operator, defined as $Y(N) := N\Omega$, is bounded: For any $N \in \mathcal{N}$, we have

$$\begin{aligned} \|Y(N)\|_{\Delta}^2 &= \|N\Omega\|^2 + \langle \Delta^{1/2}N\Omega, \Delta^{1/2}N\Omega \rangle = \|N\Omega\|^2 + \langle JN^*\Omega, JN^*\Omega \rangle \\ &\leq 2\|N\|^2, \end{aligned}$$

because $\|J\| = 1$ and $\|N^*\| = \|N\|$.

The last operator in (3.23), Δ^μ , is bounded as an operator $\mathcal{D} \rightarrow \mathcal{H}$ (see Prop. 3.2 *iii*)).

Lemma 3.5. *Let (\mathcal{N}, U, Ω) be a Borchers triple. Then the translations $U(x)$, $x \in W$, are isometries as maps on the Hilbert space \mathcal{D} .*

Proof. It is known that the commutation relations (3.4) imply that the operator $\Delta^{1/2}U(x)\Delta^{-1/2}$, $x \in W$, is defined on $\text{dom } \Delta^{-1/2}$, and coincides there with $JU(x)J = U(-x)$ [30, Thm. 2.3.1 f)]. We therefore find for $\psi \in \mathcal{D}$ and $x \in W$ the equation $\Delta^{1/2}U(x)\psi = \Delta^{1/2}U(x)\Delta^{-1/2}\Delta^{1/2}\psi = U(-x)\Delta^{1/2}\psi$, and consequently

$$\begin{aligned} \|U(x)\psi\|_{\Delta}^2 &= \|U(x)\psi\|^2 + \|\Delta^{1/2}U(x)\psi\|^2 \\ &= \|U(x)\psi\|^2 + \|U(-x)\Delta^{1/2}\psi\|^2 = \|\psi\|_{\Delta}^2, \end{aligned}$$

²The condition that a linear map X between two Banach spaces is nuclear is slightly weaker than X having summable approximation numbers [32].

where we have used that $U(x)$ is unitary on \mathcal{H} . This shows that $U(x)$ is an isometry on \mathcal{D} . Note that $U(x)$ is (except for trivial cases) not unitary because it does not have full range. \square

The product of the last two operators in the split (3.23),

$$(3.24) \quad D_{x,\mu} : \mathcal{D} \rightarrow \mathcal{H}, \quad D_{x,\mu} = \Delta^\mu U(x),$$

can however be compact (and even have approximation number that go to zero quite fast), analogously to the situation encountered in Thm 2.3.

To obtain estimates on the approximation numbers $a_n(D_{x,\mu})$, one splits \mathcal{H} and \mathcal{D} into irreducible subspaces of U . Then each subspace takes the form $\mathcal{H} = L^2(\mathbb{R}, d\theta)$, $\mathcal{D} = H^2(\mathbb{S}_{0,\pi})$, and $U_m(x) : \mathcal{D} \rightarrow \mathcal{D}$ acts by multiplication with the (analytic continuation of the) weight $w_{m,x}$ (3.6), depending on the representation type m . (Note that the analytically continued weight functions $w_{m,x} \in H^\infty(\mathbb{S}_{0,\pi})$ (3.6) are bounded and inner, for any m and x .)

Explicitly, the operator $D_{x,\mu}^{(m)} = D_{x,\mu}$ then takes the concrete form of a “weighted restriction operator”, $0 < \mu < \pi$, $x \in W$,

$$(3.25) \quad D_{x,\mu}^{(m)} : H^2(\mathbb{S}_{0,\pi}) \rightarrow L^2(\mathbb{R}),$$

$$(3.26) \quad (D_{x,\mu}^{(m)} \psi)(\theta) := w_{m,x}(\theta + i\mu) \cdot \psi(\theta + i\mu).$$

This observation warrants a more systematic analysis of weighted restriction operators on Hardy spaces on strips. Before we enter into this analysis in the next section, let us comment on the relation between the operators $D_{x,\mu}^{(m)}$ and $\Xi_{x,\mu}$ (3.22).

Estimates on the approximation numbers of $D_{x,\mu}^{(m)}$ do not imply corresponding estimates on the maps $\Xi_{x,\mu}$ (3.22). To establish bounds on the $a_n(\Xi_{x,\mu})$, one also has to take into account the multiplicities occurring in the decomposition of U into irreducibles. But for typical examples of Borchers triples, this analysis involves basically only (symmetrized) tensor powers of operators of the form $D_{x,\mu}^{(m)}$ [1]. For this reason, it is of interest to determine strong decay properties of the approximation numbers of $D_{x,\mu}^{(m)}$.

3.2 Weighted restriction operators on $H^2(\mathbb{S})$

As a slight generalization of what appeared before, we consider here the following setting: Let $\mathbb{S} \subset \mathbb{C}$ be a strip domain, which will be fixed in the following. To define the operators we want to study, we take a narrower strip $\hat{\mathbb{S}} \subset \mathbb{S}$, such that the closure of the smaller strip is contained in the

larger one, and a weight function $w \in H^\infty(\mathbb{S})$. We are then interested in the mappings $f \mapsto (w \cdot f)|_{\tilde{\mathbb{S}}}$, considered as operators $H^2(\mathbb{S}) \rightarrow H^2(\tilde{\mathbb{S}})$.

For simplicity, we will always assume that both \mathbb{S} and $\tilde{\mathbb{S}}$ are symmetric around the real axis, i.e. $\mathbb{S} = \mathbb{S}_{-b,b}$ for some $b > 0$ and $\tilde{\mathbb{S}} = \lambda\mathbb{S}$ for some $0 < \lambda < 1$. We then define

$$(3.27) \quad R_{w,\lambda} : H^2(\mathbb{S}) \rightarrow H^2(\lambda\mathbb{S}), \quad R_{w,\lambda}f := (w \cdot f)|_{\lambda\mathbb{S}}.$$

As a limiting case as $\lambda \rightarrow 0$, we also define

$$(3.28) \quad R_{w,0} : H^2(\mathbb{S}) \rightarrow L^2(\mathbb{R}), \quad R_{w,0} := (w \cdot f)_0$$

as the restriction of wf to the real line. It is clear from Prop. 3.3 *i*) and the form of the norm (3.9) that the restriction maps $R_{1,\lambda}$, $0 \leq \lambda < 1$, with trivial weight $w = 1$ are bounded. Since restriction of $f \in H^2(\mathbb{S})$ to $\lambda'\mathbb{S}$ is the same as first restricting f to $\lambda\mathbb{S}$, $\lambda > \lambda'$, and then to $\lambda'\mathbb{S}$, we find that there is a constant c such that

$$(3.29) \quad a_n(R_{w,\lambda'}) \leq c a_n(R_{w,\lambda}), \quad 0 \leq \lambda' \leq \lambda \leq 1, \quad n \in \mathbb{N}.$$

In particular, the operators $R_{w,0}$ mapping to $L^2(\mathbb{R})$ (3.28) can be estimated in terms of the $R_{w,\lambda}$, $\lambda > 0$. The latter operators map between Hardy spaces and can be reformulated as composition operators as follows.

Let $0 < \lambda < 1$ and

$$(3.30) \quad L_\lambda : H^2(\lambda\mathbb{S}) \rightarrow H^2(\mathbb{S}), \quad (L_\lambda f)(z) := \sqrt{\lambda}f(\lambda z).$$

Taking into account that the Riemann maps τ_λ for $\lambda\mathbb{S}$ and τ for \mathbb{S} are related by $\tau_\lambda = \lambda\tau$, it follows that L_λ is unitary. Furthermore, the product $\lambda^{-1/2} L_\lambda R_{1,\lambda}$ is easily seen to be the composition operator C_λ on $H^2(\mathbb{S})$ with linear symbol $z \mapsto \lambda z$. This shows that the restriction operators $R_{1,\lambda}$ are unitarily similar to composition operators. Furthermore, we note that C_λ (and thus $R_{1,\lambda}$) are *not* compact. This is so because on $H^2(\mathbb{S})$ there exist no compact composition operators at all [36]. The *weighted* operators $R_{w,\lambda}$ can however be compact, depending on the weight w , similar to the example in Thm. 2.3.

For the following analysis of the approximation numbers of $R_{w,\lambda}$, we recall that $H^2(\mathbb{S})$ is a reproducing kernel Hilbert space. Its kernel function $K_{\mathbb{S}}$ is related to the well-known Szegő kernel [35] $K_{\mathbb{D}}(z, z') = (1 - \bar{z}z')^{-1}$ of $H^2(\mathbb{D})$ by

$$(3.31) \quad K_{\mathbb{S}}(\zeta, \zeta') = \sqrt{(\tau^{-1})'(\zeta)} \cdot K_{\mathbb{D}}(\tau^{-1}(\zeta), \tau^{-1}(\zeta')) \cdot \sqrt{(\tau^{-1})'(\zeta')}.$$

To compute this explicitly for the strip $\mathbb{S} = \mathbb{S}_{-b,b}$, we insert (3.14) and get

$$(3.32) \quad K_{\mathbb{S}}(\zeta, \zeta') = \frac{\pi}{4b} \frac{1}{\cosh \frac{\pi(\bar{\zeta} - \zeta')}{4b}}.$$

We also recall that given any orthonormal basis $\{\psi_n\}_n$ of $H^2(\mathbb{S})$, we have $\sum_n \overline{\psi_n(\zeta)} \psi_n(\zeta') = K_{\mathbb{S}}(\zeta, \zeta')$. So, in particular,

$$(3.33) \quad \sum_n |\psi_n(\zeta)|^2 = \frac{\pi}{4b} \frac{1}{\cos \frac{\pi \operatorname{Im}(\zeta)}{2b}}.$$

Proposition 3.6. *Let $w \in H^\infty(\mathbb{S})$ and $0 \leq \lambda < 1$.*

i) If $w|_{\lambda\mathbb{S}} \in H^2(\lambda\mathbb{S})$ (for $\lambda > 0$) or $w_0 \in L^2(\mathbb{R})$ (for $\lambda = 0$), then $R_{w,\lambda}$ is Hilbert-Schmidt, with Hilbert-Schmidt norm

$$(3.34) \quad \|R_{w,\lambda}\|_2 = \sqrt{\frac{\pi}{4b \cos \frac{\pi\lambda}{2}}} \cdot \|w\|_{\lambda\mathbb{S}}, \quad \lambda > 0, \quad \|R_{w,0}\|_2 = \sqrt{\frac{\pi}{4b}} \cdot \|w\|_2.$$

ii) If w is non-vanishing and rapidly decreasing in the sense that for any $k \in \mathbb{N}$,

$$(3.35) \quad \sup_{\substack{\theta \in \mathbb{R} \\ -\lambda b \leq \mu \leq \lambda b}} (|w(\theta + i\mu)| (1 + \theta^2)^k) < \infty,$$

then the approximation numbers of $R_{w,\lambda}$ satisfy for any $N \in \mathbb{N}$

$$(3.36) \quad \sup_{n \in \mathbb{N}} (n^N a_n(R_{w,\lambda})) < \infty.$$

iii) If $w(\theta) \rightarrow c$, $c \neq 0$, as $\theta \rightarrow \infty$ or $\theta \rightarrow -\infty$, then $R_{w,\lambda}$ is not compact.

Proof. *i)* Let $\{\psi_n\}_n$ be some orthonormal basis of $H^2(\mathbb{S})$, and $0 < \lambda < 1$. Then, using (3.33),

$$\begin{aligned} \sum_n \|R_{w,\lambda} \psi_n\|_{\lambda\mathbb{S}}^2 &= \frac{1}{2\pi} \sum_n \int_{\mathbb{R}} d\theta (|w_{-\lambda b}(\theta)|^2 |\psi_{n,-\lambda b}(\theta)|^2 + |w_{\lambda b}(\theta)|^2 |\psi_{n,\lambda b}(\theta)|^2) \\ &= \frac{1}{8b} \int_{\mathbb{R}} d\theta \left(\frac{|w(\theta - i\lambda b)|^2}{\cos \frac{\pi\lambda}{2}} + \frac{|w(\theta + i\lambda b)|^2}{\cos \frac{\pi\lambda}{2}} \right) \\ &= \frac{\pi}{4b \cos \frac{\pi\lambda}{2}} \cdot \|w\|_{\lambda\mathbb{S}}^2. \end{aligned}$$

Since $\|R_{w,\lambda}\|_2^2 = \text{Tr}(R_{w,\lambda}^* R_{w,\lambda})$, this finishes the proof for $\lambda > 0$. The argument for $\lambda = 0$ is analogous.

ii) By (3.29), it is sufficient to show the claim for $\lambda > 0$. Let $k \in \mathbb{N}$. We find intermediate strip regions $\mathbb{S} \supset \lambda_1 \mathbb{S} \supset \lambda_2 \mathbb{S}_2 \supset \dots \supset \lambda_{k-1} \mathbb{S} \supset \lambda \mathbb{S}$ such that in each inclusion, the closure of the smaller strip is contained in the larger strip. In view of the assumption on w , we may furthermore write our weight as a product $w = w_1 \cdot w_2 \cdots w_k$, with $w_1, \dots, w_k \in H^2(\mathbb{S})$ (We can take $w_j := w^{1/k}$ for $j = 1, \dots, k$.) Thus our operator can be written as

$$R_{w,\lambda} = R_{\lambda/\lambda_{k-1}, w_k} \cdot R_{\lambda_{k-1}/\lambda_{k-2}, w_{k-1}} \cdots R_{\lambda_1, w_1}.$$

By part *i)*, each of the k factors is Hilbert-Schmidt. That is, $R_{w,\lambda}$ can be written as a product of an arbitrary number of Hilbert-Schmidt operators. This implies the claim by standard estimates on approximation numbers [32].

iii) We consider the case that $w(\theta) \rightarrow c \neq 0$ as $\theta \rightarrow +\infty$, the opposite limit is analogous. By (3.29), it is sufficient to consider the case $\lambda = 0$.

For non-zero $f \in H^2(\mathbb{S})$, we consider the sequence $f_n(\zeta) := f(\zeta - n)$. To show that $R_{w,0}$ is not compact, we show that $\{R_{w,0} f_n\}_n$ has no convergent subsequence. In fact, by dominated convergence we have

$$\|R_{w,0} f_n\|_2^2 = \int_{\mathbb{R}} d\theta |w(\theta)|^2 |f(\theta - n)|^2 \rightarrow c^2 \|f_0\|_2^2.$$

Since $c \neq 0$ and $f \neq 0$, this limit is non-zero, i.e. $\|R_{w,0} f_n\|_2 \geq c_0 > 0$ for sufficiently large n . On the other hand, we have

$$|\langle R_{w,0} f_{n_1}, R_{w,0} f_{n_2} \rangle_2| \leq \|w\|_\infty^2 \int_{\mathbb{R}} d\theta |f(\theta)| \cdot |f(\theta + n_1 - n_2)|,$$

and this converges to 0 for $n_1 - n_2 \rightarrow \infty$ by the falloff properties of Hardy space functions. Thus for large $n_1, n_2, |n_1 - n_2|$, the vectors $R_{w,0} f_{n_1}, R_{w,0} f_{n_2}$ have approximately identical non-zero length and are approximately orthogonal to each other. Thus $\{R_{w,0} f_n\}_n$ can have no convergent subsequence. \square

The situations described in item *ii)* and *iii)* of this proposition fit to the weights appearing in the massive and massless irreducible Poincaré representations, introduced in the previous section. To see this, we need to translate the weights $w_{m,x} \in H^\infty(\mathbb{S}_{0,\pi})$ (3.6) to the symmetric strip $\mathbb{S}_{-\pi/2,\pi/2}$ by shifting their argument $\zeta \rightarrow \zeta + \frac{i\pi}{2}$. In the massive case $m > 0$, this results in the weight (denoted by the same symbol)

$$(3.37) \quad w_{m,x}(\zeta) = e^{m(-x_+ e^\zeta + x_- e^{-\zeta})}, \quad \zeta \in \mathbb{S}_{-\pi/2,\pi/2}.$$

Taking into account that $\pm x_{\pm} > 0$, it is apparent that w is rapidly decreasing on any strip of the form $\lambda\mathbb{S}$, $0 < \lambda < 1$. Thus, by part *ii*) of the preceding proposition, the approximation numbers of $R_{w,\lambda}$ are rapidly decreasing for any $0 \leq \lambda < 1$.

In the massless case, however, the weight becomes

$$(3.38) \quad w_{0\pm,x}(\zeta) = e^{\mp x_{\pm} e^{\pm\zeta}}, \quad \zeta \in \mathbb{S}_{-\pi/2,\pi/2},$$

which converges to 1 as $\zeta \rightarrow \mp\infty$. Thus, by item *iii*), the weighted restriction operator $R_{w,0}$ is not compact, which fits with the known result that in the massless case, $\Xi_{x,\mu}$ is not compact.

The estimates of Prop. 3.6 *ii*) are rather imprecise, and can be improved for specific weights w . To derive sharper bounds, let us switch to the unit disc \mathbb{D} and make contact with the setting of Section 2.

By the very definition of the scalar product $\langle \cdot, \cdot \rangle_{\mathbb{S}}$, the mapping

$$(3.39) \quad V : H^2(\mathbb{S}) \rightarrow H^2(\mathbb{D}), \quad V\psi := \sqrt{\tau'} \cdot (\psi \circ \tau),$$

is unitary, with inverse $V^{-1}f = \sqrt{(\tau^{-1})'} \cdot (f \circ \tau^{-1})$. For the scaled strip $\lambda\mathbb{S}$, $0 < \lambda < 1$, we have the Riemann map $\tau_{\lambda} := \lambda\tau$, with corresponding unitary V_{λ} . With the help of these unitaries, we can transfer the operators $R_{w,\lambda}$ to the Hardy space $H^2(\mathbb{D})$ on the disc.

Let us introduce some notation first. We define $\varphi_{\lambda} = g^{-1} \circ \gamma_{\lambda} \circ g$, with $0 < \lambda < 1$ and

$$g(z) = \frac{1+z}{1-z}, \quad \gamma_{\lambda}(u) = u^{\lambda} \text{ for } \operatorname{Re} u > 0$$

as the lens map of parameter λ . Explicitly,

$$(3.40) \quad \varphi_{\lambda}(z) = \frac{(1+z)^{\lambda} - (1-z)^{\lambda}}{(1+z)^{\lambda} + (1-z)^{\lambda}}.$$

Lemma 3.7. *The operator $R_{w,\lambda}$, $0 < \lambda < 1$, is unitarily similar to a weighted lens map composition operator $M_{\hat{w}}C_{\varphi_{\lambda}}$ on $H^2(\mathbb{D})$, with lens map symbol φ_{λ} (3.40), and weight*

$$(3.41) \quad \hat{w}_{\lambda}(z) := \sqrt{\lambda} m_{\lambda}(z) w(\lambda\tau(z)), \quad m_{\lambda}(z) := \left(\frac{1 - \varphi_{\lambda}(z)^2}{1 - z^2} \right)^{1/2}.$$

Proof. The disc operator unitarily similar to $R_{w,\lambda}$ is $V_\lambda R_{w,\lambda} V^{-1} : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$. Inserting the definitions yields

$$\begin{aligned} (V_\lambda R_{w,\lambda} V^{-1} f)(z) &= \sqrt{\lambda \tau'(z)} \sqrt{(\tau^{-1})'(\lambda \tau(z))} \cdot w(\lambda \tau(z)) \cdot f(\tau^{-1}(\lambda \tau(z))) \\ &= \sqrt{\lambda} \sqrt{\frac{\tau'(z)}{\tau'(\tau^{-1}(\lambda \tau(z)))}} \cdot w(\lambda \tau(z)) \cdot f(\tau^{-1}(\lambda \tau(z))). \end{aligned}$$

We note that

$$(3.42) \quad \tau^{-1}(\lambda \cdot \tau(z)) = \tanh(\lambda \cdot \operatorname{arctanh}(z)) = \varphi_\lambda(z)$$

is the lens map. Inserting the explicit form of τ' (3.14) then yields the claimed result. \square

The weight (3.41) consists of the unbounded universal factor m_λ (which diverges as $z \rightarrow \pm 1$) and the bounded factor $w \circ \lambda \tau$. For the factor m_λ , it is easy to show that

$$(3.43) \quad |m_\lambda(z)| \leq 2 |1 - z^2|^{\frac{1}{2}(\lambda-1)}, \quad z \in \mathbb{D}.$$

In the case of the “massive weight” (3.37), we obtain after a short calculation

$$(3.44) \quad w(\lambda \tau(z)) \leq e^{-(s_+ g(z)^\lambda + s_- g(-z)^\lambda)},$$

with the parameters $s_\pm := \frac{m\pi}{2} |x_\pm| > 0$. Whereas this function decreases to zero quite fast as z approaches ± 1 , the two contact points of the lens map with the boundary of \mathbb{D} , the function m_λ only diverges mildly at these points. For the purposes of estimating approximation numbers, we may therefore reduce the parameters s_\pm a little to compensate the divergent factor m_λ , and consider weighted lens map composition operators with a weight of the form (3.44) instead. Such operators will be studied in the following section.

To conclude this section, we also transfer the operators $R_{w,0}$ (3.28), corresponding to $\lambda = 0$, to the disc. Whereas for $\lambda > 0$, we obtained weighted composition operators on the disc, the case $\lambda = 0$ corresponds to Carleson embeddings.

To see this, let us define on \mathbb{D} the measure

$$(3.45) \quad \mu_w(x + iy) := |\hat{w}_1(x)|^2 \delta(y) dx dy, \quad w_1(z) := w\left(\frac{4b}{\pi} \operatorname{arctanh} z\right),$$

supported on the real diameter of the disc.

Lemma 3.8. *Let $w \in H^\infty(\mathbb{S}_{-b,b})$.*

- i) The measure μ_w (3.45) is a Carleson measure, i.e. $H^2(\mathbb{D}) \subset L^2(\mathbb{D}, d\mu_w)$, and the embedding $J_w : H^2(\mathbb{D}) \hookrightarrow L^2(\mathbb{D}, d\mu_w)$ is bounded.*
- ii) The operator $R_{w,0}$ is unitarily similar to the embedding J_w .*

Proof. *i)* Since the weight \hat{w}_1 is bounded on \mathbb{D} , we can estimate the measure of a Carleson box as, $\xi \in \mathbb{T}$, $0 < h \leq 1$,

$$\mu[S(\xi, h)] \leq \mu[S(1, h)] = \int_{1-h}^h |\hat{w}_1(x)|^2 dx \leq \|\hat{w}_1\|_\infty^2 \cdot h$$

for $\operatorname{Re} \xi \geq 0$, and analogously for $\operatorname{Re} \xi < 0$. Thus μ is a Carleson measure (see Thm. 2.1), which implies the remaining statements in *i*).

ii) We define the unitaries

$$(3.46) \quad V_0 : L^2(\mathbb{R}) \rightarrow L^2((-1, 1), dx), \quad (V_0\psi)(x) := \frac{\psi\left(\frac{4b}{\pi} \operatorname{arctanh}(x)\right)}{\sqrt{\frac{\pi}{4b}(1-x^2)}}$$

and

$$(3.47) \quad \tilde{V} : L^2((-1, 1), dx) \rightarrow L^2(\mu), \quad (\tilde{V}\psi)(x) := \frac{\psi(x)}{\hat{w}_1(x)}.$$

Then inserting the definitions yields, $f \in H^2(\mathbb{D})$,

$$\begin{aligned} (\tilde{V}V_0R_{w,0}V^{-1}f)(x) &= \frac{1}{\hat{w}_1(x)} \frac{w\left(\frac{4b}{\pi} \operatorname{arctanh}(x)\right) (V^{-1}f)\left(\frac{4b}{\pi} \operatorname{arctanh}(x)\right)}{\sqrt{\frac{\pi}{4b}(1-x^2)}} \\ &= f(x). \end{aligned}$$

□

4 Weighted lens map composition operators

We now turn to weighted *lens map* composition operators on $H^2(\mathbb{D})$ as particular examples of weighted composition operators. As in the previous section, we define the analytic map $\varphi = \varphi_\lambda = g^{-1} \circ \gamma_\lambda \circ g$, with $0 < \lambda < 1$ and

$$g(z) = \frac{1+z}{1-z}, \quad \gamma_\lambda(u) = u^\lambda \text{ for } \operatorname{Re} u > 0$$

as the lens map of parameter λ . Explicitly,

$$\varphi(z) = \frac{(1+z)^\lambda - (1-z)^\lambda}{(1+z)^\lambda + (1-z)^\lambda}.$$

Recall that the image $\varphi(\mathbb{D}) \subset \mathbb{D}$ has exactly two non-tangential contact points with the unit circle at 1 and -1 , and in particular verifies $\|\varphi\|_\infty = 1$. Also observe that $\varphi(-z) = -\varphi(z)$ for all $z \in \mathbb{D}$.

We choose a weight of the form motivated by the applications outlined in the previous section, namely $w = w_0 \circ \varphi$ where ³

$$(4.1) \quad w_0(z) = \exp \left[- \left(\frac{1+z}{1-z} \right)^\lambda \right] \exp \left[- \left(\frac{1-z}{1+z} \right)^\lambda \right] =: w_1(z)w_{-1}(z)$$

We have $\max(\|w_1\|_\infty, \|w_{-1}\|_\infty) \leq 1$, and note that w_0 tends to 0 quite rapidly as $z \rightarrow \pm 1$. Namely, if $z \in \mathbb{D}$:

$$(4.2) \quad \operatorname{Re} z \geq 0 \Rightarrow |w_0(z)| \leq |w_1(z)| \leq \exp \left(- \frac{\delta}{|1-z|^\lambda} \right)$$

$$(4.3) \quad \operatorname{Re} z \leq 0 \Rightarrow |w_0(z)| \leq |w_{-1}(z)| \leq \exp \left(- \frac{\delta}{|1+z|^\lambda} \right)$$

where $\delta = \cos(\lambda\pi/2)$. Indeed, if for example $\operatorname{Re} z \geq 0$, we see that

$$\operatorname{Re} \left(\frac{1+z}{1-z} \right)^\lambda \geq \delta \left| \frac{1+z}{1-z} \right|^\lambda \quad \text{with } |1+z| \geq \operatorname{Re}(1+z) \geq 1.$$

We take for w_0 this double product to take both contact points ± 1 of $\varphi(\mathbb{D})$ with the unit circle into account. We note in passing (this is general) that

$$C_\varphi M_{w_0} = M_{w_0 \circ \varphi} C_\varphi = M_w C_\varphi.$$

We can now state one of our main theorems. The positive constants $0 < c < C$ are allowed to change from one line to another in what follows.

Theorem 4.1. *Let $T = M_w C_\varphi : H^2 \rightarrow H^2$ where w and φ are as above. Then T is compact, and more precisely:*

- i) *One nearly has exponential decay, namely $a_n(T) \leq C e^{-c \frac{n}{\log n}}$.*
- ii) *The previous estimate is optimal: $a_n(T) \geq c e^{-C \frac{n}{\log n}}$.*

³In comparison to (3.44), we have set the inessential parameters s_\pm to 1.

Proof. We begin with the upper bound, following the strategy of [25]. If $\gamma(t) = \varphi(e^{it})$, one easily checks [25, Lemma 2.5] that

$$(4.4) \quad |t| \leq \frac{\pi}{2} \Rightarrow 1 - |\gamma(t)| \approx |1 - \gamma(t)| \approx |t|^\lambda$$

$$(4.5) \quad \frac{\pi}{2} \leq |t| \leq \pi \Rightarrow 1 - |\gamma(t)| \approx |1 + \gamma(t)| \approx (\pi - |t|)^\lambda.$$

We fix an integer $N \geq 2$. Let B be the Blaschke product

$$\begin{aligned} B(z) &= \prod_{1 \leq k \leq \log N} \left(\frac{z - p_k}{1 - \overline{p_k}z} \frac{z - \overline{p_k}}{1 - p_k z} \right)^N \prod_{1 \leq k \leq \log N} \left(\frac{z + p_k}{1 + \overline{p_k}z} \frac{z + \overline{p_k}}{1 + p_k z} \right)^N \\ &=: B_1(z) B_{-1}(z) \end{aligned}$$

where $p_k = \gamma(t_k)$ and $t_k = \frac{\pi}{2} 2^{-(k-1)/\lambda}$. This is a Blaschke product of length $\leq 4N \log N$. Observe that $\overline{B_{\pm 1}(z)} = B_{\pm 1}(\overline{z})$. Next, set $E = BH^2$, a subspace of codimension $< [4N \log N + 1] =: M$ (with $[\cdot]$ the integer part). If $f = Bg \in E$, with $\|f\| = \|g\| = 1$, we have (remembering (2.4) and since $w = w_0 \circ \varphi$)

$$\|T(f)\|^2 = \int_{\mathbb{T}} |w_0(\varphi(u))|^2 |B(\varphi(u))|^2 |g(\varphi(u))|^2 dm(u) = \int_{\mathbb{D}} |w_0|^2 |B|^2 |g|^2 dm_\varphi.$$

Now, using again Carleson's embedding theorem for the measure

$$\mu = |w_0|^2 |B|^2 dm_\varphi,$$

as in the proof of Theorem 2.2, and the non-tangential behavior of φ near the points ± 1 (allowing us to ignore the Carleson windows centered elsewhere than in ± 1 as in [25, p. 809]), we get with help of (2.3):

$$a_M(T)^2 \lesssim \sup_{0 < h < 1} \left(\frac{I_1(h)}{h} + \frac{I_{-1}(h)}{h} \right)$$

where

$$\begin{aligned} I_1(h) &= \int_{|\gamma(t)-1| \leq Ch} |B_1(\gamma(t))|^2 |w_1(\gamma(t))|^2 dt \\ I_{-1}(h) &= \int_{|\gamma(t)+1| \leq Ch} |B_{-1}(\gamma(t))|^2 |w_{-1}(\gamma(t))|^2 dt. \end{aligned}$$

Actually, the term $I_1(h)$ takes care of the Carleson boxes $S(\xi, h)$ for which $\operatorname{Re} \xi \geq 0$ and the term $I_{-1}(h)$ of those for which $\operatorname{Re} \xi < 0$. We will estimate

only $I_1(h)$, the estimate being similar for $I_{-1}(h)$.

By interpolation, we can assume $h = 2^{-q/\lambda}$ with q a non-negative integer, and we separate two cases:

Case 1: $q > \log N$. Then, majorizing $|B_1(\gamma(t))|$ by 1 and using (4.2) as well as (4.4), we get

$$\begin{aligned} h^{-1}I_1(h) &\lesssim h^{-1} \int_{|t|^\lambda \leq Ch} |w_1(\gamma(t))|^2 dt \lesssim h^{-1} \int_{|t|^\lambda \leq Ch} \exp\left(-\frac{C}{|t|^{\lambda^2}}\right) dt \\ &\lesssim h^{\frac{1}{\lambda}-1} \exp\left(-\frac{C}{h^\lambda}\right) \lesssim \exp(-C2^q) \lesssim \exp(-CN). \end{aligned}$$

Case 2: $q \leq \log N$. Then, we majorize $|w_1(\gamma(t))|$ by 1 and estimate $|B_1(\gamma(t))|$ more accurately, with help of an obvious modification of Lemma 2.6 in [25], which we recall:

Lemma 4.2. *Set $t_k = \frac{\pi}{2} 2^{-(k-1)/\lambda}$. Then*

$$t_{\lfloor \log N \rfloor} \leq t \leq t_1 \Rightarrow |B_1(\gamma(t))| \leq \chi^N$$

where $\lfloor \cdot \rfloor$ denotes the integer part and where $\chi < 1$ only depends on λ .

We can now finish the estimate (note that $t_1 = \pi/2$):

$$h^{-1}I_1(h) \lesssim h^{-1} \int_0^{t_{\lfloor \log N \rfloor}} |w_1(\gamma(t))|^2 dt + h^{-1} \int_{t_{\lfloor \log N \rfloor}}^{t_1} |B_1(\gamma(t))|^2 dt.$$

The first term is estimated as in Case 1. And since $q \leq \log N$, the second sum is dominated by $2^{q/\lambda} \chi^{2N} \lesssim N^{1/\lambda} \exp(-CN) \lesssim \exp(-C'N)$. Putting both cases together, we get $\sup_{0 < h < 1} h^{-1}I_1(h) \lesssim \exp(-CN)$ and finally, with $M = \lfloor 4N \log N + 1 \rfloor$ and using once more (2.3):

$$a_M^2 \lesssim e^{-CN}.$$

Inverting, interpolating, and using that $(a_p)_{p \geq 1}$ is non-increasing, we finally get for all positive integers n :

$$a_n \lesssim e^{-C \frac{n}{\log n}}$$

as claimed. This ends the proof of the upper bound.

To establish the lower bound, we first observe that

$$w_0(z) = \exp(-\gamma_\lambda g(z)) \exp(-\gamma_\lambda g(-z)).$$

Since $\gamma_\lambda \gamma_\lambda = \gamma_{\lambda^2}$, we get $\gamma_\lambda g \varphi = \gamma_{\lambda^2} g$, and an explicit formula for w is

$$(4.6) \quad w(z) = \exp \left[- \left(\frac{1+z}{1-z} \right)^{\lambda^2} \right] \exp \left[- \left(\frac{1-z}{1+z} \right)^{\lambda^2} \right] =: w'_1(z) w'_{-1}(z).$$

We now apply Lemma 2.6. To that effect, we must make a good choice of the u_j 's. As in [27] for lens maps, we choose $u_j = 1 - e^{-j\varepsilon}$ where $\varepsilon > 0$ has to be adjusted, and $v_j = \varphi(u_j)$. We know from [26, Lemma 6.5] that $I_v \leq \exp(C/\varepsilon)$, and we have $\sqrt{1 - u_j^2} \geq c e^{-n\varepsilon}$. Moreover, since

$$\left(\frac{1+u_j}{1-u_j} \right)^{\lambda^2} \leq \left(\frac{2}{1-u_j} \right)^{\lambda^2},$$

we see that

$$\inf_{1 \leq j \leq n} |w'_1(u_j)| \geq \exp(-C e^{n\varepsilon}).$$

And clearly $\inf_{1 \leq j \leq n} |w'_{-1}(u_j)| \geq e^{-1}$. So that $\inf_{1 \leq j \leq n} |w(u_j)| \geq \exp(-C e^{n\varepsilon})$ (recall that $w = w'_1 w'_{-1}$). Lemma 2.6 now gives us

$$a_n(T) \gtrsim \exp \left[- C \left(e^{n\varepsilon} + n\varepsilon + \frac{1}{\varepsilon} \right) \right] \gtrsim \exp \left[- C \left(e^{n\varepsilon} + \frac{1}{\varepsilon} \right) \right].$$

We finally adjust $\varepsilon = \frac{1}{2} \frac{\log n}{n}$ to get

$$a_n(T) \gtrsim \exp \left[- C \left(\sqrt{n} + \frac{n}{\log n} \right) \right] \gtrsim \exp \left[- C \frac{n}{\log n} \right].$$

This ends the proof of Theorem 4.1. □

Acknowledgements: G. Lechner would like to thank O. Bandtlow for discussions and pointing out the article [25], which initiated this collaboration. L. Rodríguez-Piazza was also supported by the research project MTM2015-63699-P (Spanish MINECO and FEDER funds).

References

- [1] S. Alazzawi and G. Lechner. Inverse Scattering and Locality in Integrable Quantum Field Theories. *Comm. Math. Phys.* 354(3), 913-956, 2017
- [2] K. Attele. Multipliers of the range of composition operators. *Tokyo. J. Math.*, 15:185–198, 1992.
- [3] A. Bakan and S. Kaijser. Hardy spaces for the strip. *J. Math. Anal. Appl.*, 333(1):347–364, 2007.

- [4] O. Bandtlow. Resolvent estimates for operators belonging to exponential classes. *Integr. Equ. Oper. Theory*, 61(1):21–43, 2008.
- [5] H. Baumgärtel and M. Wollenberg. Causal Nets of Operator Algebras. *Akademie Verlag*, 1992.
- [6] H.-J. Borchers. The CPT theorem in two-dimensional theories of local observables. *Comm. Math. Phys.*, 143:315–332, 1992.
- [7] D. Buchholz, C. D’Antoni, and R. Longo. Nuclear Maps and Modular Structures 2: Applications to Quantum Field Theory. *Comm. Math. Phys.*, 129:115, 1990.
- [8] D. Buchholz, C. D’Antoni, and R. Longo. Nuclear maps and modular structures. I. General properties. *J. Funct. Anal.*, 88:233–250, 1990.
- [9] D. Buchholz and G. Lechner. Modular nuclearity and localization. *Ann. Henri Poincaré*, 5:1065–1080, 2004.
- [10] D. Buchholz, G. Lechner, and S. J. Summers. Warped Convolutions, Rieffel Deformations and the Construction of Quantum Field Theories. *Comm. Math. Phys.*, 304:95–123, 2011.
- [11] B. Carl and I. Stephani. Entropy, Compactness and the Approximation of Operators. *Cambridge University Press*, 1990.
- [12] I. Chalendar, E. Gallardo-Gutiérrez, and J. Partington. Weighted composition operators on the Dirichlet space. *J. Math. Anal. Appl.*, 305:183–196, 2005.
- [13] J. Clifford and M. Dabkowski. Singular values and Schmidt pairs of composition operators on the Hardy space: boundedness and spectral properties. *Math. Ann.*, 3-4:1265–1279, 2015.
- [14] M. Contreras and A. Hernández-Diáz. Weighted composition operators between different Hardy spaces. *Integr. Equ. Oper. Theory*, 46:165–188, 2003.
- [15] J.B. Conway. Functions of One Complex Variable. *Springer*, 1978.
- [16] C. Cowen and B. MacCluer. Composition Operators on Spaces of Analytic Functions. *CRC Press*, 1994.
- [17] P. L. Duren. Theory of H^p Spaces. Dover Books on Mathematics. *Dover Publications, Inc., New York*, 1970.
- [18] J. Garnett. Bounded analytic functions. *Springer*, 2007.
- [19] E. Gallardo, R. Kumar, and J. Partington. Boundedness, Compactness and Schatten-class membership of weighted composition operators. *Integr. Equ. Oper. Theory* 67 , 467-479, 2010.
- [20] R. Haag. Local Quantum Physics - Fields, Particles, Algebras. *Springer, second edition*, 1996.

- [21] S. Hyvärinen, M. Lindström, I. Nieminen, and E. Saukko. Spectra of weighted composition operators with automorphic symbols. *J. Funct. Anal.*, 265:1749–1777, 2013.
- [22] G. Lechner. Construction of Quantum Field Theories with Factorizing S-Matrices. *Comm. Math. Phys.*, 277:821–860, 2008.
- [23] G. Lechner and Roberto Longo. Localization in Nets of Standard Spaces. *Comm. Math. Phys.*, 336(1):27–61, 2015.
- [24] P. Lefèvre. Generalized Essential Norm of Weighted Composition Operators on some Uniform Algebras of Analytic Functions. *Integr. Equ. Oper. Theory*, 63:557–569, 2009.
- [25] P. Lefèvre, D. Li, H. Queffélec, and L. Rodríguez-Piazza. Some new properties of composition operators associated with lens maps. *Israel J. Math.*, 195:801–824, 2013.
- [26] D. Li, H. Queffélec, and L. Rodríguez-Piazza. On approximation numbers of composition operators. *J. Approx. Theory*, 164(4):431–459, 2012.
- [27] D. Li, H. Queffélec, and L. Rodríguez-Piazza. Estimates for approximation numbers of some classes of composition operators on the Hardy space. *Ann Acad. Sci. Fenn. Math.*, 38:547–564, 2013.
- [28] D. Li, H. Queffélec, and L. Rodríguez-Piazza. A spectral radius formula for approximation numbers of composition operators. *J. Funct. Anal.* 268 (2): 4753-4774, 2015.
- [29] D. Li, H. Queffélec, and L. Rodríguez-Piazza. Some examples of composition operators and their approximation numbers on the Hardy space of the bidisk. *Preprint*, arXiv:1706.03570, submitted.
- [30] R. Longo. Lectures on Conformal Nets - Part 1. In *Von Neumann algebras in Sibiu*, pages 33–91. Theta, 2008.
- [31] N. K. Nikolskii. Operators, functions and systems: An Easy Reading, Volume 1. In *Math. Surveys and Monographs, Vol. 92, Amer. Math. Soc., Providence, RI*, 2002.
- [32] A. Pietsch. Nuclear Locally Convex Spaces. *Springer*, 1972.
- [33] M. Reed and B. Simon. Methods of Modern Mathematical Physics I - Functional Analysis. *Academic Press*, 1972.
- [34] J. H. Shapiro. The essential norm of a composition Operator. *Annals of Math.* 125, 375-404, 1987.
- [35] J. H. Shapiro. Composition Operators and Classical Function Theory. *Springer*, 1993.
- [36] J. H. Shapiro and W. Smith. Hardy spaces that support no compact composition operators. *J. Funct. Anal.*, 205(1):62–89, 2003.

- [37] E. M. Stein and G. Weiss. Introduction to Fourier Analysis on Euclidean Spaces. *Princeton University Press*, 1971.
- [38] M. Takesaki. Theory of Operator Algebras II. *Springer*, 2003.