

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/108860/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Perry, Amy, Gordon-Smith, K., Jones, L. and Jones, Ian 2017. A visual timeline tool for tracking mood and medication perinatally in affective disorders. *Archives of Women's Mental Health* 20 (5), pp. 709-710. 10.1007/s00737-017-0743-8 file

Publishers page: <http://dx.doi.org/10.1007/s00737-017-0743-8> <<http://dx.doi.org/10.1007/s00737-017-0743-8>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



A visual timeline tool for tracking mood and medication perinatally in affective disorders

Amy Perry, K. Gordon-Smith, L. Jones & I. Jones

Women with bipolar disorder are at particularly high risk of relapse following childbirth with almost one in five experiencing an episode of postpartum psychosis following delivery (Di Florio et al. 2013; Wesseloo et al. 2016). Factors which increase risk for severe postpartum relapse in women with bipolar disorder are poorly understood.

The majority of previous research has traditionally employed retrospective methods of assessment, and/or has only measured a limited range of potential risk factors. High-quality studies investigating risk factors prospectively in pregnant women with bipolar disorder are rare, and have predominantly focused on the role of medication use in postnatal relapse (Viguera et al. 2007).

We are currently conducting a large-scale prospective pregnancy study to investigate factors which influence risk for severe postpartum mood episodes in women with bipolar disorder (UK Bipolar Disorder Research Network, www.BDRN.org). Women are recruited systematically through specialist perinatal psychiatry services nationwide, or non-systematically via national patient support charities. Lifetime and current psychopathology is assessed during pregnancy using a semi structured interview (Schedules for Clinical Assessment in Neuropsychiatry; Wing et al. 1990), and a follow-up SCAN interview is conducted at 12-week postpartum to assess psychopathology during the perinatal period.

These data are combined with further information obtained at 12-week postpartum from clinician questionnaires and medical case records to determine both the occurrence of episodes of illness and medication use across the whole perinatal period. Data are also gathered on a wide range of potential risk and protective factors including obstetric complications, sleep loss, psychosocial factors relating to pregnancy (social support for example), comorbid physical and psychiatric conditions, menstrual history, and major adverse life events occurring lifetime ever and in the perinatal period specifically. These data are highly complex and multifaceted; there is considerable variation not only between participants but also individually throughout the perinatal period. For research, but also in clinical practice, there is a need to summarize this complex information and we have therefore developed a visual timeline tool as a means of graphically representing these detailed psychiatric and medication data longitudinally and simply across the perinatal period. Figure 1 provides an example timeline generated from data collected during the second pregnancy of a participant with DSM-5 (American Psychiatric Association 2013) bipolar I disorder. This woman, who had a previous episode of postpartum psychosis, took quetiapine prophylactically throughout her pregnancy. She experienced an episode of major depression at 24 weeks gestation. Following an increase in the dose of quetiapine, she recovered and remained euthymic throughout the rest of the pregnancy. The quetiapine dose was increased again at 32 weeks of pregnancy and she delivered at 37 weeks. At 24 days postpartum, she developed an episode of postpartum psychosis requiring compulsory admission to a psychiatric mother and baby unit and the addition of further psychotropic medication.

Perinatal Timeline (weeks)




Fig. 1 A timeline representing course of illness and medication use during the perinatal period for a participant recruited to the Bipolar Disorder Research Network (BDRN) prospective pregnancy study.

We have found this timeline tool to be a simple and efficient method of collating and presenting large amounts of complex clinical data for each perinatal period. Presenting these data graphically has allowed us to visually identify potential patterns in the data that are taken forward for formal testing. In addition to its use in research, we anticipate this tool would be equally useful in clinical practice for reviewing a patient's perinatal psychiatric history.

Acknowledgements The authors would like to thank the Stanley Medical Research Institute for funding this research and all members of the Bipolar Disorder Research Network (www.bdrn.org). We also wish to express our gratitude to all the individuals who kindly gave their time to participate in this research.

Compliance with ethical standards

Funding This study is funded by the Stanley Medical Research Institute (grant number 5710002223).

Conflict of interest The authors declare that they have no conflict of interest.

References

American Psychiatric Association (2013) Diagnostic and Statistical Manual of Mental Disorders, 5th edn. American Psychiatric Publishing, Arlington, VA

Di Florio A, Forty L, Gordon-Smith K et al (2013) Perinatal episodes across the mood disorder spectrum. *JAMA Psychiatry* 70:168–175
Viguera AC, Whitfield T, Baldessarini RJ et al (2007) Risk of recurrence in women with bipolar disorder during pregnancy: prospective study of mood stabilizer discontinuation. *Am J Psychiatry* 164:1817–1824; quiz 1923. doi:[10.1176/appi.ajp.2007.06101639](https://doi.org/10.1176/appi.ajp.2007.06101639)

Wesseloo R, Kamperman AM, Munk-Olsen T et al (2016) Risk of postpartum relapse in bipolar disorder and postpartum psychosis: a systematic review and meta-analysis. *Am J Psychiatr* 173:117–127. doi:[10.1176/appi.ajp.2015.15010124](https://doi.org/10.1176/appi.ajp.2015.15010124)

Wing JK, Babor T, Brugha T et al (1990) SCAN: schedules for clinical assessment in neuropsychiatry. *Arch gen Psychiatry* 47:589–593