

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/109213/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Burnley-Hall, Nicholas, Abdul, Fairuz, Androshchuk, Vitaliy, Morris, Keith, Osseigerning, Nick, Anderson, Richard, Rees, D. Aled and James, Philip 2018. Dietary nitrate supplementation reduces circulating platelet-derived extracellular vesicles in coronary artery disease patients on clopidogrel therapy: a randomised, double-blind, placebo-controlled study. *Thrombosis and Haemostasis* 118 (01) , pp. 112-122. 10.1160/TH17-06-0394

Publishers page: <http://dx.doi.org/10.1160/TH17-06-0394>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Dietary Nitrate Supplementation Reduces Circulating Platelet-Derived Extracellular Vesicles in Coronary Artery Disease Patients on Clopidogrel Therapy: A Randomised, Double-Blind, Placebo-Controlled Study

Nicholas Burnley-Hall¹ Fairoz Abdul¹ Vitaliy Androshchuk¹ Keith Morris² Nick Ossei-Gernings³ Richard Anderson³ D. Aled Rees⁴ Philip E. James²

¹ Department of Biomedical Science, School of Medicine, Cardiff University, Cardiff, United Kingdom

² Department of Biomedical Science, Cardiff School of Health Sciences, Cardiff Metropolitan University, Cardiff, United Kingdom

³ Department of Cardiology, University Hospital of Wales, Heath Park, Cardiff, United Kingdom

⁴ Neurosciences and Mental Health Research Institute, School of Medicine, Cardiff University, Cardiff, United Kingdom

Address for correspondence Professor Philip E. James, PhD,
Department of Biomedical Science, Cardiff School of Health Sciences,
Cardiff Metropolitan University, Cardiff, CF5 2SG, United Kingdom
(e-mail: pjames@cardiffmet.ac.uk).

Abstract

Extracellular vesicles (EVs) are implicated in the pathogenesis of cardiovascular disease (CVD). Specifically, platelet-derived EVs are highly pro-coagulant, promoting thrombin generation and fibrin clot formation. Nitrate supplementation exerts beneficial effects in CVD, via an increase in nitric oxide (NO) bioavailability. Clopidogrel is capable of producing NO-donating compounds, such as S-nitrosothiols (RSNO) in the presence of nitrite and low pH. The aim of this study was to assess the effect of nitrate supplementation with versus without clopidogrel therapy on circulating EVs in coronary artery disease (CAD) patients. In this randomized, double-blind, placebo-controlled study, CAD patients with (n = 10) or without (n = 10) clopidogrel therapy received a dietary nitrate supplement (SIS nitrate gel) or identical placebo. NO metabolites and platelet activation were measured using ozone-based chemiluminescence and multiple electrode aggregometry. EV concentration and origin were determined using nanoparticle tracking analysis and time-resolved fluorescence. Following nitrate supplementation, plasma RSNO was elevated (4.7 ± 0.8 vs 0.2 ± 0.5 nM) and thrombin-receptor mediated platelet aggregation was reduced (19.9 ± 6.0 vs 4.0 ± 6.4 U) only in the clopidogrel group compared with placebo. Circulating EVs were significantly reduced in this group (1.183e11 ± 3.15e10 vs 9.93e9 ± 1.84e10 EVs/mL), specifically the proportion of CD41b EVs (2,120 ± 728 vs 235 ± 436 RFU [relative fluorescence unit]) compared with placebo. In vitro experiments demonstrated clopidogrel-SNO can reduce platelet-EV directly (6.209e10 ± 4.074e9 vs 3.94e11 ± 1.91e10 EVs/mL). In conclusion, nitrate supplementation reduces platelet-derived EVs in CAD patients on clopidogrel therapy, increasing patient responsiveness to clopidogrel. Nitrate supplementation may represent a novel approach to moderating the risk of thrombus formation in CAD patients.

Introduction

Extracellular vesicles (EVs) are small, spherical structures (0.1–1-µm diameter) enclosed within a phospholipid bilayer, generated by multiple cell types including platelets, leukocytes and endothelial cells. EVs represent a novel way of communication between cells, transporting both protein and genetic material. Recently, EVs have been implicated as a potential biomarker of cardiovascular disease (CVD), and are considered to play a crucial role in the promotion of coagulation,¹ inflammation² and cell survival.³ Elevated levels of EVs have been observed in a range of CVD states,⁴ and have been shown to correlate with poor outcomes in coronary artery disease (CAD) patients.⁵ Therefore, EVs represent an attractive target for pharmacological and/or dietary intervention.

Platelet-derived EVs have received the most attention in the field, primarily due to their abundance and reactivity.⁶ The surface of platelet-derived EVs is approximately 100-fold more pro-coagulant than the surface of activated platelets themselves.⁷ Platelet-derived EVs express phosphatidylserine

that stimulates thrombin generation and stabilizes fibrin clot formation.¹ Therefore, a method of reducing platelet-EVs might be of clinical benefit in CVD patients. Thienopyridines, such as clopidogrel, act via irreversible inhibition of the adenosine diphosphate (ADP) receptor, preventing platelet activation. In addition to their antiplatelet actions, additional properties have been reported, particularly with clopidogrel, including improvement in nitric oxide (NO) bioavailability, anti-inflammatory effects and reduced endothelial injury after percutaneous coronary intervention (PCI).⁸ Previous studies have shown that clopidogrel reduces EV production in parallel with suppression of platelet activation in patients with acute coronary syndromes (ACSs).⁹ NO plays an essential role in maintaining both endothelial and platelet homeostasis; however, NO levels are compromised in CVD, resulting in the progression of atherosclerosis and vascular dysfunction.¹⁰ In platelet physiology, NO, acting via the cyclic guanosine monophosphate (cGMP) pathway, offers an additional inhibitory pathway to thienopyridine mediated inhibition. When activated together, *ex vivo* studies have shown that they produce a synergistic anti-platelet effect, culminating in a reduction in intracellular calcium.¹¹ Ingestion of dietary nitrate elevates circulating levels of nitrite via bioconversion by commensal bacteria in the enterosalivary circulation.¹² The benefits of nitrate supplementation in CVD are well documented; it has recently been shown to improve vascular function in hypercholesterolemia patients, and reduce *ex vivo* platelet aggregation in response to ADP/collagen.^{13,14} Additionally, Lee et al have recently shown that sustained-release nitrates can act synergistically with clopidogrel to enhance platelet inhibition in patients undergoing PCIs.¹⁵ S-nitrosothiols (RSNO) are a class of compound formed by S-nitrosation of reduced sulphhydryl groups, and are able to act as NO donors, with potent anti-platelet effects.¹⁶ Our group has previously demonstrated that thienopyridines readily form RSNO,¹⁷ dependent on a low pH and nitrite(NO₂⁻) availability, both of which are present in the stomach. These thienopyridine–SNO molecules are able to mimic NO with the additional benefit that the NO “moiety” is protected and preserved from further reactive metabolism. We hypothesized that inorganic nitrate supplementation with clopidogrel therapy would increase plasma RSNO levels and subsequently reduce circulating EVs in CAD patients.

Methods

Subjects and Protocol

Twenty patients with established CAD volunteered and consented to take part in this proof-of-concept, randomized, double-blind, placebo-controlled, crossover study. Patients were split into those receiving clopidogrel 75 mg daily (n = 10) and those receiving no anti-platelet therapy, referred to henceforth as “naïve” (n = 10). Patients attended the Cardiology Day Case Unit, University Hospital of Wales (UHW), as part of their normal clinical care. All patients were randomized to receive either 200 mL (67.2 mM, confirmed by ozone-based chemiluminescence) of dietary nitrate supplement (Science in Sport GoNitrate), followed by a placebo (of identical appearance), or vice versa. We have previously shown that from ingestion of this dose of inorganic nitrate, plasma nitrate and nitrite values peak at 2 hours and have returned to baseline within 24 hours.¹⁸ We adopted a washout period of 7 days, in agreement with several other groups.^{19,20} ► Fig. 1 summarizes the study design. Blood samples were taken from the antecubital vein through an 18-g Venflon intravenous (IV) cannula into EDTA (ethylenediaminetetraacetic acid), citrate and hirudin vacutainers. Samples were taken both before ingestion of the supplement and 2 hours post-supplementation. Patients took all prescribed medication at least 1 hour prior to the study. Patients were included if they were male, over 18 years of age with stable CAD and had been fasted for >6 hours. Patients in the clopidogrel group must have been receiving clopidogrel for >1 month prior to commencement of the study. Patients were excluded if they had a clopidogrel intolerance or contraindication, were on other long-term oral anti-coagulant drugs or receiving IV or subcutaneous anti-thrombin therapy. Patients were also excluded if they had any ischemic event (ACS, stroke or transient ischemic attack) or revascularization procedure (PCI or CABG) within the preceding 3 months, chronic renal or liver disease, or an inability to give informed consent. This study conformed to the ethical principles contained in the Declaration of Helsinki. Ethical approval was provided by the South East Wales Research Ethics Committee (IRAS Project ID 102427).

Biochemical Measurements

A full blood count was measured on an ABX Pentra X120

A full blood count was measured on an ABX Pentra X120 haematology blood analyser (Horiba, Northampton, UK). Serum cholesterol, high-density lipoprotein cholesterol and triglycerides were assessed using an Aeroset automated analyser (Abbott Diagnostics, Berkshire, UK). Low-density

lipoprotein cholesterol was calculated using Friedewald's formula. C-reactive protein was assayed by nephelometry (BN-II system, Milton Keynes, UK). All biochemical measurements were performed by the Department of Medical Biochemistry, UHW.




Fig. 1 Study design. Patients were split into those receiving clopidogrel 75 mg daily ($n = 10$) and those receiving no anti-platelet therapy (naïve, $n = 10$). Patients were randomly allocated to either nitrate supplementation (Science in Sport Go⁺ Nitrates gel), followed by a placebo of identical appearance, or vice versa. A washout interval of a minimum of 7 days separated the two treatment periods. Blood samples were taken immediately before ingestion of the supplement and 2 hours post-supplementation.

Measurement of Plasma NO Metabolites

Blood samples were collected into EDTA vacutainers and promptly centrifuged (2,500g, 15 minutes, 4°C) to yield plasma which was subsequently snap frozen in liquid nitrogen and stored at -80°C until analysis. NO metabolites were assessed using ozone-based chemiluminescence. Briefly, for plasma nitrite and RSNO analysis, 5-mL tri-iodide reagent (I₃⁻) was placed in a custom-built glass purge vessel and heated at 55°C via a water bath thermostatically controlled by a hotplate. The inert carrier gas (O₂-free N₂) purging I₃⁻ was linked to a sodium hydroxide trap (1 M), connected to an NO analyser (Sievers NOA 280i; Analytix, Boldon, UK). Plasma samples (200 µL nitrite, 400 µL RSNO) were directly injected in to the purge vessel through a rubber septum injection inlet. To differentiate between nitrite and RSNO, samples were run before and after pre-treatment with 5% acidified sulphanilamide (290 mM), which removes NO₂⁻, thus allowing the selective measurement of the residual plasma RSNO. Increased accuracy of RSNO measurement was obtained by a 50-point adjacent signal averaging algorithm, improving the signal-to-noise ratio, using Origin 7.0 (OriginLab, Massachusetts). For nitrate analysis, plasma samples (20 µL) were injected into vanadium chloride heated at 90°C before detection via the NO analyser. Results were compared with a sodium nitrite/nitrate standard curve performed daily to account for day-to-day temperature fluctuations. Room temperature was 19 ± 2°C. In our hands, the limit of sensitivity for these plasma RSNO, nitrite and nitrate assays are >1, >10 and >500 nmol/L, respectively.

Platelet Aggregation

Whole blood aggregation was assessed via multiple electrode aggregometry, using the Multiplate analyser (Roche Diagnostics, Forrenstrasse, Switzerland). The adhesion and aggregation of platelets on the sensor surface enhances the electrical resistance between the two sensor electrodes within a test cell. Whole blood (300 µL) collected in a hirudin vacutainer was diluted 1:1 with saline solution and incubated at 37°C for 3 minutes and measured within 30 minutes of blood drawing. Following this, platelet activation was initiated by the addition of either ADP (6.5 µM) or thrombin receptor-activating peptide (TRAP; 32 µM). The subsequent increase in impedance due to attachment of platelets to the

electrodes is transformed to arbitrary aggregation units. The aggregation measured is quantified as the area under the curve (AUC) and expressed in units (U).

Extracellular Vesicle Isolation

Blood was collected into sodium citrate vacutainers, as recommended by the Scientific Standardisation Committee of the International Society on Thrombosis and Haemostasis.^{21,22} Blood was immediately centrifuged (2,500g, 15 minutes, 4°C) to isolate plasma, which was subsequently spun (2,500g, 15 minutes, 4°C) to obtain platelet-poor plasma (PPP). PPP was ultracentrifuged (100,000g, 60 minutes, 4°C) and the resulting EV pellet was resuspended in phosphate-buffered saline (PBS) and slow-frozen overnight at -80°C. All EV analysis was completed within 7 days of isolation. This is an established protocol that we have utilized in several clinical studies to date.^{23–25} EV Size and Concentration nEV concentration and size distribution were determined using nanoparticle tracking analysis (NTA; NanoSight LM10 system, UK) as described previously. NTA is a laser-illuminated microscopic technique equipped with a 642-nm laser and a high-sensitivity digital camera system (Orca-Flash2.8, Hamamatsu, NanoSight Ltd) that determines the brownian motion of nanoparticles in real time to assess size and concentration. Sixty-second videos were recorded and particle movement was analysed using NTA software (version 2.3). Camera sensitivity and detection threshold were 11 to 14 and 4 to 6, respectively. EV samples were diluted in EV-free sterile PBS. Samples were run in quintuplicate, from which EV distribution, size and average concentration were calculated. EV concentrations were expressed as EVs/mL plasma.

EV Cellular Origin

Time-resolved fluorescence was used to assess the origin of isolated EVs, as described previously.²³ EVs (5e10/well) were loaded into a high protein binding 96-well plate (Greiner Bio-One, Germany) overnight at 4°C, before non-specific sites were blocked with 1% bovine serum albumin for 2 hours. EVs were incubated overnight at 4°C with mouse anti-human antibodies against CD9, CD41, CD11b, CD235a and CD144 as markers of exosomes, platelets, leukocytes, erythrocytes and endothelial cells, respectively (all antibodies from Abcam, Cambridge, UK). Markers were detected using a biotinylated anti-mouse IgG secondary antibody and a streptavidin:europium conjugate (PerkinElmer, Buckinghamshire, UK).

In Vitro Platelet EV Production

The in vitro model was designed to confirm the capability of clopidogrel-SNO to inhibit platelet EV release in response to a pharmacological stimulus (ADP). Platelet-rich plasma (PRP) was isolated from the blood of healthy volunteers via sodium citrate vacutainers (200g, 20 minutes). PPP was also isolated (200g, 20 minutes, followed by 2 × 2,500g, 15 minutes) for use as a control. Platelets were stimulated with ADP (6.5 μM) and incubated for 5 minutes before NaNO₂, clopidogrel, S-nitrosoglutathione (GSNO) or clopidogrel-SNO (all 10 μM) were added and incubated for 1 hour at 37°C. This concentration was chosen to compare directly with in vitro aggregation studies in healthy blood showing the IC₅₀ for GSNO/ clopidogrel-SNO as approximately 7.5 μM. Clopidogrel-SNO was produced as described in detail previously by our group.¹⁷

Statistics

A power calculation based on prior results from EV samples from CAD patients showed that nine subjects would provide 90% power for detecting a 20% difference in circulating EVs between placebo and nitrate supplementation, assuming 10% variation, with $\alpha = 0.05$. Data were analysed using GraphPad Prism version 5.0. Data are expressed as mean ± standard error (SE). All data were assessed for both a period effect and treatment-period interaction and checked for normality using the Kolmogorov-Smirnov test. The change in measurement before and after nitrate supplementation or placebo was calculated and compared directly using a paired t-test or a Wilcoxon matched pairs test, for normally or non-normally distributed data, respectively.

Results

Subject Characteristics

Of the 20 males that participated in the study, 10 were taking clopidogrel (>1 month) and 10 were not receiving antiplatelet therapy (naïve group). The mean ages of the groups were 63.2 ± 3.6 and 62.7 ± 3.2 years, respectively. Biochemical measurements are summarized in ► Table 1. Importantly, no differences were observed in platelet count at baseline or at the follow-up visit. No differences were

seen in age, body mass index, biochemical or haematological parameters between groups. No significant period effects and treatment–period interaction were observed for any of the parameters determined in this study.

Plasma NO Metabolites

Baseline plasma nitrate, nitrite and RSNO measurements were not significantly different between naïve and clopidogrel groups (nitrate [NO₃⁻]: 31.14 ± 2.5 vs 31.34 ± 5.1 μM; NO₂⁻: 92.3 ± 9.0 vs 118.9 ± 15.7 nM; RSNO: 6.4 ± 0.5 nM vs 11.9 ± 3.4 μM). Nitrate supplementation significantly elevated plasma NO₃⁻ and NO₂⁻ compared with placebo in both naïve (ΔNO₃⁻: 252.1 ± 17.4 μM vs 3.7 ± 1.9 μM, p < 0.001; ΔNO₂⁻: 167.8 ± 40.1 vs 18.5 ± 22.6 nM, p < 0.05) (► Fig. 2a,c) and clopidogrel (ΔNO₃⁻: 256.6 ± 20.5 vs 2.0 ± 3.4 μM, p < 0.001; ΔNO₂⁻: 164.8 ± 68.5 vs 5.2 ± 12.2 nM, p < 0.05) (► Fig. 2b, d) groups. Increases in plasma nitrate and nitrite were similar between naïve and clopidogrel groups (ΔNO₃⁻ naïve: 252.1 ± 17.4 μM vs clopidogrel: 256.6 ± 20.5 μM, p > 0.05; ΔNO₂⁻ naïve: 167.8 ± 40.1 nM vs clopidogrel: 164.8 ± 68.5 nM, p > 0.05). Plasma RSNO did not change following nitrate supplementation compared with placebo in the naïve group (ΔRSNO: 0.9 ± 1.3 vs 1.3 ± 0.6 nM) (► Fig. 2e). However, plasma RSNO was significantly elevated after receiving nitrate supplementation when compared with placebo in the clopidogrel group (ΔRSNO: 4.7 ± 0.8 vs 0.2 ± 0.5 nM, p < 0.001) (► Fig. 2f). Additionally, RSNO in the clopidogrel group was significantly elevated in comparison to the naïve group (4.7 ± 0.8 vs 0.9 ± 1.3 nM, p < 0.05), respectively.

Platelet Aggregation

TRAP-mediated platelet aggregation was not significantly different between clopidogrel and naïve groups (94.8 ± 7.0 vs 99.0 ± 6.3 U, respectively). Patients receiving clopidogrel had markedly lower ADP-mediated platelet aggregation compared with the naïve group at baseline (37.9 ± 3.80 vs 60.0 ± 4.4 U, p < 0.001). Nitrate supplementation failed to significantly reduce both ADP- and TRAP-mediated platelet aggregation in the naïve group when compared with placebo (ΔADP: 7.8 ± 5.8 vs 2.1 ± 4.9 U, ΔTRAP: 21.5 ± 8.4 vs 3.8 ± 9.2 U) (► Fig. 3a). Nitrate supplementation had no effect on ADP-mediated platelet aggregation compared with placebo in patients receiving clopidogrel (ΔADP: 2.2 ± 2.2 vs 0.1 ± 1.8 U) (► Fig. 3b). However, nitrate supplementation did significantly reduce TRAP-mediated platelet aggregation in the clopidogrel group compared with placebo (ΔTRAP: 19.9 ± 6.0 vs 4.0 ± 6.4 U, p < 0.05) (► Fig. 3b). The reduction in ADP- and TRAP-mediated aggregation was similar within both the naïve and clopidogrel groups (ΔADP: naïve: 7.8 ± 5.8 U vs clopidogrel: 2.2 ± 2.2 U, p > 0.05. ΔTRAP: naïve: 21.5 ± 8.4 U vs clopidogrel: 19.9 ± 6.0 U, p > 0.05).

Table 1 Participant characteristics

Abbreviations: ACEi/ARB, angiotensin converting enzyme inhibitors/angiotensin II receptor blocker; BMI, body mass index; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; CRP, C-reactive protein; GTN, glyceryl trinitrate; HDL, high-density lipoprotein; LDL, low-density lipoprotein; MI, myocardial infarction; NSAIDs, non-steroidal anti-inflammatory drugs; TIA, transient ischemic attack.

Note: Summary of patient characteristics including age, BMI, SHIM score, cardiovascular risk factors, medications, biochemical and haematological measurements. Haematological measures were taken at the beginning of both patient visits, prior to any treatment. No statistically significant differences between the two groups were observed (p > 0.05).

	Naïve group (n = 10)		Clopidogrel group (n = 10)	
Age	62.7 ± 3.19		63.2 ± 3.66	
BMI (kg/m ²)	29.9 ± 1.49		28.2 ± 1.18	
Cardiovascular risk factors				
Diabetes mellitus	1 (10%)		2 (20%)	
Past/current smoking	3 (30%)		6 (60%)	
Hypertension	7 (70%)		8 (80%)	
Dyslipidaemia	10 (100%)		8 (80%)	
Family history of premature				
Heart disease (<65 y)	4 (40%)		2 (20%)	
Stroke/TIA	1 (10%)		2 (20%)	
Peripheral vascular disease	1 (10%)		1 (10%)	
History of MI	4 (40%)		7 (70%)	
Previous revascularization				
PCI	4 (40%)		7 (70%)	
CABG	2 (20%)		4 (40%)	
Respiratory disease (asthma/COPD)	0 (0%)		3 (30%)	
Medications				
Aspirin	7 (70%)		4 (40%)	
Clopidogrel	0 (0%)		10 (100%)	
Proton-pump inhibitor	5 (50%)		7 (70%)	
Beta blockers	4 (40%)		6 (60%)	
ACEI/ARB	6 (60%)		8 (80%)	
Statins	10 (100%)		8 (80%)	
GTN	1 (10%)		1 (10%)	
Thyroxin	1 (10%)		0 (0%)	
NSAIDs	0 (0%)		2 (20%)	
Oral anti-coagulants	1 (10%)		0 (0%)	
Calcium channel blockers	2 (20%)		1 (10%)	
Diuretics	2 (20%)		2 (20%)	
Biochemical measures				
CRP (mg/L)	5.3 ± 4.27		4.90 ± 4.63	
Total cholesterol (mmol/L)	4.58 ± 0.77		4.30 ± 0.82	
Triglycerides (mmol/L)	2.1 ± 1.80		1.47 ± 0.90	
HDL (mmol/L)	1.11 ± 0.26		1.07 ± 0.13	
LDL (mmol/L)	2.61 ± 0.54		2.57 ± 0.68	
Total cholesterol:HDL ratio	4.13 ± 1.69		4.02 ± 0.57	
Haematological measures				
	Visit 1	Visit 2	Visit 1	Visit 2
White cell count (× 10 ⁹ /L)	6.0 ± 0.50	6.3 ± 0.6	6.7 ± 0.6	6.4 ± 0.3
Haemoglobin (g/L)	149.1 ± 3.1	151.5 ± 3.6	147.5 ± 3.4	144.9 ± 3.4
Platelet count (× 10 ⁹ /L)	232.8 ± 25.0	219.4 ± 16.1	234.7 ± 13.4	237.8 ± 18.2

Fig. 2 NO metabolite measurements. The change in plasma NO metabolites 2 hours post-nitrate supplementation or placebo. (a) Naïve group plasma NO₃⁻, μM. (b) Clopidogrel group plasma NO₃⁻, μM. (c) Naïve group NO₂⁻, nM. (d) Clopidogrel group plasma NO₂⁻, nM. (e) Naïve group plasma RSNO, nM. (f) Clopidogrel group plasma RSNO, nM. Data are expressed as mean ± SEM (n = 10). p < 0.05 and □□□ p < 0.001





Fig. 3 Platelet aggregation. ADP- and TRAP-induced platelet aggregation as measured by MEA (multiple electrode aggregometry). (a) Agonist response in the naïve group. (b) Agonist response in the clopidogrel group. Data are expressed as mean \pm SEM (n = 10). $p < 0.05$.



EV Size and Concentration

Nitrate supplementation had no effect on the mode size of EVs compared with placebo in both naïve (123 ± 9.2 vs 111 ± 7.3 nm) and clopidogrel (129 ± 11.4 vs 122 ± 16.2 nm) groups, respectively. EV concentration at baseline did not differ between naïve and clopidogrel groups, respectively ($5.0 \times 10^{11} \pm 4.2 \times 10^{10}$ EVs/mL vs $4.7 \times 10^{11} \pm 3.7 \times 10^{10}$ EVs/mL). ► Fig. 4a shows the size distribution profile of EVs in both the naïve and clopidogrel group at baseline, split into 50-nm bin sizes. Nitrate supplementation did not reduce the circulating EV concentration compared with placebo in the naïve group (Δ EVs: $2.78 \times 10^{10} \pm 4.22 \times 10^{10}$ vs $1.76 \times 10^{10} \pm 1.23 \times 10^{10}$ EVs/mL). However, nitrate supplementation did significantly reduce EV concentration in the clopidogrel group compared with placebo (Δ EVs: $1.183 \times 10^{11} \pm 3.15 \times 10^{10}$ vs $9.93 \times 10^9 \pm 1.84 \times 10^{10}$ EVs/mL, $p < 0.01$) (► Fig. 4b). The reduction in circulating EV concentration was similar between naïve and clopidogrel groups ($2.78 \times 10^{10} \pm 4.22 \times 10^{10}$ vs $1.183 \times 10^{11} \pm 3.15 \times 10^{10}$ EVs/mL, $p > 0.05$), respectively.

EV Cellular Origin

Comparison between the naïve and clopidogrel groups revealed that the clopidogrel group had significantly higher baseline levels of exosomal (CD9), platelet (CD41) and leukocyte (CD11b) markers compared with the naïve group




Fig. 4 Circulating EV concentration. (a) Size distribution profile for EVs in both the naïve and clopidogrel group, split into 50-nm bin sizes. (b) Change in circulating plasma EV concentration 2 hours post-nitrate supplementation/placebo in naïve and clopidogrel groups determined by NTA (nanoparticle tracking analysis). Data are expressed as mean \pm SEM ($n = 10$). $p < 0.05$.

(CD9: $32,899 \pm 1,303$ vs $23,812 \pm 1,891$ RFU [relative fluorescence unit], $p < 0.001$; CD41: $15,753 \pm 1,372$ vs $10,064 \pm 705$ RFU, $p < 0.01$; CD11b: $14,245 \pm 1,512$ vs $9,578 \pm 1,250$ RFU, $p < 0.05$) (► Fig. 5a). No significant differences were seen in surface protein markers for exosomes, platelets, leukocytes, erythrocytes (CD235a) and endothelial cells (CD144) of EVs in the naïve group following

nitrate supplementation compared with placebo (► Fig. 5b). CD41 was significantly reduced following nitrate supplementation compared with placebo in patients receiving clopidogrel (Δ CD41: $_2,120 _728$ vs $_235 _436$ RFU, $p < 0.05$) (► Fig. 5c). Changes in surface marker levels were similar




Fig. 5 Effect of nitrate supplementation on EV surface protein. The change in content of exosomes (CD9), platelet (CD41), leukocyte (CD11b), erythrocyte (CD235a) and endothelial (CD144) in plasma EVs after nitrate supplementation versus placebo. (a) Difference in baseline levels of protein expression between naïve and clopidogrel groups ($n = 10$). (b) Change in protein expression after nitrate supplementation/placebo in the naïve group ($n = 10$). (c) Change in protein content after nitrate supplementation/placebo in the clopidogrel group ($n = 10$). Proteins were detected using a streptavidin–europium conjugate and measured using time-resolved fluorescence (relative fluorescence units [RFU]). Data are expressed as mean \pm SEM. $***p < 0.001$, $**p < 0.01$ and $*p < 0.05$. between naïve and clopidogrel groups (CD9: $_471 _625$ vs $_1,079 _744$, CD41: $_498 _423$ vs $_2,120 _728$, CD11b: $_1,053 _934$ vs $_877 _441$, CD235a: $_259 _230$ vs $_678 _354$, CD144: $_364 _354$ vs $_106 _336$, $p > 0.05$ for all comparisons).

In Vitro Experiments

Laboratory experiments were undertaken to investigate the direct effect of nitrite, clopidogrel and clopidogrel–SNO separately on platelet activation and EV generation. EV generation was increased 200% in response to addition of ADP. Following a dose–response, 10 μ M of inhibitors (sodium nitrite, clopidogrel–SNO, GSNO) were added, which showed this concentration was effective at inhibiting platelet aggregation and thus platelet-derived EV release. Indeed, aggregation measured in response to ADP was significantly inhibited by both clopidogrel–SNO and GSNO, with a similar IC₅₀ (6.56 \pm 1.08 vs 8.42 \pm 1.35 μ M). Nitrite alone was ineffective at inhibiting platelet aggregation in response to ADP unless concentrations greater than 10 mM were applied (► Fig. 6a). The addition of sodium nitrite (NaNO₂) or clopidogrel to stimulated PRP had no effect on EV production (NaNO₂: 3.83e11 \pm 1.707e10 EVs/mL, clopidogrel: 3.91e11 \pm 1.805e10 EVs/mL, vs ADP alone as control 3.94e11 \pm 1.91e10 EVs/mL). Similarly, no change in CD41 was observed (NaNO₂: 20,093 \pm 1,244 RFU, clopidogrel: 18,238 \pm 2,824 RFU, vs ADP control: 19,703 \pm 1,375 RFU, $p > 0.05$). Following the addition of clopidogrel–SNO, EV production significantly reduced compared with the ADP control (6.209e10 \pm 4.074e9 vs 3.94e11 \pm 1.91e10 EVs/mL, $p < 0.001$). GSNO also significantly reduced EV production compared with the ADP control (8.67e10 \pm 8.63e9 EVs/mL, $p < 0.001$) (► Fig. 6b). Reductions were also seen in CD41 expression following addition of both SNO molecules compared with control (GSNO: 8,946 \pm 1,460 RFU, clopidogrel–SNO: 9,713 \pm 1,214 RFU, vs ADP control: 19,703 \pm 1,375 RFU, $p < 0.01$) (► Fig. 6c).

Discussion

This study shows that acute dietary NO₃⁻ supplementation significantly reduces circulating EVs in patients with CAD already established on long-term clopidogrel therapy. Furthermore, it significantly reduced platelet activation via the thrombin receptor, accompanied by a reduction in the proportion of platelet-derived EVs. This reduction was associated with a rise in circulating RSNO only in the clopidogrel group. Previously thought of as a physiologically inert oxidative end product of NO metabolism, it is now widely accepted that NO₃⁻ and NO₂⁻ metabolism occurs in both blood and tissues to form NO and other bioactive NO metabolites, representing an alternative to the “classical” L-arginine pathway. 12 NO₃⁻ supplementation has been shown to have a variety of benefits in CVD including improving vascular function,¹³ decreasing blood pressure²⁶ and attenuating oxidative stress.²⁷

Baseline measurements of plasma NO₂⁻ prior to NO₃⁻ supplementation or placebo were noticeably lower in both the naïve (92.3 \pm 9.0 nM) and clopidogrel (118.9 \pm 15.7 nM) groups compared with the values typically observed in healthy individuals (200–300 nM) we have reported previously.²⁸ These low NO₂⁻ levels reflect the reduced NO bioavailability and degree of endothelial dysfunction seen in these patient cohorts.

Significant elevations in plasma RSNO levels following NO₃⁻ supplementation were seen only in patients taking clopidogrel. We have previously demonstrated in vitro that the low pH environment of the stomach can modify thienopyridines, to form thienopyridine–SNO molecules²⁹ that exhibit NO-like biological effects, including platelet inhibition and vasodilation.^{30,31} The low pH exposes the free thiol group present in these drugs, before biotransformation in the presence of NO₂⁻ to form RSNO compounds. These molecules provide an addition to the nitrate–nitrite–NO pathway and the effects seen upon nitrate/nitrite supplementation.³²

Patients receiving proton-pump inhibitors (PPIs; and thus an increased pH in the stomach) exhibit reduced platelet inhibition in response to clopidogrel, although the mechanism of this is felt to be multifactorial and incompletely defined.³³ Recently, Pinheiro et al have shown that oral nitrite administration in rats was associated with an increase in RSNO and a decrease in blood pressure.³⁴ Furthermore, treatment with both PPIs and the thiol-depleting agent buthionine sulfoximine attenuated the increase in plasma RSNO and blunted the anti-hypertensive effects of nitrite.³⁴




Fig. 6 In vitro platelet EV production. (a) Dose–response curve of GSNO, clopidogrel–SNO and nitrite on platelet inhibition. The IC₅₀ of GSNO and clopidogrel–SNO were similar (6.56 ± 1.08 vs $8.42 \pm 1.35 \mu\text{M}$). (b) The effect of various agents on EV production. Clopidogrel–SNO severely attenuated EV production in platelet-rich plasma. An alternative nitrosothiol, GSNO, also inhibited EV production. (c) The reduction in EV concentration was mirrored by a reduction in CD41 expression, measured by TRF. $n = 4$, $p < 0.001$ and $p < 0.01$.

However, the influence of stomach pH on RSNO formation in patients is yet to be established.

To our knowledge, this is the first study to demonstrate that dietary nitrate supplementation can reduce circulating EV concentration when administered with clopidogrel. The lack of RSNO produced in the naïve group and absence of a significant reduction in EV concentration suggest these effects are due to formation of clopidogrel–SNO molecules in the stomach. Conversely, in patients receiving clopidogrel and NO₃⁻, clopidogrel–SNO is formed which can subsequently donate NO in the plasma as previously demonstrated.³⁵ Our group have previously shown that nitrite-derived NO can reduce EV production in endothelial cells.³⁶ This is consistent with previous reports that showed impairment of NO production increased EV production in endothelial cells.³⁷ It is well established that NO elicits many of its effects, such as platelet activation, via a reduction in intracellular calcium. Conversely, many of the processes required for EV formation, such as lipid membrane remodelling and cytoskeleton disruption require increases in intracellular calcium.³⁸ Our in vitro results confirm clopidogrel–SNO, derived from NO₂⁻, inhibits platelet EV generation directly, where NO₂⁻ and clopidogrel alone had no effect. This is consistent with inhibition of platelet aggregation we have observed at these concentrations and with previously established effects of RSNO on platelets.³⁹

Interestingly, a significant reduction in EV concentration was only seen in the clopidogrel group. Whether this decrease can be attributed to the increased RSNO levels seen compared with the naïve group, or as a result of the combination of two independent effects of NO and clopidogrel, remains unclear. Clopidogrel acts via inhibition of the ADP/ P2Y₁₂ receptor, preventing the inhibition of adenylate cyclase, consequently reducing platelet aggregation. The ADP receptor has previously been shown to have a potentiating role in dense granule secretion.⁴⁰ Granule and vesicle release from platelets is mediated by common cellular machinery such as SNARE proteins, intracellular calcium levels and cytoskeletal organization.⁴¹ Thus, it is possible that the reduction in plasma EV seen in our patient cohort could be due to the combined, independent effects of NO and clopidogrel. However, given our data showing a significant reduction of platelet EV seen when clopidogrel–SNO is administered in vitro, it seems likely that the increased formation of RSNO in these patients causes the reduction in EV seen in this study.

Consistent with previous ex vivo studies, we provide evidence that nitrate supplementation can act synergistically with the P2Y₁₂ inhibitor clopidogrel resulting in augmented platelet inhibition than clopidogrel alone.¹² The significant reduction in TRAP-induced platelet activation suggests RSNO may act via elevation of cGMP, as shown previously.⁴²

Measurement of surface protein content revealed that the proportion of platelet-derived EVs decreased in the clopidogrel group. This specific reduction in platelet-derived EVs is in keeping with the hypothesis that the combined effects of RSNO and clopidogrel are responsible for the reduction in EV concentration seen in this study. Both cGMP and cyclic adenosine monophosphate are established synergistic mediators of platelet inhibition,⁴³ and it has previously been established that combined NO and clopidogrel have a synergistic effect on platelet activation.¹¹

Platelet aggregation plays a key role in the development of atherosclerosis, and anti-platelet therapy is well established in the treatment of CVD. Under physiological conditions, adhesion of platelets to the endothelium is inhibited by endogenous production of NO, highlighted by the reduced NO bioavailability observed in CVD states. Previous studies have also shown a reduction in platelet activation ex vivo following nitrate supplementation.²⁶ These authors also showed that prevention of the enterosalivary bioconversion of NO₃⁻ to NO₂⁻ prevented the expected rise in plasma NO₂⁻, the decrease in blood pressure and the inhibition of platelet aggregation.²⁶

This study highlights the ability of a simple dietary intervention, in combination with clopidogrel, to reduce platelet activation, as well as circulating, pro-coagulant EVs in a CVD cohort. Thus, inorganic nitrates are capable of increasing patient responsiveness to clopidogrel. High on-treatment platelet reactivity is common in approximately 15 to 40% of patients prescribed clopidogrel.⁴⁴ As formation of clopidogrel–SNO does not require activation of the pro-drug, all patients on clopidogrel would benefit from increases in NO bioavailability, regardless of their clopidogrel metabolism. This may explain the various pleiotropic effects of clopidogrel that have been observed in recent studies.⁸ Ultimately, reducing the circulating, pro-coagulant EVs in CAD patients offers a new approach to moderating of the risk of thrombus formation and thus a myocardial infarction. This may also be of therapeutic

interest in other CVD cohorts, as well as cancer patients, where cancer-associated thrombosis is the second leading cause of death.⁴⁵

This study has several limitations. First, despite the robust design of a placebo-controlled, double-blind, crossover study, the small sample size of groups ($n = 10$), powered to enable the study to detect significant changes in EV concentration, limits the overall power of the study in terms of association between factors. Thus, there are several nonsignificant trends which may have been significant in a larger sample population. Although not statistically significant, a trend to decreased platelet aggregation was observed in patients not receiving clopidogrel. A large-scale study will be required to fully investigate the effect of NO₃⁻ alone on platelets; however, this observation was not accompanied by a decrease in EV or increase in RSNO. Second, the concentration of circulating RSNO attained in patients receiving clopidogrel is considerably lower than those used in *in vitro* models. The intentions of the *in vitro* studies were not to mimic the *in vivo* concentrations of these molecules, but rather a proof-of-principle to demonstrate the capability of clopidogrel–SNO to inhibit platelet EV release in response to addition of a pharmacological agent. The authors also recognize small differences in both cardiovascular risk factors and medications between the naïve and clopidogrel groups, rendering it difficult to conclude that differences in outcomes seen in this study are due to clopidogrel treatment or other potential confounding factors. Patients in both groups had been prescribed PPIs, which may have affected increased pH in the stomach toward neutrality and thus may have interfered with clopidogrel–SNO formation. However, it is important to note that PPI raises stomach pH from approximately 2 to 3 to about 4 to 6.^{46–48} At this pH, RSNO formation is only marginally reduced, as shown previously.¹⁷ Furthermore, nitrate supplementation was given about 1 hour after patients took their clopidogrel. This may not be optimal timing to facilitate the formation of clopidogrel–SNO. However, studies have highlighted the delayed transit of clopidogrel in CVD patients, visualizing intact tablets within the stomach 1.5 to 2 hours following ingestion.⁴⁹ Thus, significant levels of clopidogrel would remain in the stomach while simultaneous increases in nitrite within the stomach also occur, facilitating clopidogrel–SNO formation. Comparisons between naïve and clopidogrel groups showed a similar reduction in platelet aggregation, EV concentration and EV surface markers following nitrate supplementation. Thus, due to the lack of RSNO produced by the naïve group, it is unlikely that the formation of RSNO is the sole mechanism responsible for the beneficial effects of nitrite. Finally, our study investigated the acute effects of a single nitrate supplement. Future studies should address the effect of chronic nitrate supplementation on EV populations in relation to RSNO and platelet activity in patients.

In summary, this study shows that acute nitrate supplementation can reduce platelet-derived EVs and residual platelet activity only in patient's receiving clopidogrel and nitrate concomitantly. Increases in plasma RSNO are observed only in these patients. These results suggest that dietary NO₃⁻ supplementation could provide an additional adjunct to platelet inhibition with clopidogrel in patients at risk from CVD. Medical Biochemistry, UHW, for performing the biochemical measurements on the patient samples. Finally, the authors would like to thank the nursing team for overseeing the care of the patients at both the cardiology outpatients' clinic and the cardiac day care unit at the University Hospital of Wales.

Conflicts of Interests

None declared.

Funding

This work was funded by a Health and Care Research Wales Scholarship awarded to NBH and a clinical fellowship funded by the Nott Legacy for Thrombosis Research awarded to F. A.

References

- 1 Zubairova LD, Nabiullina RM, Nagaswami C, et al. Circulating microparticles alter formation, structure, and properties of fibrin clots. *Sci Rep* 2015;5:17611
- 2 Puddu P, Puddu GM, Cravero E, Muscari S, Muscari A. The involvement of circulating microparticles in inflammation, coagulation and cardiovascular diseases. *Can J Cardiol* 2010;26(04): 140–145
- 3 Kurachi M, Mikuni M, Ishizaki Y. Extracellular vesicles from vascular endothelial cells promote survival, proliferation and motility of oligodendrocyte precursor cells. *PLoS One* 2016;11(07):e0159158

- 4 Jansen F, Nickenig G, Werner N. Extracellular vesicles in cardiovascular disease: potential applications in diagnosis, prognosis, and epidemiology. *Circ Res* 2017;120(10):1649–1657
- 5 Sinning J-M, Losch J, Walenta K, Böhm M, Nickenig G, Werner N. Circulating CD31p/Annexin Vp microparticles correlate with cardiovascular outcomes. *Eur Heart J* 2011;32(16):2034–2041
- 6 Aatonen MT, Ohman T, Nyman TA, Laitinen S, Grönholm M, Siljander PR. Isolation and characterization of platelet-derived extracellular vesicles. *J Extracell Vesicles* 2014;3:24692
- 7 Sinauridze EI, Kireev DA, Popenko NY, et al. Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thromb Haemost* 2007;97(03):425–434
- 8 Adamski P, Koziński M, Ostrowska M, et al. Overview of pleiotropic effects of platelet P2Y12 receptor inhibitors. *Thromb Haemost* 2014;112(02):224–242
- 9 Behan MWH, Fox SC, Heptinstall S, Storey RF. Inhibitory effects of P2Y12 receptor antagonists on TRAP-induced platelet aggregation, procoagulant activity, microparticle formation and intracellular calcium responses in patients with acute coronary syndromes. *Platelets* 2005;16(02):73–80
- 10 Anderson TJ. Nitric oxide, atherosclerosis and the clinical relevance of endothelial dysfunction. *Heart Fail Rev* 2003;8(01):71–86
- 11 Kirkby NS, Lundberg MH, Chan MV, et al. Blockade of the purinergic P2Y12 receptor greatly increases the platelet inhibitory actions of nitric oxide. *Proc Natl Acad Sci U S A* 2013;110(39): 15782–15787
- 12 Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 2008;7(02):156–167
- 13 Velmurugan S, Gan JM, Rathod KS, et al. Dietary nitrate improves vascular function in patients with hypercholesterolemia: a randomized, double-blind, placebo-controlled study. *Am J Clin Nutr* 2016;103(01):25–38
- 14 Velmurugan S, Kapil V, Ghosh SM, et al. Antiplatelet effects of dietary nitrate in healthy volunteers: involvement of cGMP and influence of sex. *Free Radic Biol Med* 2013;65:1521–1532
- 15 Lee DH, Kim MH, Guo LZ, et al. Concomitant nitrates enhance clopidogrel response during dual anti-platelet therapy. *Int J Cardiol* 2016;203:877–881
- 16 Megson IL, Sogo N, Mazzei FA, Butler AR, Walton JC, Webb DJ. Inhibition of human platelet aggregation by a novel S-nitrosothiol is abolished by haemoglobin and red blood cells in vitro: implications for anti-thrombotic therapy. *Br J Pharmacol* 2000;131 (07):1391–1398
- 17 Bundhoo SS, Anderson RA, Sagan E, et al. Direct formation of thienopyridine-derived nitrosothiols—just add nitrite!. *Eur J Pharmacol* 2011;670(2–3):534–540
- 18 James PE, Willis GR, Allen JD, Winyard PG, Jones AM. Nitrate pharmacokinetics: Taking note of the difference. *Nitric Oxide* 2015;48:44–50
- 19 McQuillan JA, Casadio JR, Dulson DK, Laursen PB, Kilding AE. The effect of nitrate supplementation on cycling performance in the heat in well-trained cyclists. *Int J Sports Physiol Perform* 2017;19:1–22
- 20 Kerley CP, Cahill K, Bolger K, et al. Dietary nitrate supplementation in COPD: an acute, double-blind, randomized, placebo-controlled, crossover trial. *Nitric Oxide* 2015;44:105–111
- 21 Witwer KW, Buzás EI, Bemis LT, et al. Standardization of sample collection, isolation and analysis methods in extracellular vesicle research. *J Extracell Vesicles* 2013;2:20360
- 22 Lacroix R, Judicone C, Mooberry M, Boucekine M, Key NS, Dignat-George F; The ISTH SSC Workshop. Standardization of pre-analytical variables in plasma microparticle determination: results of the International Society on Thrombosis and Haemostasis SSC Collaborative workshop. *J Thromb Haemost* 2013;2:12207
- 23 Connolly KD, Guschina IA, Yeung V, et al. Characterisation of adipocyte-derived extracellular vesicles released pre- and postadipogenesis. *J Extracell Vesicles* 2015;4:29159
- 24 Willis GR, Connolly K, Ladell K, et al. Young women with polycystic ovary syndrome have raised levels of circulating annexin V-positive platelet microparticles. *Hum Reprod* 2014;29(12): 2756–2763
- 25 Connolly KD, Willis GR, Datta DBN, et al. Lipoprotein-apheresis reduces circulating microparticles in individuals with familial hypercholesterolemia. *J Lipid Res* 2014;55(10):2064–2072
- 26 Webb AJ, Patel N, Loukogeorgakis S, et al. Acute blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 2008;51(03): 784–790
- 27 Carlström M, Persson AEG, Larsson E, et al. Dietary nitrate attenuates oxidative stress, prevents cardiac and renal injuries, and reduces blood pressure in salt-induced hypertension. *Cardiovasc Res* 2011;89(03):574–585
- 28 Willis GR, Udiawar M, Evans WD, Blundell HL, James PE, Rees DA. Detailed characterisation of circulatory nitric oxide and free radical indices—is there evidence for abnormal cardiovascular homeostasis in young women with polycystic ovary syndrome? *BJOG* 2014;121(13):1596–1603

- 29 Anderson RA, Bundhoo S, James PE. A new mechanism of action of thienopyridine antiplatelet drugs - a role for gastric nitrosothiol metabolism? *Atherosclerosis* 2014;237(01):369–373
- 30 Richardson G, Hicks SL, O'Byrne S, et al. The ingestion of inorganic nitrate increases gastric S-nitrosothiol levels and inhibits platelet function in humans. *Nitric Oxide* 2002;7(01):24–29
- 31 Liu T, Schroeder HJ, Wilson SM, et al. Local and systemic vasodilatory effects of low molecular weight S-nitrosothiols. *Free Radic Biol Med* 2016;91:215–223
- 32 Ingram TE, Pinder AG, Bailey DM, Fraser AG, James PE. Low-dose sodium nitrite vasodilates hypoxic human pulmonary vasculature by a means that is not dependent on a simultaneous elevation in plasma nitrite. *Am J Physiol Heart Circ Physiol* 2010;298(02): H331–H339
- 33 Kwok CS, Loke YK. Effects of proton pump inhibitors on platelet function in patients receiving clopidogrel: a systematic review. *Drug Saf* 2012;35(02):127–139
- 34 Pinheiro LC, Amaral JH, Ferreira GC, et al. Gastric S-nitrosothiol formation drives the antihypertensive effects of oral sodium nitrite and nitrate in a rat model of renovascular hypertension. *Free Radic Biol Med* 2015;87:252–262
- 35 Arnette DR, Stamler JS. NO₂⁻, NO, and NO⁻ donation by S-nitrosothiols: implications for regulation of physiological functions by S-nitrosylation and acceleration of disulfide formation. *Arch Biochem Biophys* 1995;318(02):279–285
- 36 Burnley-Hall N, Willis G, Davis J, Rees DA, James PE. Nitrite-derived nitric oxide reduces hypoxia-inducible factor 1 α -mediated extracellular vesicle production by endothelial cells. *Nitric Oxide* 2017;63:1–12
- 37 Wang J-M, Wang Y, Huang J-Y, et al. C-Reactive protein-induced endothelial micro particle generation in HUVECs is related to BH4-dependent NO formation. *J Vasc Res* 2007;44(03):241–248
- 38 Montoro-García S, Shantsila E, Marín F, Blann A, Lip GY. Circulating micro particles: new insights into the biochemical basis of micro particle release and activity. *Basic Res Cardiol* 2011;106(06):911–923
- 39 Bundhoo S, Sagan E, James PE, Anderson RA. Clopidogrel results in favourable changes in nitric oxide metabolism in patients undergoing percutaneous coronary intervention. *Thromb Haemost* 2014;111(02):373–374
- 40 Dangelmaier C, Jin J, Smith JB, Kunapuli SP. Potentiation of thromboxane A₂-induced platelet secretion by Gi signalling through the phosphoinositide-3 kinase pathway. *Thromb Haemost* 2001;85(02):341–348
- 41 Flaumenhaft R. Molecular basis of platelet granule secretion. *Arterioscler Thromb Vasc Biol* 2003;23(07):1152–1160
- 42 Arima T, Ohshima Y, Mizuno T, Kitamura Y, Segawa T, Nomura Y. Cyclic GMP elevation by 5-hydroxytryptamine is due to nitric oxide derived from endogenous nitrosothiol in NG108-15 cells. *Biochem Biophys Res Commun* 1996;227(02):473–478
- 43 Gambaryan S, Geiger J, Schwarz UR, et al. Potent inhibition of human platelets by cGMP analogs independent of cGMP-dependent protein kinase. *Blood* 2004;103(07):2593–2600
- 44 Gurbel PA, Tantry US. Drug insight: Clopidogrel non responsiveness. *Nat Clin Pract Cardiovasc Med* 2006;3(07):387–395
- 45 Brose KMJ, Lee AYY. Cancer-associated thrombosis: prevention and treatment. *Curr Oncol* 2008;15(Suppl 1):S58–S67
- 46 Tutuian R, Katz PO, Bochenek W, Castell DO. Dose-dependent control of intragastric pH by pantoprazole, 10, 20 or 40 mg, in healthy volunteers. *Aliment Pharmacol Ther* 2002;16(04): 829–836
- 47 Shin JS, Lee JY, Cho KH, et al. The pharmacokinetics, pharmacodynamics and safety of oral doses of ilaprazole 10, 20 and 40 mg and esomeprazole 40 mg in healthy subjects: a randomised, open-label crossover study. *Aliment Pharmacol Ther* 2014;40(05):548–561
- 48 Gan KH, Geus WP, Lamers CB, Heijerman HG. Effect of omeprazole 40 mg once daily on intraduodenal and intragastric pH in H. pylori-negative healthy subjects. *Dig Dis Sci* 1997;42(11): 2304–2309
- 49 Ghobrial J, Gibson CM, Pinto DS. Delayed clopidogrel transit during myocardial infarction evident on angiography. *J Invasive Cardiol* 2015;27(05):E68–E69