

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/115187/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Pugh, Richard, Harrison, Wendy, Battle, Ceri, Hancock, Chris and Szakmany, Tamas 2017. Ventilator-associated events...perhaps not the answer. *Journal of the Intensive Care Society* 18 (3), pp. 263-264. 10.1177/1751143717694917 file

Publishers page: <http://dx.doi.org/10.1177/1751143717694917>
<<http://dx.doi.org/10.1177/1751143717694917>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Ventilator associated events... perhaps not the answer

Richard Pugh, Wendy Harrison, Ceri Battle, Chris Hancock, Tamas Szakmany

In the most recent issue of JICS, Dr Thomas has rightly argued that ventilator-associated pneumonia (VAP) can be difficult to diagnose and that efforts to limit ventilator-associated morbidity must extend beyond VAP prevention (1). We share his concerns, but are unconvinced that adopting the CDC NHSN ventilator-associated event (VAE) surveillance system recently introduced in the United States (2) is currently the right approach for our population (3).

In Wales, there has been a national surveillance programme to identify episodes of VAP since 2008, originally using the Hospitals in Europe Link for Infection Control through Surveillance definitions (HELICS, classified as PN 1 to 5, according to radiological, systemic, pulmonary and microbiological criteria). We recently investigated triggers for initiating antibiotic therapy for suspected ventilator-associated respiratory tract infection (VARTI) and the implications of variation in clinical practice for VAP surveillance (4). Among 282 invasively ventilated patients admitted to ICU for 48 hours or more, 33 developed VARTI – the main features having been volume and purulence of sputum, inflammatory markers and radiographic changes. In less than 50% cases was chest radiography performed at time of diagnosis, which precludes diagnosis of VAP according to HELICS definition – though these episodes were associated with a significantly prolonged duration of mechanical ventilation and ICU stay.

However, applying the new CDC NHSN definitions to our cohort, we found no overlap between VAE and suspected VARTI. Outside of the UK, other investigators have also described potential limitations to VAE surveillance. A ventilator-associated complication (VAC) is identified when there is sustained increase in FiO₂ or PEEP after a stable baseline of 48 hours or more. However, many patients do not meet the definition of VAC because fluctuations in FiO₂ and PEEP prevent them achieving this stable baseline (5). Variations in practice may affect whether a patient develops a VAC depending on preference for PEEP level or ventilator mode, such as APRV. Previous studies have also found low level of agreement between episodes of VAC and VAP; indeed patients meeting VAC criteria have a range of pathology which might include VAP, but alternatively may represent fluid overload, ARDS or atelectasis. Interventions required to prevent and treat such heterogeneous pathology are likely to differ and it may be difficult to understand the contribution of different processes to patient outcomes associated with VAC.

There are distinctions between the diagnostic needs of the clinician at the bedside and the surveillance administrator - and it is possible to perceive a risk of disengagement with a quality improvement programme that does not emphasise clinical discrimination. Furthermore, although VAE data is highly objective, in a paper-based clinical environment, the workload associated with screening should not be underestimated. Our approach has not been to dispense with HELICS definitions of VAP at this stage, but – appreciating the variation in performance of chest x-ray and subjectivity in interpretation – to supplement with an additional category “PNO” where there is pulmonary, systemic and microbiological evidence of VARTI but without radiographic evidence. We look forward to reporting further on this issue.

References

1. Thomas M. Ventilator associated events should trump ventilator associated pneumonia. *Journal of the Intensive Care Society*, 2016; 17(4): 358.
2. Magill SS, Klompas M, Balk R, et al. Developing a new, national approach to surveillance for ventilator-associated events. *Crit Care Med* 2013;41(11):2467–75.
3. Pugh R, Szakmany T. Ventilator-associated pneumonia surveillance: complex, complicated and still searching for the right tool. *British Journal of Intensive Care* 2014; 24(1):20-25
4. Pugh R, Harrison W, Harris S, et al. Is HELICS the right way? Lack of chest radiography limits ventilator-associated pneumonia surveillance in Wales. *Frontiers in Microbiology* 2016; 7:1271
5. Fan Y, Gao F, Wu Y, Zhang J, Zhu M, X L. Does ventilator-associated event surveillance detect ventilator-associated pneumonia in intensive care units? A systematic review and meta-analysis. *Critical Care* 2016; 20:338