

## Viewpoints

# The Cognitive Thalamus as a Gateway to Mental Representations

 Mathieu Wolff<sup>1,2</sup> and  Seralynne D. Vann<sup>3</sup>

<sup>1</sup>Centre National de la Recherche Scientifique, INCIA, Unité Mixte de Recherche 5287, Bordeaux, France, <sup>2</sup>University of Bordeaux, INCIA, Unité Mixte de Recherche 5287, Bordeaux, France, and <sup>3</sup>School of Psychology, Cardiff University, Cardiff, CF10 3AT, United Kingdom

Historically, the thalamus has been viewed as little more than a relay, simply transferring information to key players of the cast, the cortex and hippocampus, without providing any unique functional contribution. In recent years, evidence from multiple laboratories researching different thalamic nuclei has contradicted this idea of the thalamus as a passive structure. Dated models of thalamic functions are being pushed aside, revealing a greater and far more complex contribution of the thalamus for cognition. In this *Viewpoints* article, we show how recent data support novel views of thalamic functions that emphasize integrative roles in cognition, ranging from learning and memory to flexible adaption. We propose that these apparently separate cognitive functions may indeed be supported by a more general role in shaping mental representations. Several features of thalamocortical circuits are consistent with this suggested role, and we highlight how divergent and convergent thalamocortical and corticothalamic pathways may complement each other to support these functions. Furthermore, the role of the thalamus for subcortical integration is highlighted as a key mechanism for maintaining and updating representations. Finally, we discuss future areas of research and stress the importance of incorporating new experimental findings into existing knowledge to continue developing thalamic models. The presence of thalamic pathology in a number of neurological conditions reinforces the need to better understand the role of this region in cognition.

## Introduction

For over half a century, learning and memory have been intimately associated with the hippocampal formation, often leaving the functional contribution of other brain regions overlooked. However, the thalamus also has a long-standing link to memory. Indeed, damage within this region invariably occurs in Korsakoff syndrome, one of the key symptoms of which is a dense amnesia (Kopelman et al., 2009). The co-occurring diencephalic damage in this condition was noted as early as the end of the 19th century (Gudden, 1896), with a more explicit link between diencephalic damage and memory subsequently made by Gamper (1928). However, it was not until later in the 20th century that these brain regions began to gain further interest, by which time there was already a widespread focus on the medial temporal lobe for memory function, following the reports of Patient H.M. (Delay and Brion, 1969; Victor et al., 1971; Scoville and Milner, 2000). Thus, although the severity of memory impairments is often largely

comparable between temporal lobe and diencephalic amnesia (e.g., Hunkin et al., 1994; Shaw and Aggleton, 1995; Caulo et al., 2005), the role of the thalamus, and the diencephalon in general, has largely been disregarded. In a similar vein, the neural bases of reasoning, thought, and cognition are generally considered to be supported by the cortex, the prefrontal cortex (PFC) in particular (Donoso et al., 2014), with little, if any, role for subcortical areas. Thus, in terms of cognition, the thalamus has typically been viewed as a supporting member of the cast that acts simply as a relay for the main players (i.e., the hippocampus and the neocortex). Within these models, the thalamus has taken on a passive role, simply transferring information without providing any unique contribution to the system. In recent years, however, evidence has emerged that contradicts this idea of a passive relay and highlights a central role for the thalamus in cognition.

There are inherent difficulties in attempting to generate global models of thalamic functions because the thalamus is not a unitary structure. It comprises a large number of nuclei, each with different anatomical connectivity and functional properties. The idea of the thalamus as a heterogeneous structure with only a small number of nuclei supporting the canonical sensory-motor relay function was first established by the pioneering work of Guillery and Sherman (Sherman and Guillery, 1996). These authors further developed their model over the years, proposing a dichotomy of thalamic functions based primarily on the main type of afferents received by thalamic nuclei (for a comprehensive perspective of this work and of the major contribution of Ray Guillery, who sadly passed away last year, see Murray Sherman, 2018). Those nuclei receiving driver input (i.e., capable of directly eliciting neuronal activity) from the cortex are called higher-

Received June 27, 2018; revised Oct. 24, 2018; accepted Oct. 28, 2018.

M.W. was supported by Independent Investigator National Alliance for Research on Schizophrenia and Depression Grant 27402 and French Agency for Research THALAME Grant ANR-14-CE13-0029. S.D.V. was supported by Wellcome Trust Senior Research Fellowship in Biomedical Sciences WT090954AIA. We thank John Aggleton for helpful discussions.

The authors declare no competing financial interests.

Correspondence should be addressed to either of the following: Dr. Mathieu Wolff, Centre National de la Recherche Scientifique, INCIA, Unité Mixte de Recherche 5287, Bordeaux, France, E-mail: mathieu.wolff@u-bordeaux.fr; or Dr. Seralynne D. Vann, School of Psychology, Cardiff University, Cardiff, CF10 3AT, United Kingdom, E-mail: vannsd@cardiff.ac.uk.

<https://doi.org/10.1523/JNEUROSCI.0479-18.2018>

Copyright © 2019 Wolff et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License Creative Commons Attribution 4.0 International, which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

order nuclei and are thought to actively participate in cortical functioning (Sherman, 2016). In contrast, thalamic nuclei receiving driver input from subcortical regions are considered first-order thalamic nuclei (i.e., textbook relay thalamic nuclei). Other researchers characterized some thalamic nuclei as limbic on the basis of their connectivity with the cingulate cortex and their contribution to cognition rather than purely sensory-motor processes (Vogt and Gabriel, 1993). Both higher-order and limbic thalamic nuclei appear necessary for cognition (Vogt and Gabriel, 1993; Varela, 2014), but neither classification includes all nuclei that support this role; therefore, from a behavioral perspective, the term “cognitive thalamus” more accurately captures the essence of those thalamic nuclei that primarily support cognitive functions.

In this *Viewpoints* article, we will describe a revised model of the thalamus wherein, instead of merely acting as relays, thalamic nuclei contribute to cortical functioning and higher-order cognition, ranging from learning and memory to flexible adaptation. We will discuss the possibility that these apparently separate cognitive functions may indeed be supported by a more general role of the thalamus in maintaining and updating mental representations. The anterior thalamic nuclei (ATn) and the mediodorsal thalamus (MD) will serve as the main examples to illustrate this view. Given that extensive reviews of these thalamic nuclei are available (Bradfield et al., 2013a; Jankowski et al., 2013; Aggleton and Nelson, 2015; Dillingham et al., 2015a; Mitchell, 2015; Wolff et al., 2015a; Ouhaz et al., 2018; Pergola et al., 2018), our aim is not to give a detailed analysis of these areas but to highlight general functional principles that may transcend specific nuclei and so be relevant for the cognitive thalamus as a whole. We will also consider the role of corticothalamic versus thalamocortical projections and the integration of thalamocortical loops with other cortical and subcortical networks. In doing so, we hope to provide a general overview of the current state of knowledge and to identify areas where future research is needed.

### The cognitive thalamus


#### *Learning and memory*

Memory was probably the first cognitive function formally associated with the thalamus. Both the ATn and MD have been implicated in the memory impairments associated with the Korsakoff syndrome (Victor et al., 1971; Harding et al., 2000), but their individual contributions to learning and memory appear quite different (Bradfield et al., 2013a; Mitchell, 2015; Wolff et al., 2015a). Indeed, it had been proposed that a double dissociation existed between ATn and MD functions, with ATn supporting recollective memory and MD supporting familiarity-based memory (Aggleton and Brown, 1999). Although recent data suggest that this model does not entirely capture the mnemonic contribution of the MD (Danet et al., 2017), experimental manipulations in rodents have established clear distinctions between the types of memory processes supported by these different thalamic regions (Bradfield et al., 2013a; Wolff et al., 2015a).

ATn lesions in rodents produce striking impairments across spatial memory tasks, with the severity of deficit often comparable with that seen following hippocampal lesions (Warburton and Aggleton, 1999; Aggleton and Nelson, 2015). Impairments are found on reference and working memory, as well as path integration tasks (Warburton et al., 1997; Warburton and Aggleton, 1999; Frohardt et al., 2006). Thus, ATn lesions appear to disrupt the processing of environmental cues and the updating and monitoring of the animal's position within the environment. These spatial impairments are consistent with the electrophysio-

logical properties of the ATn, as this structure contains a number of spatially responsive cells encoding information, such as orientation, spatial location, and running speed (Taube, 1995; Tsanov et al., 2011b; Jankowski et al., 2015; Laurens et al., 2016). In contrast, the recognition of single items does not appear to require the ATn, although the ATn may be important for reducing interference between multiple similar items (Law and Smith, 2012; Nelson and Vann, 2017). Furthermore, ATn lesion-induced impairments are found when animals are required to combine item memory with additional features, such as temporal order and location (Parker and Gaffan, 1997; Wilton et al., 2001; Wolff et al., 2006; Dumont and Aggleton, 2013; Nelson and Vann, 2014). While a similar pattern of deficits can be found following MD lesions (Cross et al., 2012) MD lesions also impair the ability to discriminate the temporal order of two items, whereas temporal order memory impairments following ATn lesions only emerge when multiple items are used (Mitchell and Dalrymple-Alford, 2005; Nelson and Vann, 2017).

Although MD lesions can disrupt performance on spatial memory tasks, this does not appear to arise from impairments of spatial memory per se, but rather from impairments of strategic aspects of the task (Hunt and Aggleton, 1998a). There is an ongoing assumption that MD is particularly important for working memory because of its connections with the PFC (Watanabe and Funahashi, 2012; Funahashi, 2013; Halassa and Kastner, 2017; Parnaudeau et al., 2018) and because delay-dependent cells are found in the primate MD (Funahashi, 2013). Cells displaying delay-dependent activity have also been found in the rodent MD, but the findings are far more variable with some studies showing activity at delays comparable with cells within the dorsomedial PFC (dmPFC) (Bolkan et al., 2017) and others showing no delay activity (Han et al., 2013; Miller et al., 2017). Although MD damage in primates disrupts working memory, these deficits are most often found in combination with other memory or executive deficits, suggesting that working memory itself may not be specifically compromised (Watanabe and Funahashi, 2012; Baxter, 2013). In rodents, data from delayed nonmatching-to-place tasks (i.e., spatial alternation) bring little support for the idea that the MD contributes to working memory. This task takes advantage of rodents' natural tendency to search in novel locations for food and requires rats to alternate between arms of a T-shaped maze, often for a reward. Although this behavioral task appears simple, it indeed relies on multiple cognitive processes and can be solved using several different strategies (Dudchenko, 2001). Deficits can thus reflect poor spatial memory, but several other factors can also affect performance, including impulsivity, motivation, reward-response associations, and interference sensitivity. When impaired performance is observed after MD damage, it has often been reported as transient or nonspecific (Stokes and Best, 1990; Hunt and Aggleton, 1991, 1998a; Alexinsky, 2001; Chauveau et al., 2005). Importantly, several experiments performed in different laboratories found delayed nonmatching-to-place performance to be unaffected after thalamic damage, even when damage was substantial and long delays were included (Neave et al., 1993; Hunt and Aggleton, 1998b; Mitchell and Dalrymple-Alford, 2005; Alcaraz et al., 2016b). Experimental data supporting the opposite view (i.e., a role for the MD in delayed nonmatching-to-place), mostly come from recent chemogenetic and optogenetic interventions conducted in mice in which impairments were found at longer delays (Parnaudeau et al., 2013; Bolkan et al., 2017), as well as during the acquisition of the nonmatching-to-place task (Parnaudeau et al., 2013). The apparent discrepancy between these findings and earlier studies may



**Figure 1.** Different degrees of complexity for thalamocortical architecture. Basically defined as the reciprocal projections between prefrontal and thalamic areas (**A**), the thalamocortical loop includes an additional layer constituted by the reticular thalamic nucleus (TRN, **B**). This area is one of the main sources of thalamic inhibition. Both thalamocortical and corticothalamic pathways send collaterals to the TRN. In addition, the TRN sends supplemental inhibitory projections to other thalamic nuclei (**C**), not included in the actual loop, thus opening this loop, which may allow gating of specific thalamocortical inputs (see also Fig. 3).

arise for a number of reasons. The most pronounced impairments in these mouse studies are found during longer delays in well-trained animals and thus might reflect impairments in additional factors, such as impulsivity. The specificity of viral spread within the thalamus may also be an issue in mice, and potential encroachment into adjacent thalamic nuclei, such as the ATn, could contribute to the findings (Hunt and Aggleton, 1998b; Mitchell and Chakraborty, 2013; Aggleton and Nelson, 2015; Wolff et al., 2015a), given that damage to the ATn, but not MD, severely impairs spatial working memory (Alcaraz et al., 2016b). Together, the overall picture appears to be that MD is not necessary for working memory but may contribute to additional aspects of task performance, such as delay monitoring or habit formation when animals are overtrained.

#### Shaping mental representations

Decades ago, Tolman (1948) coined the term “cognitive map” to refer to a highly organized knowledge database that allows flexible actions. Cognitive maps can be considered mental representations requiring the combination of external cues with internal states to generate accurate depictions of general rules and/or associative laws. These representations are vital for animals to successfully interact with the world (Ramos, 2014). Although it is possible to dissociate thalamic nuclei on the basis of their distinct cognitive functions (Mitchell and Dalrymple-Alford, 2005, 2006; Wolff et al., 2008b, 2015a, b; Bradfield et al., 2013a; Moreau et al., 2013), this does not contradict the idea of an overall involvement of the cognitive thalamus in shaping mental representations. For instance, while the ATn and the MD belong to distinct functional circuits, they are both considered important for directing attention to task-relevant behavioral features (Wolff et al., 2015b; Wright et al., 2015), which is required to build task-relevant mental constructs. Moreover, thalamic damage often impairs memory acquisition, suggesting further that forming meaningful representation requires thalamic integrity (Cermak et al., 1980; Vann and Aggleton, 2003; Wolff et al., 2008b; Marchand et al., 2014; Sweeney-Reed et al., 2014).

Even after initial learning is established, the thalamus continues to play an important role, possibly by monitoring and updating current information within a changing environment. For example, ATn damage appears to be particularly detrimental when elements of flexibility are required to solve ongoing challenges: the ability to reach a previously learned position from a new start is disproportionately impaired by ATn lesions (Wolff et al., 2008a), as is spatial alternation when the animal is released from opposite arms for sample and test trials (Warburton et al., 1997; Loukavenko et al., 2007, 2016). A common feature of these experimental situations is that animals must track changes in task demands and update their current frame of reference accordingly

to maintain successful performance. The MD also appears to be particularly important when successful performance requires the update of action-outcome or stimulus-outcome associations, as shown in rodents (Corbit et al., 2003; Ostlund and Balleine, 2008; Bradfield et al., 2013a; Parnaudeau et al., 2015; Alcaraz et al., 2016b, 2018) and also in primates (Mitchell et al., 2007; Izquierdo and Murray, 2010; Browning et al., 2015; Chakraborty et al., 2016; Wicker et al., 2018).

Together, these data highlight a role for thalamic nuclei in monitoring, maintaining, and updating mental constructs, in contrast to previous views, which have emphasized the dominant role of cortical areas (Wilson et al., 2010, 2014; Markov et al., 2013). Increasing evidence indicates instead that close functional interactions between cortical and thalamic areas are essential to shape these representations to address ongoing challenges (Cross et al., 2012; Parnaudeau et al., 2013; Browning et al., 2015; Bolkan et al., 2017; Miller et al., 2017; Schmitt et al., 2017; Alcaraz et al., 2018; Marton et al., 2018). To better understand the nature of these interactions, it is necessary to consider specific features of the organization of thalamocortical circuits.

#### The thalamocortical loop

One hallmark of thalamocortical circuits is the reciprocity of projections between cortical and thalamic areas. This has been viewed as “reentry,” a process whereby two or more brain regions concurrently stimulate, and are stimulated by, each other. This reciprocal and parallel processing supports the synchronization of neuronal firing required for rapid neural integration. The binding of activity across a number of regions is thought to underpin the conscious processing of stimuli, which is necessary to form a unified mental construct (e.g., a scene or visual representation) (Tononi and Edelman, 1998; Tononi et al., 1998; Edelman and Gally, 2013). As a result of recent technical advances, it is now possible to selectively target projection-defined neurons, which has opened up new possibilities in assessing the functional role of thalamocortical *versus* corticothalamic pathways. Two recent studies have used this approach and have shown that reciprocal pathways between MD and the dmPFC can be functionally differentiated (Bolkan et al., 2017; Alcaraz et al., 2018). Thus, thalamocortical and corticothalamic pathways may play complementary but dissociable roles in cognition. Unlike thalamocortical projections, which are mostly ipsilateral, corticothalamic projections also provide substantial contralateral innervation at the thalamic level (Preuss and Goldman-Rakic, 1987; Négyessy et al., 1998; Bradfield et al., 2013a; Mathiasen et al., 2017). These projections include collaterals to the reticular thalamic nucleus, which in turn provides lateral inhibition for virtually any thalamic nucleus (Pinault, 2004; Halassa and Acsády, 2016) (Fig. 1). The functional relevance of this organization is discussed below.

#### Corticothalamic pathways: directing cognitive resources

A cardinal feature of higher-order thalamic nuclei is that they receive both a modulatory input and a driver input from the cortex (from layers 6 and 5, respectively) (Usrey and Sherman, 2018). This organization suggests important functional roles for corticofugal pathways, possibly implementing additional and indirect corticocortical routes through the thalamus (Jones, 1998; Sherman, 2005, 2012). This view is, however, largely speculative

and mostly derived from neurophysiological studies of sensory-motor functions (Sherman, 2016). But even sensory mechanisms can contribute to cognition: they may be viewed as enabling abstraction of relevant information, thus helping to represent the external world in a meaningful way (Cudeiro and Sillito, 2006). Branching of thalamocortical pathways at the level of the reticular thalamic nucleus may enable the gating of salient thalamic inputs by minimizing the importance of those that are currently irrelevant (McCormick and von Krosigk, 1992; Zikopoulos and Barbas, 2007; Stillova et al., 2015), thus providing a possible mechanism of focused attention (Béhuret et al., 2015; Wimmer et al., 2015). This view is rooted in the ideas initially developed by Crick (1984) of the reticular thalamic nucleus acting as an attentional searchlight. Interestingly, increased modulation of corticothalamic pathways has been found to parallel increased attentional demand in humans (Jagtap and Diwadkar, 2016). The dynamic nature of the excitatory-inhibitory balance at the thalamic level depends on current behavioral demand, with critical dependence on corticothalamic pathways and their collaterals to the reticular thalamic nucleus (Crandall et al., 2015; Li and Ebner, 2016). It is thus possible that cortical projections to the thalamus directly adjust the gain and the tuning precision of thalamic cells as required by ongoing behavioral demands (Mease et al., 2014; Guo et al., 2017a).

Beyond their role in directing attentional resources, corticothalamic pathways have also been linked to processes underpinning learning. For example, direct evidence for a causal involvement of corticothalamic pathways in learning has been reported in an appetitive Pavlovian conditioning task. Optogenetic manipulation of the projections from PFC to the paraventricular thalamic nucleus during task acquisition affected the conditioned response, highlighting a role for this pathway in the encoding of predictive environmental cues (Otis et al., 2017). Furthermore, dmPFC-to-MD pathways have been demonstrated to support upcoming choice either in a spatial working memory task (Bolkan et al., 2017) or when the retrieval of current goal value is required for successful responding (Alcaraz et al., 2018). Importantly, these corticothalamic pathways also promote behavioral flexibility (Nakayama et al., 2018), especially when rule switching is required (Marton et al., 2018). Collectively, these data suggest a central role for corticothalamic pathways in cognition, and their functional relevance seems to range from directing attention to solving cognitive challenges.

#### *Thalamocortical pathways: more than a relay*

Central to almost all definitions of thalamic function is the concept that this region is a “relay.” Even those nuclei that are considered to have a more cognitive role are still considered to be principally involved in relaying information either between cortical sites or between medial temporal lobe and neocortex. This description of thalamic function attributes little or no additional role for these nuclei other than acting as a waystation. However, this clearly underestimates and oversimplifies the role of the thalamus. The idea of thalamic regions monitoring and updating information and providing an active contribution rather than a passive relay is not in fact new. The MD was previously suggested to be involved in mediating cognitive aspects of odor-guided tasks rather than transmitting sensory information (Eichenbaum et al., 1980). The latter possibility was considered because the MD links the piriform cortex (the primary olfactory cortex) with the orbitofrontal associative cortex (Courtiol and Wilson, 2015). But even when using an odor-guided behavioral assay, it appears that

task-related features, rather than purely sensory information, are represented by MD cells (Courtiol and Wilson, 2016).

More recently, other evidence has emerged that also supports the idea of nonrelay contributions of the thalamus. For example, Schmitt et al. (2017) recently showed that the MD is able to sustain cortical representations rather than relaying information. These data emphasize a role for the thalamus in controlling cortical connectivity to maintain rule representation (Halassa and Kastner, 2017; Nakajima and Halassa, 2017). A causal relationship between MD-PFC activity and social dominance behavior was also recently established, underscoring further the importance of thalamic inputs for cortical functions (Zhou et al., 2017). It should be noted that the importance of sustained thalamocortical activity during delay has been observed to instruct future actions in other thalamocortical circuits (Guo et al., 2017b), suggesting that sustaining cortical activity may constitute an essential role of thalamic inputs.


Like the MD, the ATn does not passively relay information. Instead these nuclei show long-term, input-dependent modification of their responses, which can amplify the convergent inputs from different sources (Tsanov et al., 2011a). Behavioral data are also consistent with a nonrelay function for the ATn: the behavioral effects of ATn lesions can be more pronounced than lesions disrupting any of their individual inputs (e.g., Aggleton et al., 1995; Warburton and Aggleton, 1999; Sziklas and Petrides, 2000; Wright et al., 2015; Powell et al., 2017), suggesting that no single pathway supports all cognitive aspects of ATn function. Therefore, it is unlikely that the ATn is a simple relay in the traditional sense, but instead integrates information from midbrain, diencephalic, hippocampal, and cortical regions (Tsanov et al., 2011a; Vann and Nelson, 2015; Mathiasen et al., 2017). This view is also supported by numerous studies showing the importance of ATn inputs for driving activity in their cortical target, the retrosplenial cortex. In rodents, ATn lesions disrupt a number of markers of activity in the retrosplenial cortex (Dupire et al., 2013; Mendez-Lopez et al., 2013; Aggleton and Nelson, 2015). Furthermore, the retrosplenial cortex is hypoactive in patients with thalamic damage (Reed et al., 2003). Importantly, retrosplenial cortex activity changes are not simply a result of deafferentation (Garden et al., 2009; Vann and Albasser, 2009; Frizzati et al., 2016), but likely reflect the loss of functional coupling between the ATn and retrosplenial cortex. Indeed, the close functional correspondence between the ATn, hippocampus, and retrosplenial cortex highlights the possible importance of the thalamus in synchronizing activity across multiple regions (Corcoran et al., 2016; Eichenbaum, 2017; Halassa and Kastner, 2017). Indeed, this function may be paramount for its role in updating existing representations.

#### **Beyond the thalamocortical loop**

At this point, it seems appropriate to expand more broadly our views of thalamic functions, looking beyond the thalamocortical loop. How exactly do these loops integrate with other cortical and subcortical circuits?

#### *Cortical integration*

Thalamocortical projections are both divergent and convergent (Rubio-Garrido et al., 2009). This feature of thalamocortical architecture provides an ideal basis for integration both within and across cortical regions (Fig. 2). Keeping with the example of the MD, several parallel thalamocortical pathways originate in MD and target distinct prefrontal areas (Groenewegen, 1988; Alcaraz et al., 2016a). In rodents, there is a clear topography as lateral MD neurons innervate the dorsal wall of the PFC while medial MD neu-



**Figure 2.** In addition to the reciprocity of projections, divergence (**A**) and convergence (**B**) are two prominent features of thalamocortical organization. Whereas the former underscores the possibility that multiple thalamic microcircuits act in parallel to achieve cognitive functions, the latter offers enhanced integrative properties. Regarding convergence, MD innervation of the dmPFC and the OFC originates from essentially separate neuronal populations (light blue), even though some MD cells branch to several prefrontal areas. Sub, Submedial thalamic nucleus.

rons predominantly contact its ventral wall. In contrast, MD cells located in the central segment essentially innervate the orbitofrontal cortex (OFC) (Alcaraz et al., 2016a; Murphy and Deutch, 2018). In addition to this distinct topography across MD, individual MD neurons also innervate several PFC regions, with collaterals contacting multiple cortical layers (Kuramoto et al., 2017b). Together, this illustrates the highly divergent nature of these thalamocortical projections.

By targeting different cell types and different neuronal compartments (Delevich et al., 2015; Collins et al., 2018), the MD can influence varying aspects of PFC functioning. For example, the MD projections to the dmPFC and the OFC have been proposed to support instrumental and Pavlovian associations, respectively (Ostlund and Balleine, 2008; Alcaraz et al., 2016a). An intriguing possibility is that such organization enables simultaneous instruction of distinct cortical areas through different pathways, further highlighting the multiple thalamic microcircuits that may act in parallel to support cognition (Rikhye et al., 2018).

Moving onto convergence, cortical regions are typically innervated by more than one thalamic nucleus. The PFC receives afferent projections from a large number of thalamic nuclei, including the intralaminar, midline, and anterior nuclei (Barbas et al., 1991). Taking the OFC as an example, the main thalamic efferents to the OFC originate from the MD and the much-less-known submedial thalamic nucleus (Yoshida et al., 1992; Alcaraz et al., 2016a; Kuramoto et al., 2017a). At present, the functional relevance of this type of organization is unclear, although both thalamic nuclei appear to support successful updating of stimulus–outcome associations, a cardinal function of the OFC (Ostlund and Balleine, 2008; Alcaraz et al., 2015). At a cellular level, however, it is not known whether the inputs from MD and the submedial thalamic nucleus actually converge on single cortical neurons. Convergence onto a single cortical cell could account for the synergistic amplification of signals, which may be particularly important for accentuating behaviorally relevant environmental features. The more general process of neuronal convergence would encourage the integration of different information streams, enabling the development of more detailed multisensory mental representations (Man et al., 2013).

Disentangling the functional contribution of diverging and converging thalamic inputs will be an important objective for future research, especially considering that these features of thalamocortical organization appear to be largely conserved between species (Padberg et al., 2009).

#### Subcortical integration

While interactions between the thalamus and cortex are critical for cognition, there is increasing evidence that thalamic nuclei

may also have a role in integrating subcortical information. The ATn receives inputs from the hippocampal formation directly, mainly via the fornix (Dillingham et al., 2015b), as well as indirectly, via the mammillary bodies. However, this does not result in redundancy or a replication of information because the inputs are from distinct hippocampal populations (Amin et al., 2010; Christiansen et al., 2016), suggesting that the ATn may have a role in combining these separate hippocampal inputs. Projections from the mammillary bodies predominantly act as drivers to the ATn, whereas the cortical/hippocampal inputs have a modulatory role, specifically in the case of the anterodorsal nucleus (Somogyi et al., 1978; Petrof and Sherman, 2009). Consistent with this distinction between inputs, direct hippocampal inputs to the ATn elicit long-term depression, whereas the projections from the mammillary bodies elicit long-term potentiation (Tsanov et al., 2011a).

In addition to being a convergence point for separate hippocampal inputs, the ATn receives direct and indirect inputs from the midbrain tegmentum. The direct cholinergic input arises from the laterodorsal tegmental nucleus, whereas the indirect inputs come from the dorsal and ventral tegmental nuclei of Gudden. These midbrain inputs appear crucial for learning and memory (Mitchell et al., 2002; Taube, 2007; Vann, 2009; Clark et al., 2013; Vann, 2013), and the indirect inputs are critical for the head-direction and theta signals found in the ATn (Satoh and Fibiger, 1986; Bassant and Poindessous-Jazat, 2001; Kocsis et al., 2001; Bassett et al., 2007; Zakowski et al., 2017).

Other thalamic nuclei also exhibit a close partnership with subcortical areas, especially the striatum. The dorsal striatum is the recipient of a strong glutamatergic innervation originating from both the centro-median/parafascicular complex and the intralaminar group (Galvan and Smith, 2011). These thalamic groups are crucial for the role of the basal ganglia (BG) role in behavioral flexibility; they provide behaviorally relevant information and have been linked to goal-directed action selection in both rodents and humans (Brown et al., 2010; Kato et al., 2011; Liebermann et al., 2013; Schepers et al., 2017; Kato et al., 2018). For instance, removal of the parafascicular nucleus (Pf) in rats disrupted goal-directed behaviors by preventing rats from updating action–outcome associations (Bradfield et al., 2013b). Importantly, this manipulation was also found to reduce intrinsic activity of striatal cholinergic interneurons specifically, suggesting the existence of a direct link between behavioral flexibility and Pf inputs to these striatal cholinergic interneurons (Bradfield et al., 2013b; Yamanaka et al., 2018). Using cross-unilateral lesions of the Pf and striatum, Bradfield and Balleine (2017) showed that the impairment in updating action–outcome associations was due to an inability to use internal state to access the appropriate associations; in contrast, the use of external context to control accurate goal-directed action selection remained intact (Bradfield and Balleine, 2017). Lack of flexibility after inhibition of this thalamostriatal pathway is also evident in other paradigms, such as the five-choice serial reaction time task (Saund et al., 2017). These findings further highlight the importance of the thalamus in monitoring and updating mental representations in response to changing circumstances.

This latter consideration certainly calls for a critical assessment of how thalamocortical loops and the BG integrate. The thalamostriatal projection described above has only been minimally featured in classic models of BG functioning (Bostan and Strick, 2018). Instead, the emphasis has been on thalamic nuclei as a motor output, even though this does not address the contribution of thalamus-to-BG pathways (Goldberg et al., 2013). Such a model is supported by the fact that the motor thalamus, that is, the ventral anterior/ventral lateral thalamic areas, is one of the main recipients of BG output. But even this admittedly motor thalamic region has been shown to play a crucial role in performance monitoring (Seifert et al., 2011; Ullsperger et al., 2014).

The other main thalamic recipient of BG projections is the MD. This BG-MD pathway has also shown to be necessary for higher-order cognition (Leung and Balleine, 2015). As a result, a more integrative picture of the thalamus is now emerging, especially when considering the existence of multiple and interacting cortico-BG-thalamo-cortical loops (Haber and Calzavara, 2009; Bell and Shine, 2016). Interestingly, both corticostriatal and corticothalamic pathways were recently demonstrated to support behavioral flexibility, whereas corticocortical pyramidal prefrontal neurons were not (Nakayama et al., 2018), further demonstrating the key contribution of subcortical areas for high-order cognition. Together, these data suggest a critical role for the thalamus in integrating subcortical as well as cortical inputs to support cognitive functions.

#### *The thalamus: a bridge between the medial temporal lobe and frontal cortex*

To account for the myriad cognitive symptoms and memory impairments elicited by thalamic damage, an interesting early proposal was that the thalamus acts as a link between the medial temporal lobe and the frontal lobe (Warrington and Weiskrantz, 1982). Accordingly, disconnecting the hippocampal inputs to the ATn impairs performance on spatial memory tasks (Warburton et al., 2000). Furthermore, ATn damage in rodents and medial diencephalic damage in patients reduce activity in the frontal cortex (Kapur, 1994; Jenkins et al., 2002; Reed et al., 2003; Caulo et al., 2005; Vann and Albasser, 2009; Ozyurt et al., 2014). However, ATn lesion effects are unlikely to be due to the disruption of a one-way flow of information from the hippocampal formation to the frontal cortex as originally suggested. Indeed, the importance of the return projections from the ATn to the hippocampus has been recently highlighted (Vann and Nelson, 2015). For example, memory elicited event-related potentials in the ATn have been found to precede those in the hippocampus, suggesting a flow of information from ATn to the hippocampus, rather than the reverse (Stillova et al., 2015). The activity changes found in the frontal cortex following thalamic damage may therefore reflect a role for ATn in coordinating and synchronizing activity across hippocampal and cortical regions (Sweeney-Reed et al., 2014).

The MD could also be seen as link between the medial temporal lobe, as a recipient of projections originating from the BLA (technically a medial temporal lobe structure), and the OFC (Timbie and Barbas, 2015; Wolff et al., 2015a). It is therefore interesting to note that lesions to either MD or the BLA produce qualitatively different impairments when the contingency between predictive cues and their outcome is modified (Ostlund and Balleine, 2008). In that study, rats were trained to respond to two stimuli that reliably predicted a reward. Subsequently, one of the stimuli became noncontingently paired with reward, whereas the other remained a reliable predictor. After this manipulation,


rats with BLA lesions did not adapt their behavior and continued to exhibit a positive conditioned response to both contingent and noncontingent stimuli. In contrast, rats with MD lesions exhibited a nonspecific effect with reduced responding to both the noncontingent and contingent stimuli (i.e., they ceased to respond to cues that still reliably predicted their outcome) (Ostlund and Balleine, 2008). This suggests that adaptive cognition requires the functional contribution of multiple complementary pathways. Further work is thus warranted to examine the functional contribution of the connections between these areas and their cortical target, the OFC.

The reuniens nucleus (Re) is another thalamic nucleus that has recently received much attention due to its role in connecting prefrontal and temporal lobe areas: it projects to multiple frontal areas and is the main source of thalamic afferents to the hippocampus. Because all of these projections are reciprocal, the Re is thought to be a major hub, orchestrating functional exchanges between frontal areas and the hippocampus, especially in the absence of direct inputs from the PFC to the hippocampus (McKenna and Vertes, 2004; Vertes et al., 2006; Hoover and Vertes, 2012; Varela et al., 2014). Because of these unique properties, an unusually large number of studies have been conducted recently to assess the possible functions of these Re projections. There are a number of excellent review papers available on the subject (Cassel et al., 2013; Cassel and Pereira de Vasconcelos, 2015; Pereira de Vasconcelos and Cassel, 2015; Vertes, 2015). Briefly, Re dysfunction affects contextual fear memory (Xu and Sudhof, 2013; Vetere et al., 2017), impairs spatial memory consolidation (Loureiro et al., 2012), either increases or decreases behavioral flexibility (Cholvin et al., 2013; Linley et al., 2016; Prasad et al., 2017; Viena et al., 2018), impedes goal-directed spatial navigation (Ito et al., 2015), or delayed nonmatching-to-place (Hallock et al., 2013, 2016; Viena et al., 2018), and even affects visual threat responses (Salay et al., 2018). Given these diverse findings, continued research is needed before it is possible to generate global functional models of this region. Nevertheless, an interesting proposal, and a recurring theme across thalamic nuclei is that the Re may be particularly important for synchronizing PFC and hippocampal activity, facilitating functional exchanges between these areas (Eichenbaum, 2017). The wide array of functions that appear to be Re-dependent may therefore reflect the diverse nature of the tasks that require functional cooperation between PFC and hippocampal areas.

#### **Future directions**

At this point, it should be evident that cognitive functions are supported by distributed neural circuits among which thalamic nuclei play highly integrative roles. In this *Viewpoints* article, we examined recent evidence implicating thalamic nuclei in high-order cognitive functions and have attempted to highlight how many of these new findings support aspects of early cognitive views of thalamic functions (e.g., Warrington and Weiskrantz, 1982; Sherman and Guillery, 1996; Aggleton and Brown, 1999). We therefore propose that a central role for the cognitive thalamus is to shape mental representations, either by maintaining relevant mental constructs online or by updating those no longer relevant for ongoing challenges (Fig. 3). This proposition builds on many diverse experimental findings and focuses on general thalamic functions rather than specific contributions of individual nuclei.

The thalamus has been increasingly implicated in neurological disorders that present with cognitive dysfunction, including schizophrenia (Pinault, 2011; Uhlhaas et al., 2013; Anticevic et



**Figure 3.** *A*, Divergence, convergence, and gating are essential functional principles of thalamocortical circuits. Divergent and convergent thalamocortical pathways may promote parallel functioning and integrative processing at the cortical level, respectively. In addition, returning corticothalamic pathways are able to gate relevant thalamocortical inputs through lateral inhibition at the level of the TRN. This architecture may contribute to the maintenance (*B*) or updating (*C*) of cortical representations. In the latter case, some mental constructs may no longer be relevant (outlined in gray dashed line), whereas others become more prominent (blue solid arrows).

al., 2014), drug addiction (Balleine et al., 2015), Korsakoff syndrome (Victor et al., 1971; Harding et al., 2000), Alzheimer's disease (Braak and Braak, 1991), and Down syndrome (Karlsen et al., 2014; Perry et al., 2018). Therefore, new models of thalamic function may better explain the pattern of deficits associated with these conditions.

While we share the belief that thalamus research has entered a new era and that virtually any brain region has a thalamic story to tell (Acsády, 2017), we also anticipate the danger of being overwhelmed by data that may be difficult to replicate or to extend. Relying on standardized behavioral assays may be key, as is in-depth analysis of the literature. Technical developments progress at a much higher rate than conceptual advances. The latter can only be supported by a better analysis of how new data integrate with the preexisting knowledge. In doing so, conceptual views may be enriched, revised, or even drastically changed if novel technical approaches are used to test some of their predictions.

As discussed at several points throughout this article, there are areas of thalamic research where our current knowledge is woefully lacking. Filling these missing gaps could prove invaluable in testing and advancing current models of thalamic function. First, the functional relevance of divergent and convergent thalamocortical or corticothalamic pathways needs to be addressed more systematically. Until recently, this has been technically impossible due to methodological limitations. But by using a combination of chemogenetic and/or optogenetic approaches, it is now possible to test the relative contributions of these separate pathways. There is no single, definitive experiment to address this issue; instead, we believe the combined effort across multiple laboratories will be essential to advancements in this area. Second, although a large part of this *Viewpoints* article has emphasized functional interactions between the cortex and the thalamus, we also highlighted the importance of the thalamus for subcortical integration. There is a clear need for additional data to better appreciate how the thalamus interacts with the BG and other subcortical areas. Furthermore, the advances in multisite *in vivo* recordings will help address the contribution of the thalamus for coordinating and synchronizing brainwide activity. Finally, work on the thalamus has been performed either by researchers primarily interested in sensory-motor nuclei or by those interested in cognition. Surprisingly, these two worlds have been quite separate and relatively unaware of the progress made in the other field. We believe in the potential of uniting efforts in promoting a cognitive view of the thalamus, from sensory salience to adaptive cognition.

## References

- Acsády L (2017) The thalamic paradox. *Nat Neurosci* 20:901–902. [CrossRef Medline](#)
- Aggleton JP, Brown MW (1999) Episodic memory, amnesia, and the hippocampal-anterior thalamic axis. *Behav Brain Sci* 22:425–444; discussion 444–489. [Medline](#)
- Aggleton JP, Nelson AJ (2015) Why do lesions in the rodent anterior thalamic nuclei cause such severe spatial deficits? *Neurosci Biobehav Rev* 54:131–144. [CrossRef Medline](#)
- Aggleton JP, Neave N, Nagle S, Hunt PR (1995) A comparison of the effects of anterior thalamic, mamillary body and fornix lesions on reinforced spatial alternation. *Behav Brain Res* 68:91–101. [CrossRef Medline](#)
- Alcaraz F, Marchand AR, Vidal E, Guillou A, Faugère A, Coutureau E, Wolff M (2015) Flexible use of predictive cues beyond the orbitofrontal cortex: role of the submedial thalamic nucleus. *J Neurosci* 35:13183–13193. [CrossRef Medline](#)
- Alcaraz F, Marchand AR, Courtand G, Coutureau E, Wolff M (2016a) Parallel inputs from the mediodorsal thalamus to the prefrontal cortex in the rat. *Eur J Neurosci* 44:1972–1986. [CrossRef Medline](#)
- Alcaraz F, Naneix F, Desfosses E, Marchand AR, Wolff M, Coutureau E (2016b) Dissociable effects of anterior and mediodorsal thalamic lesions on spatial goal-directed behavior. *Brain Struct Funct* 221:79–89. [CrossRef Medline](#)
- Alcaraz F, Fresno V, Marchand AR, Kremer EJ, Coutureau E, Wolff M (2018) Thalamocortical and corticothalamic pathways differentially contribute to goal-directed behaviors in the rat. *eLife* 7:e32517. [CrossRef Medline](#)
- Alexinsky T (2001) Differential effect of thalamic and cortical lesions on memory systems in the rat. *Behav Brain Res* 122:175–191. [CrossRef Medline](#)
- Amin E, Wright N, Poirier GL, Thomas KL, Erichsen JT, Aggleton JP (2010) Selective lamina dysregulation in granular retrosplenial cortex (area 29) after anterior thalamic lesions: an *in situ* hybridization and transneuronal tracing study in rats. *Neuroscience* 169:1255–1267. [CrossRef Medline](#)
- Anticevic A, Cole MW, Repovs G, Murray JD, Brumbaugh MS, Winkler AM, Savic A, Krystal JH, Pearlson GD, Glahn DC (2014) Characterizing thalamo-cortical disturbances in schizophrenia and bipolar illness. *Cereb Cortex* 24:3116–3130. [CrossRef Medline](#)
- Balleine BW, Morris RW, Leung BK (2015) Thalamocortical integration of instrumental learning and performance and their disintegration in addiction. *Brain Res* 1628:104–116. [CrossRef Medline](#)
- Barbas H, Henion TH, Dermon CR (1991) Diverse thalamic projections to the prefrontal cortex in the rhesus monkey. *J Comp Neurol* 313:65–94. [CrossRef Medline](#)
- Bassant MH, Poindessous-Jazat F (2001) Ventral tegmental nucleus of Gudden: a pontine hippocampal theta generator? *Hippocampus* 11:809–813. [CrossRef Medline](#)
- Bassett JP, Tullman ML, Taube JS (2007) Lesions of the tegmentomammillary circuit in the head direction system disrupt the head direction signal in the anterior thalamus. *J Neurosci* 27:7564–7577. [CrossRef Medline](#)
- Baxter MG (2013) Mediodorsal thalamus and cognition in non-human primates. *Front Syst Neurosci* 7:38. [CrossRef Medline](#)
- Béhuret S, Deleuze C, Bal T (2015) Corticothalamic synaptic noise as a

- mechanism for selective attention in thalamic neurons. *Front Neural Circuits* 9:80. [CrossRef Medline](#)
- Bell PT, Shine JM (2016) Subcortical contributions to large-scale network communication. *Neurosci Biobehav Rev* 71:313–322. [CrossRef Medline](#)
- Bolkan SS, Stujenske JM, Parnaudeau S, Spellman TJ, Rauffenbart C, Abbas AI, Harris AZ, Gordon JA, Kellendonk C (2017) Thalamic projections sustain prefrontal activity during working memory maintenance. *Nat Neurosci* 20:987–996. [CrossRef Medline](#)
- Bostan AC, Strick PL (2018) The basal ganglia and the cerebellum: nodes in an integrated network. *Nat Rev Neurosci* 19:338–350. [CrossRef Medline](#)
- Braak H, Braak E (1991) Alzheimer's disease affects limbic nuclei of the thalamus. *Acta Neuropathol* 81:261–268. [CrossRef Medline](#)
- Bradfield LA, Balleine BW (2017) Thalamic control of dorsomedial striatum regulates internal state to guide goal-directed action selection. *J Neurosci* 37:3721–3733. [CrossRef Medline](#)
- Bradfield LA, Hart G, Balleine BW (2013a) The role of the anterior, mediodorsal, and parafascicular thalamus in instrumental conditioning. *Front Syst Neurosci* 7:51. [CrossRef Medline](#)
- Bradfield LA, Bertran-Gonzalez J, Chieng B, Balleine BW (2013b) The thalamostriatal pathway and cholinergic control of goal-directed action: interlacing new with existing learning in the striatum. *Neuron* 79:153–166. [CrossRef Medline](#)
- Brown HD, Baker PM, Ragozzino ME (2010) The parafascicular thalamic nucleus concomitantly influences behavioral flexibility and dorsomedial striatal acetylcholine output in rats. *J Neurosci* 30:14390–14398. [CrossRef Medline](#)
- Browning PG, Chakraborty S, Mitchell AS (2015) Evidence for mediodorsal thalamus and prefrontal cortex interactions during cognition in macaques. *Cereb Cortex* 25:4519–4534. [CrossRef Medline](#)
- Cassel JC, Pereira de Vasconcelos A (2015) Importance of the ventral midline thalamus in driving hippocampal functions. *Prog Brain Res* 219:145–161. [CrossRef Medline](#)
- Cassel JC, Pereira de Vasconcelos A, Loureiro M, Cholvin T, Dalrymple-Alford JC, Vertes RP (2013) The reuniens and rhomboid nuclei: neuroanatomy, electrophysiological characteristics and behavioral implications. *Prog Neurobiol* 111:34–52. [CrossRef Medline](#)
- Caulo M, Van Hecke J, Toma L, Ferretti A, Tartaro A, Colosimo C, Romani GL, Uncini A (2005) Functional MRI study of diencephalic amnesia in Wernicke-Korsakoff syndrome. *Brain* 128:1584–1594. [CrossRef Medline](#)
- Cermak LS, Uhly B, Reale L (1980) Encoding specificity in the alcoholic Korsakoff patient. *Brain Lang* 11:119–127. [CrossRef Medline](#)
- Chakraborty S, Kolling N, Walton ME, Mitchell AS (2016) Critical role for the mediodorsal thalamus in permitting rapid reward-guided updating in stochastic reward environments. *eLife* 5:e13588. [CrossRef Medline](#)
- Chauveau F, Célérier A, Ognard R, Pierard C, Béracochéa D (2005) Effects of ibotenic acid lesions of the mediodorsal thalamus on memory: relationship with emotional processes in mice. *Behav Brain Res* 156:215–223. [CrossRef Medline](#)
- Cholvin T, Loureiro M, Cassel R, Cosquer B, Geiger K, De Sa Nogueira D, Raingard H, Robelin L, Kelche C, Pereira de Vasconcelos A, Cassel JC (2013) The ventral midline thalamus contributes to strategy shifting in a memory task requiring both prefrontal cortical and hippocampal functions. *J Neurosci* 33:8772–8783. [CrossRef Medline](#)
- Christiansen K, Dillingham CM, Wright NF, Saunders RC, Vann SD, Aggleton JP (2016) Complementary subicular pathways to the anterior thalamic nuclei and mammillary bodies in the rat and macaque monkey brain. *Eur J Neurosci* 43:1044–1061. [CrossRef Medline](#)
- Clark BJ, Rice JP, Akers KG, Candelaria-Cook FT, Taube JS, Hamilton DA (2013) Lesions of the dorsal tegmental nuclei disrupt control of navigation by distal landmarks in cued, directional, and place variants of the Morris Water Task. *Behav Neurosci* 127:566–581. [CrossRef Medline](#)
- Collins DP, Anastasiades PG, Marlin JJ, Carter AG (2018) Reciprocal circuits linking the prefrontal cortex with dorsal and ventral thalamic nuclei. *Neuron* 98:366–379.e4. [CrossRef Medline](#)
- Corbit LH, Muir JL, Balleine BW (2003) Lesions of mediodorsal thalamus and anterior thalamic nuclei produce dissociable effects on instrumental conditioning in rats. *Eur J Neurosci* 18:1286–1294. [CrossRef Medline](#)
- Corcoran KA, Frick BJ, Radulovic J, Kay LM (2016) Analysis of coherent activity between retrosplenial cortex, hippocampus, thalamus, and anterior cingulate cortex during retrieval of recent and remote context fear memory. *Neurobiol Learn Mem* 127:93–101. [CrossRef Medline](#)
- Courtiole E, Wilson DA (2015) The olfactory thalamus: unanswered questions about the role of the mediodorsal thalamic nucleus in olfaction. *Front Neural Circuits* 9:49. [CrossRef Medline](#)
- Courtiole E, Wilson DA (2016) Neural representation of odor-guided behavior in the rat olfactory thalamus. *J Neurosci* 36:5946–5960. [CrossRef Medline](#)
- Crandall SR, Cruikshank SJ, Connors BW (2015) A corticothalamic switch: controlling the thalamus with dynamic synapses. *Neuron* 86:768–782. [CrossRef Medline](#)
- Crick F (1984) Function of the thalamic reticular complex: the searchlight hypothesis. *Proc Natl Acad Sci U S A* 81:4586–4590. [CrossRef Medline](#)
- Cross L, Brown MW, Aggleton JP, Warburton EC (2012) The medial dorsal thalamic nucleus and the medial prefrontal cortex of the rat function together to support associative recognition and recency but not item recognition. *Learn Mem* 20:41–50. [CrossRef Medline](#)
- Cudeiro J, Sillito AM (2006) Looking back: corticothalamic feedback and early visual processing. *Trends Neurosci* 29:298–306. [CrossRef Medline](#)
- Danet L, Pariente J, Eustache P, Raposo N, Sibon I, Albuicher JF, Bonneville F, Péran P, Barbeau EJ (2017) Medial thalamic stroke and its impact on familiarity and recollection. *eLife* 6:e28141. [CrossRef Medline](#)
- Delay J, Brion S (1969) *Le Syndrome de Korsakoff*. Paris: Masson.
- Delevich K, Tucciarone J, Huang ZJ, Li B (2015) The mediodorsal thalamus drives feedforward inhibition in the anterior cingulate cortex via parvalbumin interneurons. *J Neurosci* 35:5743–5753. [CrossRef Medline](#)
- Dillingham CM, Frizzati A, Nelson AJ, Vann SD (2015a) How do mammillary body inputs contribute to anterior thalamic function? *Neurosci Biobehav Rev* 54:108–119. [CrossRef Medline](#)
- Dillingham CM, Erichsen JT, O'Mara SM, Aggleton JP, Vann SD (2015b) Fornical and nonfornical projections from the rat hippocampal formation to the anterior thalamic nuclei. *Hippocampus* 25:977–992. [CrossRef Medline](#)
- Donoso M, Collins AG, Koechlin E (2014) Human cognition: foundations of human reasoning in the prefrontal cortex. *Science* 344:1481–1486. [CrossRef Medline](#)
- Dudchenko PA (2001) How do animals actually solve the T maze? *Behav Neurosci* 115:850–860. [CrossRef Medline](#)
- Dumont JR, Aggleton JP (2013) Dissociation of recognition and recency memory judgments after anterior thalamic nuclei lesions in rats. *Behav Neurosci* 127:415–431. [CrossRef Medline](#)
- Dupire A, Kant P, Mons N, Marchand AR, Coutureau E, Dalrymple-Alford J, Wolff M (2013) A role for anterior thalamic nuclei in affective cognition: interaction with environmental conditions. *Hippocampus* 23:392–404. [CrossRef Medline](#)
- Edelman GM, Gally JA (2013) Reentry: a key mechanism for integration of brain function. *Front Integr Neurosci* 7:63. [CrossRef Medline](#)
- Eichenbaum H (2017) Prefrontal-hippocampal interactions in episodic memory. *Nat Rev Neurosci* 18:547–558. [CrossRef Medline](#)
- Eichenbaum H, Shedlack KJ, Eckmann KW (1980) Thalamocortical mechanisms in odor-guided behavior: I. Effects of lesions of the mediodorsal thalamic nucleus and frontal cortex on olfactory discrimination in the rat. *Brain Behav Evol* 17:255–275. [CrossRef Medline](#)
- Frizzati A, Milczarek MM, Sengpiel F, Thomas KL, Dillingham CM, Vann SD (2016) Comparable reduction in Zif268 levels and cytochrome oxidase activity in the retrosplenial cortex following mammillothalamic tract lesions. *Neuroscience* 330:39–49. [CrossRef Medline](#)
- Frohardt RJ, Basset JP, Taube JS (2006) Path integration and lesions within the head direction cell circuit: comparison between the roles of the anterodorsal thalamus and dorsal tegmental nucleus. *Behav Neurosci* 120:135–149. [CrossRef Medline](#)
- Funahashi S (2013) Thalamic mediodorsal nucleus and its participation in spatial working memory processes: comparison with the prefrontal cortex. *Front Syst Neurosci* 7:36. [CrossRef Medline](#)
- Galvan A, Smith Y (2011) The primate thalamostriatal systems: anatomical organization, functional roles and possible involvement in Parkinson's disease. *Basal Ganglia* 1:179–189. [CrossRef Medline](#)
- Gamper E (1928) Zur frage der polioencephalitis der chronischen alkoholiker. anatomische befunde beim chronischem korsakow und ihre beziehungen zum klinischen bild. *Deutsche Z Nervenheilkd* 102:122–129. [CrossRef](#)
- Garden DL, Massey PV, Caruana DA, Johnson B, Warburton EC, Aggleton JP, Bashir ZI (2009) Anterior thalamic lesions stop synaptic plasticity in retrosplenial cortex slices: expanding the pathology of diencephalic amnesia. *Brain* 132:1847–1857. [CrossRef Medline](#)

- Goldberg JH, Farries MA, Fee MS (2013) Basal ganglia output to the thalamus: still a paradox. *Trends Neurosci* 36:695–705. [CrossRef Medline](#)
- Groenewegen HJ (1988) Organization of the afferent connections of the mediodorsal thalamic nucleus in the rat, related to the mediodorsal-prefrontal topography. *Neuroscience* 24:379–431. [CrossRef Medline](#)
- Gudden H (1896) Klinische und anatomische Beiträge zur Kenntnis der multiplen Alkoholneuritis nebst Bemerkungen über die Regenerationsvorgänge im peripheren Nervensystem. *Arch Psychiatr* 28:643–741.
- Guo W, Clause AR, Barth-Marion A, Polley DB (2017a) A corticothalamic circuit for dynamic switching between feature detection and discrimination. *Neuron* 95:180–194.e5. [CrossRef Medline](#)
- Guo ZV, Inagaki HK, Daie K, Druckmann S, Gerfen CR, Svoboda K (2017b) Maintenance of persistent activity in a frontal thalamocortical loop. *Nature* 545:181–186. [CrossRef Medline](#)
- Haber SN, Calzavara R (2009) The cortico-basal ganglia integrative network: the role of the thalamus. *Brain Res Bull* 78:69–74. [CrossRef Medline](#)
- Halassa MM, Acsády L (2016) Thalamic inhibition: diverse sources, diverse scales. *Trends Neurosci* 39:680–693. [CrossRef Medline](#)
- Halassa MM, Kastner S (2017) Thalamic functions in distributed cognitive control. *Nat Neurosci* 20:1669–1679. [CrossRef Medline](#)
- Hallock HL, Wang A, Shaw CL, Griffin AL (2013) Transient inactivation of the thalamic nucleus reuniens and rhomboid nucleus produces deficits of a working-memory dependent tactile-visual conditional discrimination task. *Behav Neurosci* 127:860–866. [CrossRef Medline](#)
- Hallock HL, Wang A, Griffin AL (2016) Ventral midline thalamus is critical for hippocampal-prefrontal synchrony and spatial working memory. *J Neurosci* 36:8372–8389. [CrossRef Medline](#)
- Han J, Lee JH, Kim MJ, Jung MW (2013) Neural activity in mediodorsal nucleus of thalamus in rats performing a working memory task. *Front Neural Circuits* 7:128. [CrossRef Medline](#)
- Harding A, Halliday G, Caine D, Kril J (2000) Degeneration of anterior thalamic nuclei differentiates alcoholics with amnesia. *Brain* 123:141–154. [CrossRef Medline](#)
- Hoover WB, Vertes RP (2012) Collateral projections from nucleus reuniens of thalamus to hippocampus and medial prefrontal cortex in the rat: a single and double retrograde fluorescent labeling study. *Brain Struct Funct* 217:191–209. [CrossRef Medline](#)
- Hunkin NM, Parkin AJ, Longmore BE (1994) Aetiological variation in the amnesic syndrome: comparisons using the list discrimination task. *Neuropsychologia* 32:819–825. [CrossRef Medline](#)
- Hunt PR, Aggleton JP (1991) Medial dorsal thalamic lesions and working memory in the rat. *Behav Neural Biol* 55:227–246. [CrossRef Medline](#)
- Hunt PR, Aggleton JP (1998a) Neurotoxic lesions of the dorsomedial thalamus impair the acquisition but not the performance of delayed matching to place by rats: a deficit in shifting response rules. *J Neurosci* 18:10045–10052. [CrossRef Medline](#)
- Hunt PR, Aggleton JP (1998b) An examination of the spatial working memory deficit following neurotoxic medial dorsal thalamic lesions in rats. *Behav Brain Res* 97:129–141. [CrossRef Medline](#)
- Ito HT, Zhang SJ, Witter MP, Moser EI, Moser MB (2015) A prefrontal-thalamo-hippocampal circuit for goal-directed spatial navigation. *Nature* 522:50–55. [CrossRef Medline](#)
- Izquierdo A, Murray EA (2010) Functional interaction of medial mediodorsal thalamic nucleus but not nucleus accumbens with amygdala and orbital prefrontal cortex is essential for adaptive response selection after reinforcer devaluation. *J Neurosci* 30:661–669. [CrossRef Medline](#)
- Jagtap P, Diwadkar VA (2016) Effective connectivity of ascending and descending frontothalamic pathways during sustained attention: complex brain network interactions in adolescence. *Hum Brain Mapp* 37:2557–2570. [CrossRef Medline](#)
- Jankowski MM, Ronnqvist KC, Tsanov M, Vann SD, Wright NF, Erichsen JT, Aggleton JP, O'Mara SM (2013) The anterior thalamus provides a subcortical circuit supporting memory and spatial navigation. *Front Syst Neurosci* 7:45. [CrossRef Medline](#)
- Jankowski MM, Passecker J, Islam MN, Vann S, Erichsen JT, Aggleton JP, O'Mara SM (2015) Evidence for spatially-responsive neurons in the rostral thalamus. *Front Behav Neurosci* 9:256. [CrossRef Medline](#)
- Jenkins TA, Dias R, Amin E, Brown MW, Aggleton JP (2002) Fos imaging reveals that lesions of the anterior thalamic nuclei produce widespread limbic hypoactivity in rats. *J Neurosci* 22:5230–5238. [CrossRef Medline](#)
- Jones EG (1998) Viewpoint: the core and matrix of thalamic organization. *Neuroscience* 85:331–345. [CrossRef Medline](#)
- Kapur N (1994) The mammillary bodies revisited: their role in human memory functioning. In: *Neuropsychological explorations of memory and cognition: essays in honor of Nelson Butters. Critical issues in neuropsychology* (Cermak LS, ed), pp 159–189. New York: Plenum.
- Karlsen AS, Korbo S, Uylings HB, Pakkenberg B (2014) A stereological study of the mediodorsal thalamic nucleus in down syndrome. *Neuroscience* 279:253–259. [CrossRef Medline](#)
- Kato S, Kuramochi M, Kobayashi K, Fukabori R, Okada K, Uchigashima M, Watanabe M, Tsutsui Y, Kobayashi K (2011) Selective neural pathway targeting reveals key roles of thalamostriatal projection in the control of visual discrimination. *J Neurosci* 31:17169–17179. [CrossRef Medline](#)
- Kato S, Fukabori R, Nishizawa K, Okada K, Yoshioka N, Sugawara M, Maejima Y, Shimomura K, Okamoto M, Eifuku S, Kobayashi K (2018) Action selection and flexible switching controlled by the intralaminar thalamic neurons. *Cell Rep* 22:2370–2382. [CrossRef Medline](#)
- Kocsis B, Di Prisco GV, Vertes RP (2001) Theta synchronization in the limbic system: the role of Gudden's tegmental nuclei. *Eur J Neurosci* 13:381–388. [CrossRef Medline](#)
- Kopelman MD, Thomson AD, Guerrini I, Marshall EJ (2009) The Korsakoff syndrome: clinical aspects, psychology and treatment. *Alcohol Alcohol* 44:148–154. [CrossRef Medline](#)
- Kuramoto E, Iwai H, Yamanaka A, Ohno S, Seki H, Tanaka YR, Furuta T, Hioki H, Goto T (2017a) Dorsal and ventral parts of thalamic nucleus submedius project to different areas of rat orbitofrontal cortex: a single neuron-tracing study using virus vectors. *J Comp Neurol* 525:3821–3839. [CrossRef Medline](#)
- Kuramoto E, Pan S, Furuta T, Tanaka YR, Iwai H, Yamanaka A, Ohno S, Kaneko T, Goto T, Hioki H (2017b) Individual mediodorsal thalamic neurons project to multiple areas of the rat prefrontal cortex: a single neuron-tracing study using virus vectors. *J Comp Neurol* 525:166–185. [CrossRef Medline](#)
- Laurens J, Kim B, Dickman JD, Angelaki DE (2016) Gravity orientation tuning in macaque anterior thalamus. *Nat Neurosci* 19:1566–1568. [CrossRef Medline](#)
- Law LM, Smith DM (2012) The anterior thalamus is critical for overcoming interference in a context-dependent odor discrimination task. *Behav Neurosci* 126:710–719. [CrossRef Medline](#)
- Leung BK, Balleine BW (2015) Ventral pallidal projections to mediodorsal thalamus and ventral tegmental area play distinct roles in outcome-specific pavlovian-instrumental transfer. *J Neurosci* 35:4953–4964. [CrossRef Medline](#)
- Li L, Ebner FF (2016) Cortex dynamically modulates responses of thalamic relay neurons through prolonged circuit-level disinhibition in rat thalamus in vivo. *J Neurophysiol* 116:2368–2382. [CrossRef Medline](#)
- Liebermann D, Ploner CJ, Kraft A, Kopp UA, Ostendorf F (2013) A dysexecutive syndrome of the medial thalamus. *Cortex* 49:40–49. [CrossRef Medline](#)
- Linley SB, Gallo MM, Vertes RP (2016) Lesions of the ventral midline thalamus produce deficits in reversal learning and attention on an odor texture set shifting task. *Brain Res* 1649:110–122. [CrossRef Medline](#)
- Loukavenko EA, Ottley MC, Moran JP, Wolff M, Dalrymple-Alford JC (2007) Towards therapy to relieve memory impairment after anterior thalamic lesions: improved spatial working memory after immediate and delayed postoperative enrichment. *Eur J Neurosci* 26:3267–3276. [CrossRef Medline](#)
- Loukavenko EA, Wolff M, Poirier GL, Dalrymple-Alford JC (2016) Impaired spatial working memory after anterior thalamic lesions: recovery with cerebrolysin and enrichment. *Brain Struct Funct* 221:1955–1970. [CrossRef Medline](#)
- Loureiro M, Cholvin T, Lopez J, Merienne N, Latreche A, Cosquer B, Geiger K, Kelche C, Cassel JC, Pereira de Vasconcelos A (2012) The ventral midline thalamus (reuniens and rhomboid nuclei) contributes to the persistence of spatial memory in rats. *J Neurosci* 32:9947–9959. [CrossRef Medline](#)
- Man K, Kaplan J, Damasio H, Damasio A (2013) Neural convergence and divergence in the mammalian cerebral cortex: from experimental neuroanatomy to functional neuroimaging. *J Comp Neurol* 521:4097–4111. [CrossRef Medline](#)
- Marchand A, Faugère A, Coutureau E, Wolff M (2014) A role for anterior thalamic nuclei in contextual fear memory. *Brain Struct Funct* 219:1575–1586. [CrossRef Medline](#)

- Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H (2013) Cortical high-density counterstream architectures. *Science* 342:1238406. [CrossRef Medline](#)
- Marton T, Seifkar H, Luongo FJ, Lee AT, Sohal VS (2018) Roles of prefrontal cortex and mediodorsal thalamus in task engagement and behavioral flexibility. *J Neurosci*. Advance online publication. Retrieved Feb. 7, 2018. doi: 10.1523/JNEUROSCI.1728–17.2018. [CrossRef Medline](#)
- Mathiasen ML, Dillingham CM, Kinnavane L, Powell AL, Aggleton JP (2017) Asymmetric cross-hemispheric connections link the rat anterior thalamic nuclei with the cortex and hippocampal formation. *Neuroscience* 349:128–143. [CrossRef Medline](#)
- McCormick DA, von Krosigk M (1992) Corticothalamic activation modulates thalamic firing through glutamate “metabotropic” receptors. *Proc Natl Acad Sci U S A* 89:2774–2778. [CrossRef Medline](#)
- McKenna JT, Vertes RP (2004) Afferent projections to nucleus reuniens of the thalamus. *J Comp Neurol* 480:115–142. [CrossRef Medline](#)
- Mease RA, Krieger P, Groh A (2014) Cortical control of adaptation and sensory relay mode in the thalamus. *Proc Natl Acad Sci U S A* 111:6798–6803. [CrossRef Medline](#)
- Mendez-Lopez M, Arias JL, Bontempi B, Wolff M (2013) Reduced cytochrome oxidase activity in the retrosplenial cortex after lesions to the anterior thalamic nuclei. *Behav Brain Res* 250C:264–273. [CrossRef Medline](#)
- Miller RL, Francoeur MJ, Gibson BM, Mair RG (2017) Mediodorsal thalamic neurons mirror the Activity of Medial Prefrontal Neurons Responding to Movement and Reinforcement during a Dynamic DNMTTP Task. *eNeuro* 4:ENEURO.0196–17.2017. [CrossRef Medline](#)
- Mitchell AS (2015) The mediodorsal thalamus as a higher order thalamic relay nucleus important for learning and decision-making. *Neurosci Biobehav Rev* 54:76–88. [CrossRef Medline](#)
- Mitchell AS, Chakraborty S (2013) What does the mediodorsal thalamus do? *Front Syst Neurosci* 7:37. [CrossRef Medline](#)
- Mitchell AS, Dalrymple-Alford JC (2005) Dissociable memory effects after medial thalamus lesions in the rat. *Eur J Neurosci* 22:973–985. [CrossRef Medline](#)
- Mitchell AS, Dalrymple-Alford JC (2006) Lateral and anterior thalamic lesions impair independent memory systems. *Learn Mem* 13:388–396. [CrossRef Medline](#)
- Mitchell AS, Dalrymple-Alford JC, Christie MA (2002) Spatial working memory and the brainstem cholinergic innervation to the anterior thalamus. *J Neurosci* 22:1922–1928. [CrossRef Medline](#)
- Mitchell AS, Browning PG, Baxter MG (2007) Neurotoxic lesions of the medial mediodorsal nucleus of the thalamus disrupt reinforcer devaluation effects in rhesus monkeys. *J Neurosci* 27:11289–11295. [CrossRef Medline](#)
- Moreau PH, Tsenkina Y, Lecourtier L, Lopez J, Cosquer B, Wolff M, Dalrymple-Alford J, Cassel JC (2013) Lesions of the anterior thalamic nuclei and intralaminar thalamic nuclei: place and visual discrimination learning in the water maze. *Brain Struct Funct* 218:657–667. [CrossRef Medline](#)
- Murphy MJ, Deutch AY (2018) Organization of afferents to the orbitofrontal cortex in the rat. *J Comp Neurol* 526:1498–1526. [CrossRef Medline](#)
- Murray Sherman S (2018) My prolonged collaboration with Ray Guillery. *Eur J Neurosci*. Advance online publication. Retrieved Feb. 7, 2018. doi: 10.1111/ejn.13903. [CrossRef Medline](#)
- Nakajima M, Halassa MM (2017) Thalamic control of functional cortical connectivity. *Curr Opin Neurobiol* 44:127–131. [CrossRef Medline](#)
- Nakayama H, Ibañez-Tallon I, Heintz N (2018) Cell-type-specific contributions of medial prefrontal neurons to flexible behaviors. *J Neurosci* 38:4490–4504. [CrossRef Medline](#)
- Neave N, Sahgal A, Aggleton JP (1993) Lack of effect of dorsomedial thalamic lesions on automated tests of spatial memory in the rat. *Behav Brain Res* 55:39–49. [CrossRef Medline](#)
- Négyessy L, Hámori J, Bentivoglio M (1998) Contralateral cortical projection to the mediodorsal thalamic nucleus: origin and synaptic organization in the rat. *Neuroscience* 84:741–753. [CrossRef Medline](#)
- Nelson AJ, Vann SD (2014) Mammillothalamic tract lesions disrupt tests of visuo-spatial memory. *Behav Neurosci* 128:494–503. [CrossRef Medline](#)
- Nelson AJD, Vann SD (2017) The importance of mammillary body efferents for recency memory: towards a better understanding of diencephalic amnesia. *Brain Struct Funct* 222:2143–2156. [CrossRef Medline](#)
- Ostlund SB, Balleine BW (2008) Differential involvement of the basolateral amygdala and mediodorsal thalamus in instrumental action selection. *J Neurosci* 28:4398–4405. [CrossRef Medline](#)
- Otis JM, Namboodiri VM, Matan AM, Voets ES, Mohorn EP, Kosyk O, McHenry JA, Robinson JE, Resendez SL, Rossi MA, Stuber GD (2017) Prefrontal cortex output circuits guide reward seeking through divergent cue encoding. *Nature* 543:103–107. [CrossRef Medline](#)
- Ouhaz Z, Fleming H, Mitchell AS (2018) Cognitive functions and neurodevelopmental disorders involving the prefrontal cortex and mediodorsal thalamus. *Front Neurosci* 12:33. [CrossRef Medline](#)
- Ozyurt J, Lorenzen A, Gebhardt U, Warmuth-Metz M, Müller HL, Thiel CM (2014) Remote effects of hypothalamic lesions in the prefrontal cortex of craniopharyngoma patients. *Neurobiol Learn Mem* 111:71–80. [CrossRef Medline](#)
- Padberg J, Cerkevich C, Engle J, Rajan AT, Recanzone G, Kaas J, Krubitzer L (2009) Thalamocortical connections of parietal somatosensory cortical fields in macaque monkeys are highly divergent and convergent. *Cereb Cortex* 19:2038–2064. [CrossRef Medline](#)
- Parker A, Gaffan D (1997) The effect of anterior thalamic and cingulate cortex lesions on object-in-place memory in monkeys. *Neuropsychologia* 35:1093–1102. [CrossRef Medline](#)
- Parnaudeau S, O’Neill PK, Bolkan SS, Ward RD, Abbas AI, Roth BL, Balsam PD, Gordon JA, Kellendonk C (2013) Inhibition of mediodorsal thalamus disrupts thalamofrontal connectivity and cognition. *Neuron* 77:1151–1162. [CrossRef Medline](#)
- Parnaudeau S, Taylor K, Bolkan SS, Ward RD, Balsam PD, Kellendonk C (2015) Mediodorsal thalamus hypofunction impairs flexible goal-directed behavior. *Biol Psychiatry* 77:445–453. [CrossRef Medline](#)
- Parnaudeau S, Bolkan SS, Kellendonk C (2018) The mediodorsal thalamus: an essential partner of the prefrontal cortex for cognition. *Biol Psychiatry* 83:648–656. [CrossRef Medline](#)
- Pereira de Vasconcelos A, Cassel JC (2015) The nonspecific thalamus: a place in a wedding bed for making memories last? *Neurosci Biobehav Rev* 54:175–196. [CrossRef Medline](#)
- Pergola G, Danet L, Pitel AL, Carlesimo GA, Segobin S, Pariente J, Suchan B, Mitchell AS, Barbeau EJ (2018) The regulatory role of the human mediodorsal thalamus. *Trends Cogn Sci* 22:1011–1025. [CrossRef Medline](#)
- Perry JC, Pakkenberg B, Vann SD (2018) Striking reduction in neurons and glial cells in anterior thalamic nuclei of older patients with Down’s syndrome. *bioRxiv* 449678. doi: <https://doi.org/10.1101/449678>.
- Petrof I, Sherman SM (2009) Synaptic properties of the mammillary and cortical afferents to the anterodorsal thalamic nucleus in the mouse. *J Neurosci* 29:7815–7819. [CrossRef Medline](#)
- Pinault D (2004) The thalamic reticular nucleus: structure, function and concept. *Brain Res Brain Res Rev* 46:1–31. [CrossRef Medline](#)
- Pinault D (2011) Dysfunctional thalamus-related networks in schizophrenia. *Schizophr Bull* 37:238–243. [CrossRef Medline](#)
- Powell AL, Nelson AJ, Hindley E, Davies M, Aggleton JP, Vann SD (2017) The rat retrosplenial cortex as a link for frontal functions: a lesion analysis. *Behav Brain Res* 335:88–102. [CrossRef Medline](#)
- Prasad JA, Abela AR, Chudasama Y (2017) Midline thalamic reuniens lesions improve executive behaviors. *Neuroscience* 345:77–88. [CrossRef Medline](#)
- Preuss TM, Goldman-Rakic PS (1987) Crossed corticothalamic and thalamocortical connections of macaque prefrontal cortex. *J Comp Neurol* 257:269–281. [CrossRef Medline](#)
- Ramos RT (2014) The concepts of representation and information in explanatory theories of human behavior. *Front Psychol* 5:1034. [CrossRef Medline](#)
- Reed LJ, Lasserson D, Marsden P, Stanhope N, Stevens T, Bello F, Kingsley D, Colchester A, Kopelman MD (2003) FDG-PET findings in the Wernicke-Korsakoff syndrome. *Cortex* 39:1027–1045. [CrossRef Medline](#)
- Rikhye RV, Wimmer RD, Halassa MM (2018) Toward an integrative theory of thalamic function. *Annu Rev Neurosci* 41:163–183. [CrossRef Medline](#)
- Rubio-Garrido P, Pérez-de-Manzo F, Porrero C, Galazo MJ, Clascá F (2009) Thalamic input to distal apical dendrites in neocortical layer 1 is massive and highly convergent. *Cereb Cortex* 19:2380–2395. [CrossRef Medline](#)
- Salay LD, Ishiko N, Huberman AD (2018) A midline thalamic circuit determines reactions to visual threat. *Nature* 557:183–189. [CrossRef Medline](#)
- Satoh K, Fibiger HC (1986) Cholinergic neurons of the laterodorsal tegmental nucleus: efferent and afferent connections. *J Comp Neurol* 253:277–302. [CrossRef Medline](#)
- Saund J, Dautan D, Rostron C, Urcelay GP, Gerdjikov TV (2017) Thalamic

- inputs to dorsomedial striatum are involved in inhibitory control: evidence from the five-choice serial reaction time task in rats. *Psychopharmacology (Berl)* 234:2399–2407. [CrossRef Medline](#)
- Schepers IM, Beck AK, Bräuer S, Schwabe K, Abdallat M, Sandmann P, Dengler R, Rieger JW, Krauss JK (2017) Human centromedian-parafascicular complex signals sensory cues for goal-oriented behavior selection. *Neuroimage* 152:390–399. [CrossRef Medline](#)
- Schmitt LI, Wimmer RD, Nakajima M, Happ M, Mofakham S, Halassa MM (2017) Thalamic amplification of cortical connectivity sustains attentional control. *Nature* 545:219–223. [CrossRef Medline](#)
- Scoville WB, Milner B (2000) Loss of recent memory after bilateral hippocampal lesions: 1957. *J Neuropsychiatry Clin Neurosci* 12:103–113. [CrossRef Medline](#)
- Seifert S, von Cramon DY, Imperati D, Tittgemeyer M, Ullsperger M (2011) Thalamocingulate interactions in performance monitoring. *J Neurosci* 31:3375–3383. [CrossRef Medline](#)
- Shaw C, Aggleton JP (1995) Evidence for the independence of recognition and recency memory in amnesic subjects. *Cortex* 31:57–71. [CrossRef Medline](#)
- Sherman SM (2005) Thalamic relays and cortical functioning. *Prog Brain Res* 149:107–126. [CrossRef Medline](#)
- Sherman SM (2012) Thalamocortical interactions. *Curr Opin Neurobiol* 22:575–579. [CrossRef Medline](#)
- Sherman SM (2016) Thalamus plays a central role in ongoing cortical functioning. *Nat Neurosci* 16:533–541. [CrossRef Medline](#)
- Sherman SM, Guillery RW (1996) Functional organization of thalamocortical relays. *J Neurophysiol* 76:1367–1395. [CrossRef Medline](#)
- Somogyi G, Hajdu F, Tömböl T, Madarasz M (1978) Limbic projections to the cat thalamus: a horseradish peroxidase study. *Acta Anat (Basel)* 102:68–73. [CrossRef Medline](#)
- Stillova K, Jurak P, Chladek J, Chrastina J, Halamek J, Bockova M, Golde-mundova S, Riha I, Rektor I (2015) The role of anterior nuclei of the thalamus: a subcortical gate in memory processing. An intracerebral recording study. *PLoS One* 10:e0140778. [CrossRef Medline](#)
- Stokes KA, Best PJ (1990) Mediodorsal thalamic lesions impair “reference” and “working” memory in rats. *Physiol Behav* 47:471–476. [CrossRef Medline](#)
- Sweeney-Reed CM, Zaehle T, Voges J, Schmitt FC, Buentjen L, Kopitzki K, Esslinger C, Hinrichs H, Heinze HJ, Knight RT, Richardson-Klavehn A (2014) Corticothalamic phase synchrony and cross-frequency coupling predict human memory formation. *eLife* 3:e05352. [CrossRef Medline](#)
- Sziklas V, Petrides M (2000) Selectivity of the spatial learning deficit after lesions of the mammillary region in rats. *Hippocampus* 10:325–328. [CrossRef Medline](#)
- Taube JS (1995) Head direction cells recorded in the anterior thalamic nuclei of freely moving rats. *J Neurosci* 15:70–86. [CrossRef Medline](#)
- Taube JS (2007) The head direction signal: origins and sensory-motor integration. *Annu Rev Neurosci* 30:181–207. [CrossRef Medline](#)
- Timbie C, Barbas H (2015) Pathways for emotions: specializations in the amygdalar, mediodorsal thalamic, and posterior orbitofrontal network. *J Neurosci* 35:11976–11987. [CrossRef Medline](#)
- Tolman EC (1948) Cognitive maps in rats and men. *Psychol Rev* 55:189–208. [CrossRef Medline](#)
- Tononi G, Edelman GM (1998) Consciousness and complexity. *Science* 282:1846–1851. [CrossRef Medline](#)
- Tononi G, Edelman GM, Sporns O (1998) Complexity and coherency: integrating information in the brain. *Trends Cogn Sci* 2:474–484. [CrossRef Medline](#)
- Tsanov M, Vann SD, Erichsen JT, Wright N, Aggleton JP, O’Mara SM (2011a) Differential regulation of synaptic plasticity of the hippocampal and the hypothalamic inputs to the anterior thalamus. *Hippocampus* 21:1–8. [CrossRef Medline](#)
- Tsanov M, Chah E, Vann SD, Reilly RB, Erichsen JT, Aggleton JP, O’Mara SM (2011b) Theta-modulated head direction cells in the rat anterior thalamus. *J Neurosci* 31:9489–9502. [CrossRef Medline](#)
- Uhlhaas PJ, Roux F, Singer W (2013) Thalamocortical synchronization and cognition: implications for schizophrenia? *Neuron* 77:997–999. [CrossRef Medline](#)
- Ullsperger M, Danielmeier C, Jochem G (2014) Neurophysiology of performance monitoring and adaptive behavior. *Physiol Rev* 94:35–79. [CrossRef Medline](#)
- Usrey WM, Sherman SM (2018) Corticofugal circuits: communication lines from the cortex to the rest of the brain. *J Comp Neurol*. Advance online publication. Retrieved Mar. 10, 2018. doi: 10.1002/cne.24423. [CrossRef Medline](#)
- Vann SD (2009) Gudden’s ventral tegmental nucleus is vital for memory: re-evaluating diencephalic inputs for amnesia. *Brain* 132:2372–2384. [CrossRef Medline](#)
- Vann SD (2013) Dismantling the Papez circuit for memory in rats. *eLife* 2:e00736. [CrossRef Medline](#)
- Vann SD, Aggleton JP (2003) Evidence of a spatial encoding deficit in rats with lesions of the mammillary bodies or mammillothalamic tract. *J Neurosci* 23:3506–3514. [CrossRef Medline](#)
- Vann SD, Albasser MM (2009) Hippocampal, retrosplenial, and prefrontal hypoactivity in a model of diencephalic amnesia: evidence towards an interdependent subcortical-cortical memory network. *Hippocampus* 19:1090–1102. [CrossRef Medline](#)
- Vann SD, Nelson AJ (2015) The mammillary bodies and memory: more than a hippocampal relay. *Prog Brain Res* 219:163–185. [CrossRef Medline](#)
- Varela C (2014) Thalamic neuromodulation and its implications for executive networks. *Front Neural Circuits* 8:69. [CrossRef Medline](#)
- Varela C, Kumar S, Yang JY, Wilson MA (2014) Anatomical substrates for direct interactions between hippocampus, medial prefrontal cortex, and the thalamic nucleus reuniens. *Brain Struct Funct* 219:911–929. [CrossRef Medline](#)
- Vertes RP (2015) Major diencephalic inputs to the hippocampus: supra-mammillary nucleus and nucleus reuniens. circuitry and function. *Prog Brain Res* 219:121–144. [CrossRef Medline](#)
- Vertes RP, Hoover WB, Do Valle AC, Sherman A, Rodriguez JJ (2006) Efferent projections of reuniens and rhomboid nuclei of the thalamus in the rat. *J Comp Neurol* 499:768–796. [CrossRef Medline](#)
- Vetere G, Kenney JW, Tran LM, Xia F, Steadman PE, Parkinson J, Josselyn SA, Frankland PW (2017) Chemogenetic Interrogation of a brain-wide fear memory network in mice. *Neuron* 94:363–374.e4. [CrossRef Medline](#)
- Victor M, Adams R, Collins G (1971) *The Wernicke–Korsakoff syndrome*. Philadelphia: Davis.
- Viena TD, Linley SB, Vertes RP (2018) Inactivation of nucleus reuniens impairs spatial working memory and behavioral flexibility in the rat. *Hippocampus* 28:297–311. [CrossRef Medline](#)
- Vogt BA, Gabriel M (1993) *Neurobiology of cingulate cortex and limbic thalamus*. Boston: Birkhäuser.
- Warburton EC, Aggleton JP (1999) Differential deficits in the Morris water maze following cytotoxic lesions of the anterior thalamus and fornix transection. *Behav Brain Res* 98:27–38. [CrossRef Medline](#)
- Warburton EC, Baird AL, Aggleton JP (1997) Assessing the magnitude of the allocentric spatial deficit associated with complete loss of the anterior thalamic nuclei in rats. *Behav Brain Res* 87:223–232. [CrossRef Medline](#)
- Warburton EC, Baird AL, Morgan A, Muir JL, Aggleton JP (2000) Disconnecting hippocampal projections to the anterior thalamus produces deficits on tests of spatial memory in rats. *Eur J Neurosci* 12:1714–1726. [CrossRef Medline](#)
- Warrington EK, Weiskrantz L (1982) Amnesia: a disconnection syndrome? *Neuropsychologia* 20:233–248. [CrossRef Medline](#)
- Watanabe Y, Funahashi S (2012) Thalamic mediodorsal nucleus and working memory. *Neurosci Biobehav Rev* 36:134–142. [CrossRef Medline](#)
- Wicker E, Turchi J, Malkova L, Forcelli PA (2018) Mediodorsal thalamus is required for discrete phases of goal-directed behavior in macaques. *eLife* 7:e37325. [CrossRef Medline](#)
- Wilson CR, Gaffan D, Browning PG, Baxter MG (2010) Functional localization within the prefrontal cortex: missing the forest for the trees? *Trends Neurosci* 33:533–540. [CrossRef Medline](#)
- Wilson RC, Takahashi YK, Schoenbaum G, Niv Y (2014) Orbitofrontal cortex as a cognitive map of task space. *Neuron* 81:267–279. [CrossRef Medline](#)
- Wilton LA, Baird AL, Muir JL, Honey RC, Aggleton JP (2001) Loss of the thalamic nuclei for “head direction” impairs performance on spatial memory tasks in rats. *Behav Neurosci* 115:861–869. [CrossRef Medline](#)
- Wimmer RD, Schmitt LI, Davidson TJ, Nakajima M, Deisseroth K, Halassa MM (2015) Thalamic control of sensory selection in divided attention. *Nature* 526:705–709. [CrossRef Medline](#)
- Wolff M, Gibb SJ, Dalrymple-Alford JC (2006) Beyond spatial memory: the anterior thalamus and memory for the temporal order of a sequence of odor cues. *J Neurosci* 26:2907–2913. [CrossRef Medline](#)
- Wolff M, Loukavenko EA, Will BE, Dalrymple-Alford JC (2008a) The ex-

- tended hippocampal-diencephalic memory system: enriched housing promotes recovery of the flexible use of spatial representations after anterior thalamic lesions. *Hippocampus* 18:996–1007. [CrossRef Medline](#)
- Wolff M, Gibb SJ, Cassel JC, Dalrymple-Alford JC (2008b) Anterior but not intralaminar thalamic nuclei support allocentric spatial memory. *Neurobiol Learn Mem* 90:71–80. [CrossRef Medline](#)
- Wolff M, Alcaraz F, Marchand AR, Coutureau E (2015a) Functional heterogeneity of the limbic thalamus: from hippocampal to cortical functions. *Neurosci Biobehav Rev* 54:120–130. [CrossRef Medline](#)
- Wolff M, Faugère A, Desfosses É, Coutureau É, Marchand AR (2015b) Mediodorsal but not anterior thalamic nuclei lesions impair acquisition of a conditional discrimination task. *Neurobiol Learn Mem* 125:80–84. [CrossRef Medline](#)
- Wright NF, Vann SD, Aggleton JP, Nelson AJ (2015) A critical role for the anterior thalamus in directing attention to task-relevant stimuli. *J Neurosci* 35:5480–5488. [CrossRef Medline](#)
- Xu W, Südhof TC (2013) A neural circuit for memory specificity and generalization. *Science* 339:1290–1295. [CrossRef Medline](#)
- Yamanaka K, Hori Y, Minamimoto T, Yamada H, Matsumoto N, Enomoto K, Aosaki T, Graybiel AM, Kimura M (2018) Roles of centromedian parafascicular nuclei of thalamus and cholinergic interneurons in the dorsal striatum in associative learning of environmental events. *J Neural Transm (Vienna)* 125:501–513. [CrossRef Medline](#)
- Yoshida A, Dostrovsky JO, Chiang CY (1992) The afferent and efferent connections of the nucleus submedius in the rat. *J Comp Neurol* 324:115–133. [CrossRef Medline](#)
- Zakowski W, Zawistowski P, Braszka L, Jurkowlanec E (2017) The effect of pharmacological inactivation of the mammillary body and anterior thalamic nuclei on hippocampal theta rhythm in urethane-anesthetized rats. *Neuroscience* 362:196–205. [CrossRef Medline](#)
- Zhou T, Zhu H, Fan Z, Wang F, Chen Y, Liang H, Yang Z, Zhang L, Lin L, Zhan Y, Wang Z, Hu H (2017) History of winning remodels thalamo-PFC circuit to reinforce social dominance. *Science* 357:162–168. [CrossRef Medline](#)
- Zikopoulos B, Barbas H (2007) Circuits for multisensory integration and attentional modulation through the prefrontal cortex and the thalamic reticular nucleus in primates. *Rev Neurosci* 18:417–438. [CrossRef Medline](#)