

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/117412/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Yang, Qiu E., Tansawai, Uttapoln, Andrey, Diego O., Wang, Shaolin, Wang, Yang, Sands, Kirsty, Kiddee, Anong, Assawatheptawee, Kanit, Bunchu, Nophawan, Hassan, Brekhna, Walsh, Timothy Rutland and Niumsup, Pannika R. 2019. Environmental dissemination of mcr-1 positive Enterobacteriaceae by *Chrysomya* spp. (common blowfly): An increasing public health risk. *Environment International* 122 , pp. 281-290. 10.1016/j.envint.2018.11.021 file

Publishers page: <http://dx.doi.org/10.1016/j.envint.2018.11.021>  
<<http://dx.doi.org/10.1016/j.envint.2018.11.021>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Environmental dissemination of *mcr-1* positive Enterobacteriaceae by *Chrysomya* spp.**  
2 **(common blowfly): an increasing public health risk**

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Qiu E Yang<sup>1</sup>, Uttapoln Tansawai<sup>2</sup>, Diego O. Andrey<sup>1,3</sup>, Shaolin Wang<sup>4</sup>, Yang Wang<sup>4</sup>, Kirsty Sands<sup>1</sup>, Anong Kiddee<sup>2</sup>, Kanit Assawatheptawee<sup>2</sup>, Nophawan Bunchu<sup>2</sup>, Brekhna Hassan<sup>1</sup>, Timothy Rutland Walsh<sup>1</sup>, Pannika R. Niumsup<sup>2</sup>

**Address information:**

1. Department of Medical Microbiology and Infectious Disease, Division of Infection and Immunity, Cardiff University, Cardiff CF14 4XN, UK
2. Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000, Thailand.
3. Service of Infectious Diseases, Geneva University Hospitals and Faculty of Medicine, 1211 Geneva, Switzerland
4. Beijing Advanced Innovation Centre for Food Nutrition and Human Health, College of Veterinary Medicine, China Agricultural University, Beijing, 100193, China

\*To whom correspondence should be addressed: Qiu-E Yang ([YangQe@cardiff.ac.uk](mailto:YangQe@cardiff.ac.uk)) or Timothy R. Walsh ([WalshTR@cardiff.ac.uk](mailto:WalshTR@cardiff.ac.uk))

## 26 **Abstract**

27 Until recently, the role of insects, and particularly flies, in disseminating antimicrobial  
28 resistance (AMR) has been poorly studied. In this study, we screened blowflies (*Chrysomya*  
29 spp.) from different areas near the city of Phitsanulok, Northern Thailand, for the presence of  
30 AMR genes and in particular, *mcr-1*, using whole genome sequencing (WGS). In total, 48 *mcr-*  
31 *1*-positive isolates were recovered, consisting of 17 *mcr-1*-positive *Klebsiella pneumoniae*  
32 (MCRPKP) and 31 *mcr-1*-positive *Escherichia coli* (MCRPEC) strains. The 17 MCRPKP were  
33 shown to be clonal (ST43) with few single poly nucleomorphs (SNPs) by WGS analysis. In *in-*  
34 *vitro* models, the MCRPKP were shown to be highly virulent. In contrast, 31 recovered  
35 MCRPEC isolates are varied, belonging to 12 different sequence types shared with those  
36 causing human infections. The majority of *mcr-1* gene are located on IncX4 plasmids (29/48,  
37 60.42%), sharing an identical plasmid backbone. These findings highlight the contribution of  
38 flies to the AMR contagion picture in low- and middle-income countries and the challenges of  
39 tackling global AMR.

40

## 41 **Highlights**

- 42 • WGS-based analysis of *mcr-1*-carrying isolates from blowflies (*Chrysomya* spp.)  
43 provides evidence that flies serve as an active vector for the environmental spread of  
44 *mcr-1*-mediated colistin resistance pathogens.
- 45 • Hyper-virulent *mcr-1*-carrying *Klebsiella pneumoniae* isolates were identified from  
46 blowflies, which post an acute public health risk.
- 47 • Diversity of antibiotic resistance genes including *qnrS1*, *fosA*, *bla*<sub>CTX-M-55/14</sub> and *floR*,  
48 was detected in *mcr-1* positive strains recovered from blowflies indicating the  
49 environmental spread of multi-drug resistant pathogens.

50

51 **Keywords:** blow flies; *mcr-1* gene; *Klebsiella pneumoniae*; IncX4 plasmid; multidrug

52 resistance

53 **Running Title:** Environmental dissemination of *mcr-1* gene through blow flies

## 54 **Introduction**

55 There is growing public health concern for environmental dispersal of antibiotic resistance.  
56 Historically, we have focused on resistomes in human and animal gut microbiomes, where  
57 antimicrobial resistance (AMR) and pathogens can be widely spread via animal and human  
58 wastes. However, there is increasing evidence indicating the important role of environmental  
59 factors including wastewater[1,2], wildlife [3] and flies [4], in the dissemination of AMR in  
60 different environments. In particular, flies have been recently recognized as potential reservoirs  
61 exacerbating the spread of antibiotic resistance and pathogens among animals, environments  
62 and human, as they can move freely, often unnoticed, among different public health sectors  
63 including hospitals, human communities and animal farms [4-6]. It has been demonstrated that  
64 houseflies are involved in the mechanical transmission of nosocomial infections with multidrug  
65 resistance bacteria in hospital environments, such as *Shigella* spp.; *Escherichia coli*; *Klebsiella*  
66 spp. and *Enterobacter* spp. [4,5]. According to a study analyzing the antibiotic resistome of  
67 swine manure, the larvae (*Musca domestica*) gut microbiome was significantly affected the  
68 resistant genotypes in manure-borne community [7].

69 Since the first discovery of a mobile colistin resistance mechanism (MCR-1) in November  
70 2015 [8], *mcr*-mediated colistin resistance has been globally reported in Gram-negative  
71 pathogens [9]. Additionally,, the co-resistance of colistin with other last-line antibiotics, has  
72 revealed the emergence of extensively drug resistant (XDR)strains that are virtually untreatable  
73 [10,11]. Recent studies reported housefly and blowfly are also responsible for the spread of  
74 *mcr-1* gene conferring colistin resistance in a Chinese university hospital [12], as well as in pig  
75 farms in Germany [13]. Among them, *mcr-1* was most commonly found in *E. coli* isolates,  
76 although several other Enterobacteriaceae including *K. pneumoniae* have been detected in other  
77 source such as animals [14] and human clinical isolates [15]. *K. pneumoniae* is known to be a  
78 leading cause of hospital-acquired infections, such as pneumonia, post-surgical wound and

79 urinary tract infections [16]. It is especially problematic in hospitals when becoming resistant  
80 to colistin, a last-resort antibiotic, leaves very limited therapeutic options. There is a marked  
81 paucity of our understanding on the transmission of *mcr-1*-positive *Enterobacteriaceae*  
82 (MCRPE), mainly in *E. coli* and *K. pneumoniae*, to negate the threat to human health posed by  
83 MCRPE isolates. With this aim, we investigated the carriage of *mcr-1* positive isolates from  
84 blowflies collected from different areas in the city of Phitsanulok in Thailand, using whole-  
85 genome sequence to look for associations between MCRPE strains carrying *mcr-1*-linked  
86 plasmids recovered from blowflies (*Chrysomya* spp.) and human clinical isolates.

87

88

## 89 **Materials and Methods**

### 90 **Bacterial isolates from blow flies**

91 A total of 300 blow flies were trapped at three different locations in Northern Thailand: a local  
92 market in an urban community, a rural area and a suburb of the city Phitsanulok. These  
93 locations are approximately 10 kilometers apart. Blow flies were collected by the use of a  
94 sterile sweeping net. Individual fly was kept in a sterile plastic tube and sacrificed by placing  
95 on ice for 30 min. They were identified to species level by using the taxonomic keys as  
96 described by Kurahashi and Bunchu [17]. Only *Chrysomya megacephala* flies [18], the most  
97 abundant blow flies in Thailand were selected for further analysis. The flies were individually  
98 pulverized in enriched peptone water for 30 min and then aliquots of the resultant suspensions  
99 (100  $\mu$ l) were plated on Eosin-Methylene-Blue (EMB)-agar plates supplemented with 2 mg/l  
100 colistin and incubated at 37°C overnight. One to three representative colonies with different  
101 colors from each plates were purified and subsequently screened for *mcr-1* gene by PCR. The  
102 *mcr-1*-positive bacteria were sub-cultured in liquid nutrient broth for 18 h before DNA  
103 extraction for species identification and whole-genome sequence. Minimum inhibitory

104 concentrations (MICs) of colistin for 48 *mcr-1*-bearing isolates was performed by using broth  
105 microdilution, in accordance with the guideline of the European Committee on Antimicrobial  
106 Susceptibility Testing (EUCAST), reference strain *E. coli* ATCC25922 served as a quality  
107 control.

108

### 109 **Conjugation experiments**

110 To investigate the transferability of *mcr-1*-carrying plasmids, we performed conjugation  
111 assays with sodium-azide resistance *E. coli* J53 as the recipient strain. Briefly, overnight  
112 cultures of 30 randomly selected *mcr-1*-producing donors (strains with transfer frequency  
113 showed in Table 1) and the recipient *E. coli* J53 strain were 1:2 mixed and incubated in 37 °C  
114 for 16-20 h. After incubation, we subsequently ten-fold serial diluted the mixed culture in  
115 sterile saline and aliquoted 100 µl of diluted culture onto selective agar plates containing 2  
116 mg/l colistin and 150 mg/l sodium azide. The *mcr-1*-positive transconjugants were confirmed  
117 by PCR and transfer frequency was calculated by the number of transconjugants per recipient.  
118 Plasmid analysis were done by whole genome sequence as described below.

119

### 120 **Whole-genome sequencing and bioinformatics analysis**

121 Total gDNA was extracted from an overnight culture (2 ml) on a QIAcube automated system  
122 (Qiagen, Germany) with QIAamp DNA Microbiome kit (Qiagen, Germany), followed by  
123 gDNA quantity measurement by fluorometric methods using a Qubit (ThermoFisher  
124 Scientific). Genomic DNA libraries are constructed using the NexteraXT kit (Illumina),  
125 according to manufacturer's instruction. Paired end sequencing was performed using the  
126 Illumina MiSeq platform (MiSeq Reagent V3 Kit; 2 × 300 cycles). Raw sequence reads were  
127 trimmed using Trim Galore and the genomes were *de novo*-assembled into contigs using  
128 SPAdes (3.9.0) with pre-defined kmers set.

129

130 **Bioinformatics analysis:** The CGE platform (<http://www.genomicepidemiology.org/>) were  
131 used for analysis of multilocus sequence typing (MLST-1.8), acquired resistance genes  
132 (ResFinder 3.1, all antibiotic resistance databases were selected with a cut-off value of 95%  
133 identity and 80% minimum coverage) and incompatibility group of plasmids (PlasmidFinder-  
134 1.3 version, using Enterobacteriaceae database with parameters of minimum 95% identity and  
135 85% query coverage). All contigs were searched for *mcr-I* using standalone BLAST analysis,  
136 the putative coding sequences containing *mcr-I* gene were obtained using ORF finder  
137 programs (Geneious 10.0.7). Draft genome sequences were aligned and then applied for  
138 phylogenetic analysis using Parsnp in the Harvest package, and phylogenetic trees was  
139 visualized by iTOL (<https://itol.embl.de/>). MCRPKP strain p38 recovered from healthy human  
140 feces in Thailand was served as a reference strain in the SNPs analysis of 17 MCRPKP strains  
141 in this study.

142 Primer walking was performed to fill the gap in *mcr-I*-carrying contigs from strain PN105  
143 with primers (PN\_IncX4\_forward: CGACCTTTAAGTCGTATTTGCAAGT;  
144 PN\_IncX4\_reverse: ATTGCGCCCGTAGTTCGCTA, Tm 60°C) and the complete plasmid  
145 sequence were constructed by *de novo* assembly using Geneious (10.0.7). The circular  
146 comparisons among *mcr-I*-related IncX4 plasmid backgrounds were performed using BLAST Ring  
147 Image Generator (BRIG v0.9555). Briefly, *mcr-I*-containing contigs were extracted from genomic data,  
148 and a fully sequenced *mcr-I*-linked IncX4 plasmid was served as a central reference sequence. In this  
149 study, plasmid IncX4 from strain PN105 and pMR0617mcr (GenBank No. CP024041), act as reference  
150 sequences. The similarity of between the central reference sequence and other *mcr-I*-positive contigs  
151 from studied MCRPE strains, shows as concentric rings with representative colors.

152

153 **Virulence factors of *K. pneumoniae* isolates**



154 Based on whole-genome sequence data, we have developed a database with publicly  
155 available genomes [NTUH-K2044 (Genbank accession No. AP006725.1, AB117611.1),  
156 pK2044 (Genbank accession No. NC\_006625.1), pLVPK (Genbank accession No.  
157 NC\_005249.1), allantoin metabolism (Genbank accession No. AB115590.1), SB3193  
158 (Genbank accession No. LK022716.1), *uge* CDS (Genbank accession No. AY294624.1),  
159 Kp52.145 (Genbank accession No. FO834906.1), SB4536\_2858 (Genbank accession No.  
160 HG518478.1), pO26-Vir (Genbank accession No. NC\_012487.1), IHE3034 (Genbank  
161 accession No. AM229678.1), *kvg* operon (Genbank accession No. AJ250891.2)] to determine  
162 the key virulence factors in *K. pneumoniae* strains. Our database includes a set of virulence  
163 genes: capsular biosynthesis genes (*wzy/magA*, *K2A*) [20,21]; mucoid factor regulator (*rmpA*,  
164 *rmpA2*) [22]; allantoin metabolism operons (*allABCDRS*, *gcl* and *glxRK*) [23]; an iron-uptake  
165 system (*kfuABC*) [24]; two-component operon (*kvgAS*) [25]; gene clusters for siderophores  
166 dependent iron acquisition (aerobactin *iucBCD-iutA*, yersiniabactin *ybtAEPQSTUX-irp1-irp2-*  
167 *fyuA*, colibactin *clbBCDEFGHHIJKLMNOPQR*, salmochellin *iroBCDN*, enterobactin  
168 *entABCDEF*) [26]; gene clusters of type I and type III fimbriae (*fimABCDEF* and  
169 *mrkABCDFHIJ*, respectively) [27,28]; outer membrane lipoprotein (*ycfM*); serum resistance  
170 factor (*traT*) [29]; hemolysin transport protein (*hlyABCD*) [30]; urease operon (*ureABCDEFG*)  
171 [31] and type IV secretory system gene cluster (*virB1* to *B11*) [32]. Annotation of genes with  
172 75% identity to reference sequences was performed by Geneious (10.2; Biomatters  
173 Ltd.). Capsular (KL) loci were evaluated using Kaptive platform  
174 (<http://kaptive.holtlab.net/jobs>) [33].

175 A *Galleria mellonella* model has been used for virulence test for MCRPKP isolates. Log-  
176 phase cultures of *K. pneumoniae* strains were washed with sterile saline twice, followed by  
177 standardization of bacterial concentrations to approximately  $1 \times 10^9$  -  $1 \times 10^5$  CFU/mL and 10  
178  $\mu$ l were injected into each *G. mellonella* larvae at a final inoculum ranging from  $1 \times 10^6$  -  $1 \times 10^4$

179 cfu/ml, as described previously [34]. The ST23 *Klebsiella pneumoniae* A58300 strain,  
180 harboring the K1 capsule serotype (hyperviscosity phenotype widely associated with  
181 hypervirulent strains) served as virulent control strain in the *G. mellonella* model [35].

182

183

## 184 **Results**

### 185 **Details of isolates**

186 Overall, we recovered 48 MCRPE isolates from 300 collected blowflies (16.0% flies  
187 positivity for *mcr-1*), consisting of 31 *mcr-1*-positive *E. coli* (MCRPEC) and 17 *mcr-1*-positive  
188 *K. pneumoniae* (MCRPKP). Bacterial species were determined by whole genome sequencing  
189 data. Among them, 4 MCRPE strains were recovered from local market in urban community,  
190 16 from rural area and 28 from suburb area. MICs of colistin for all MCR-1-producing isolates  
191 are ranging from 4-16 mg/l (**Table 1**).

192

### 193 **Whole genome sequencing (WGS) analysis**

194 A total of 48 MCRPE isolates were sequenced using Illumina Miseq platform. The distinct  
195 MCRPEC isolates belonged to 12 STs (**Table 1**): ST10(n=7), ST648(n=5), ST549(n=4),  
196 ST58(n=3), ST181(n=3), ST218(n=2), ST201(n=1), ST162(n=1), ST457(n=1), ST1244(n=1),  
197 ST2345(n=1), ST2705(n=1) and ST5487(n=1). Most interestingly, all 17 MCRPKP isolates  
198 belonged to ST43, thus we further determined the clonal relationship of 17 ST43 *K.*  
199 *pneumoniae* isolates by SNPs analysis using Parsnps software. Phylogenetic tree analysis for  
200 17 strains based on their raw sequencing reads showed that their core genome differed only by  
201 a few SNPs (the numbers of differences in SNPs are up to 15, Table S1), suggesting the clonal  
202 dissemination of ST43 *K. pneumoniae* isolates in *Chrysomya* spp. from Thailand.

203 Analysis of genomic accessory modules including acquired resistance genes, virulence  
204 factors and metal resistance genes, showed significant variations in resistance gene content.  
205 Apart from *mcr-1* gene, multiple antibiotic resistance genes were identified in most of the  
206 isolates, with the average number 9.35 and 7 in MCRPEC strains and MCRPKP strains,  
207 respectively. In the 31 MCRPEC collection, 25 difference resistance genes were identified by  
208 ResFinder 3.1 (<https://cge.cbs.dtu.dk/services/ResFinder/>, updated on 2018-09-10), conferring  
209 resistance to nearly all currently available antibiotics, such as  $\beta$ -lactams, aminoglycoside,  
210 chloramphenicol, fluoroquinolones and sulfonamide (**Fig.1**). The most prevalence resistance  
211 genes are *mdfA* resistant to macrolide (n=28, 90.32%), followed by gene *aadA2* conferring  
212 streptomycin resistance (n=26, 83.87%). Besides *aadA2* gene, several genes resistant to  
213 aminoglycoside were detected: *aadA1* (n=16), *aadA17* (n=2), *aac(3')-IId* (n=8), *aph(3')-Ib*  
214 (n=11), *aph(6')-Id* (n=11) and *aph(3')-Ia* (n=1). Furthermore, the plasmid-mediated  
215 fluoroquinolone resistance gene, *qnrS1*, was found in 18 MCRPEC isolates, and three  $\beta$ -  
216 lactamase-producing genes, *bla<sub>TEM-1b</sub>*, *bla<sub>CTX-M-55</sub>* and *bla<sub>CTX-M-14</sub>* are detected in 19 (61.29%),  
217 6 (19.35%) and 1 (3.23%) isolates, respectively. Additionally, resistant genes responsible to  
218 other groups of antibiotics were observed in Fig. 1, including tetracycline resistance (*tetA*, *tetB*  
219 and *tetM*), phenicol resistance (*cmlA*, *floR* and *catA*), and sulphonamine resistance (*sul2* and  
220 *sul3*).

221 In contrast, **identical resistant genotypes among MCRPKP strains were observed, and** seven  
222 acquired antibiotic resistance genes were detected in 17 MCRPKP strains, namely, *mcr-1*,  
223 *mcr-8*, *qnrS1*, *bla<sub>TEM-1b</sub>*, *tetA*, *bla<sub>SHV-40</sub>* encoding SHV  $\beta$ -lactamase and *fosA* mediating  
224 fosmycin resistance.

225

226 ***mcr-1*-associated plasmid types and transferability of *mcr-1* gene**

227 *De novo* bacterial genome assembly was performed and the *mcr-1*-carrying contigs were  
228 analyzed. Replication origin are located in the *mcr-1* contigs, allowing to analyze  
229 incompatibility groups of these plasmids by using PlasmidFinder  
230 (<https://cge.cbs.dtu.dk/services/PlasmidFinder/>). In 35 out of 48 isolates, replicon sequence  
231 type of *mcr-1*-harbouring plasmids could be identified: IncX4 (n=29, 12 *E. coli* and all 17 *K.*  
232 *pneumoniae* isolates), IncHI1A (n=2), IncHI1B (n=3) and IncHI1A-IncHI1B (n=1).  
233 Representative 10 *mcr-1*-bearing IncX4 plasmids obtained from *K. pneumoniae* isolates were  
234 probed for *mcr-1* gene using S1-PFGE. As shown in **Fig.2**, the 10 *mcr-1* genes were all located  
235 on a ~32-kb IncX4 plasmid. PCR was performed to fill the gap in *mcr-1*-carrying contigs, as a  
236 result, complete sequencings of 26 IncX4-*mcr-1*-carrying plasmids were achieved (**Fig. 3**).  
237 Alignment of 26 *mcr-1*-carrying IncX4 plasmids visualized by software BRIG v0.9555 showed  
238 that all *mcr-1*-carrying plasmid share the identical plasmid backbone, including the typical  
239 region encoding ~11kb T4ss conjugation system and a toxin-antitoxin system *hicAB*. More  
240 importantly, in **Fig. 3B**, the backbone of *mcr-1*-linked IncX4 plasmids from blowflies, are  
241 highly similar to those recovered from other sources including companion animals, human  
242 feces and poultry, further suggesting that this type of IncX4 plasmid facilitate the transmission  
243 of *mcr-1* gene. Furthermore, the transferability of *mcr-1*-bearing plasmids were performed by  
244 conjugation with *E.coli* J53 as a recipient. We randomly selected 31 *mcr-1*-positive isolates as  
245 donors, containing four different *mcr-1*-linked Inc-type plasmids: IncX4 plasmids (n=8, three  
246 *mcr-1*- positive *Klebsiella* isolates and five *mcr-1*- positive *E. coli*), IncHI1B plasmids (n=3),  
247 IncHI1A (n=2) and IncHI1A\_HI1B (n=1) (**Table 1**). 12 out of 14 *mcr-1*-bearing plasmids were  
248 successfully transferred to *E. coli* J53, IncX4-*mcr-1* plasmids are able to transferred into the  
249 recipient at a higher frequency (mean  $1.46 \times 10^{-3}$  in *E. coli* and mean  $2.11 \times 10^{-5}$  in *K.*  
250 *pneumoniae*), compared to other *mcr-1*-related IncHI1 plasmid types ( $2.77 \times 10^{-7}$ ), IncHI1A  
251 (mean  $2.27 \times 10^{-7}$ ), IncHI1B ( $1.85 \times 10^{-7}$ ) and IncHI1A-IncHI1B ( $8.7 \times 10^{-8}$ ) (**Table 1**).

252

### 253 **Virulence factors in MCRPKP and virulence loss in *G. mellonella* model**

254 *K. pneumoniae* is recognized as a serious threat to patients due to the emergence of MDR  
255 strains associated with hospital outbreaks and that they demonstrate a number of virulence  
256 factors associated with bacterial pathogenicity and poor patient outcome [36]. There are at  
257 least five groups of pathogenicity factors found in 17 MCRPKP isolates (**Table S1**), these  
258 include gene clusters associated with serum resistance (*traT*); adhesins (type I fimbrial operon  
259 *fimABCDEFGH*, and type III fimbrial operon *mrkBCDEF*); lipopolysaccharide (*wabGHN*);  
260 siderophore systems enterobactin-*entABCDEF*S, and aerobactin-*iucABCD-iutA*. Using  
261 PlasmidFinder (<https://cge.cbs.dtu.dk/services/PlasmidFinder/>), IncFIB-plasmid-related  
262 aerobactin was found in at least 14 isolates, suggesting potential aerobactin-mediated virulence  
263 transferability. Capsular synthesis loci matched type KL61 (*wzi/wzc* 412/61 typing) (Kaptive).  
264 Similarly, the iron acquisition operons *kfuABC*, *iroE*, urease-synthesis operon *ureABCDEFG*  
265 associated with gastric ulceration and urinary stone formation, and gene *ycfM* encoding surface  
266 protein were identified in all MCRPKP isolates. At least 10 different virulence factors were  
267 found in the 17 ST43 MCRPKP strains, which we further analysed by using a *G. mellonella*  
268 model [34]. The effect of different inoculum of 10 randomly selected ST43 MCRPKP strains  
269 was assessed in this model using K1 *rmpA*-positive *K. pneumoniae* A58300, as a hypervirulent  
270 strain reference strain [35]. As shown in **Fig. 4**, after 12-hour post-infection, with an inoculum  
271 of approx.  $1 \times 10^6$  CFU, 100% of mortality was observed with the K1 strain and all 10 ST43  
272 MCRPKP strains. At an inoculum of  $1 \times 10^5$  CFU, the survival rate was 70% with K1 strain  
273 and 0% with 10 ST43 MCRPKP strains. With an inoculum of  $1 \times 10^4$  CFU, 100% survival was  
274 seen with the K1 strain but only 20% with ST43. The consistency between genotypic virulence  
275 factors and the reproducible results of the *G. mellonella* infection model suggest that the ST43  
276 MCRPKP strains recovered from blowflies are highly virulent clones.

277

278 Based on our previous study, the acquisition of *mcr-1*-carrying plasmid leads to virulence  
279 loss in *E. coli* strain [34]. In this study, three *mcr-1*-carrying plasmids were transferred into a  
280 clinical susceptible *K. pneumoniae* strain ff101 and a KPC-positive *K. pneumoniae* strain p35,  
281 followed by infection of *G. mellonella* larvae with an inoculum of  $\sim 1 \times 10^5$  CFU. As shown in  
282 **Fig. 5**, two *K. pneumoniae* strains ff101 and p35 caused more than 80% and 90% of mortality  
283 after 72h infection, respectively. After acquiring of IncX4-*mcr-1* plasmid, the survivals of  
284 larvae have been increased to 40% - 80% with strain ff101, and survivals are more than five  
285 times higher from  $\sim 10\%$  to  $\sim 50\%$  with strain IncX4-*mcr-1*-carrying p35 strain at 72h after  
286 infection, suggesting that IncX4-*mcr-1* plasmid are responsible to reduce bacterial virulence.

287

288

## 289 **Discussion**

290 The *mcr-1* gene was first discovered in *E. coli*, which has become the major host of *mcr-1*  
291 gene, and has subsequently been found in all continents crossing more than 50 countries [8,37].  
292 From 'one health' perspective, environmental factors seem to be closely associated with the  
293 health of human and animals [38], for instance, the heavy use of antibiotic in livestock or  
294 human and their entry into sewage system, is considered as the major cause of resistance  
295 developing in zoonotic bacteria. MCRPE isolates have been mainly recovered from animal  
296 samples [39] and infrequently from human normal flora and clinical samples [40,41], there is  
297 a dearth of evidence on the link between these different populations. Here we present evidence  
298 that blowflies serve an environmental pathway for the transmission of MCR-positive bacteria  
299 including human pathogens. The importance of blowflies in the dissemination of MDR bacteria  
300 is only becoming recognized and represents an additional public health concern. Due to their  
301 habitation and their association with food animals and human, flies present a critical but under-

302 valued link between the environment and human communities. It has been previously identified  
303 that flies can carry the same ESBL-producing *E. coli* clone as found in chicken manure in the  
304 Netherlands [42], and identical antibiotic resistance genes were characterized from both flies  
305 and swine feces [43]. In a modelling study, eight calves were exposed to flies, which were  
306 inoculated with *E. coli* O157:H7, after 24 hours, fecal samples from all calves and drinking  
307 water were positive for *E. coli* O157:H7 [44], suggesting that flies act an effective vector for  
308 the spread of bacteria between animals and the synanthropic environment through feeding and  
309 defecation. In our study, in the region Northern Thailand, 16% (48/300) of studied blowflies  
310 possessed *mcr-I* gene, predominantly located on IncX4 plasmids (29/48, 60.42%) with higher  
311 frequency (up to  $5.93 \times 10^{-3}$ ), when comparing to other *mcr-I*-linked Inc types, such as IncHI1A  
312 (mean  $2.27 \times 10^{-7}$ , Table 1). This high transferability of *mcr-I*-bearing plasmids from insect-  
313 borne bacteria indicates that these bacteria can act as environmental reservoirs of MCR-  
314 positive bacteria, which can potentially become human pathogens. So far, no less than 14  
315 different *mcr-I* bearing plasmids incompatibility types have been identified with  
316 approximately 35.2% of published *mcr-I*-carrying plasmids belonging to IncX4 plasmid,  
317 which has been circulating in human, animal and environmental sectors [45]. More importantly,  
318 the identical nucleotide sequences of 26 *mcr-I*-carrying IncX4 plasmid in our study share an  
319 identical plasmid backbone to that obtained from human and animal samples in Thailand (**Fig.**  
320 **3B**), with typical IncX4-plasmid housekeeping functions and an accessory *-mcr-I-pap2-*  
321 cassette [34], further implying that IncX4-type plasmids serve as a *mcr-I* gene pool in Thailand.  
322 The conjugative function of these *mcr-I*-linked plasmids allow *mcr-I* gene to spread  
323 horizontally in and/or cross the species. Many insert elements have been recognized as a ‘copy-  
324 out-paste-in’ mechanism, which can facilitate the acquisition and mobilization of antibiotic  
325 resistance genes between bacterial pathogens [46]. For example, two copies or one copy of  
326 *ISAp11* flanking in *mcr-I-papA* segment, is actively involved in capture and dissemination of

327 *mcr-1* genes [47-49]. Interestingly, lacking *ISAp11* or other insert element were found in *mcr-*  
328 *1*-associated IncX4 genetic context in this study (Fig. 3), which lead to the hypothesis that  
329 *ISAp11* initially mediates the movement of *mcr-1* genes, and lose one or both copies of *ISAp11*  
330 during subsequent recombination [9,46,48].

331

332 WGS analysis provided comprehensive information for the *mcr-1*-carrying bacteria and their  
333 phylogenetic relationship. Twelve different STs were identified in 31 MCRPEC strains, which  
334 is consistent with other studies that MCRPEC isolates are highly diverse [9,50,51], but ST10-  
335 like *E. coli* seems to represented the higher proportion in MCRPEC isolates (7/31, 22.58%).  
336 *E. coli* ST10 frequently recovered from meat products [52], food-borne animals [53] and  
337 human clinical samples [54], has been strongly associated with human infections and ESBL-  
338 production [55]. Interestingly, in a surveillance study, ESBL-producing *E.coli* ST10 is the most  
339 predominant lineage obtained from a military medical center in America [56]. *E. coli* ST10 is  
340 common among MCRPEC isolates [45], recovered from human [57], animals [58] and  
341 environmental sectors [59]. Apart from *mcr-1* gene, a variety of acquired resistance genes were  
342 detected in all MCRPEC isolates (Fig.1), including plasmid-mediated quinolone resistance  
343 gene (*qnrS1*) and ESBL-dependent *bla*<sub>CTX-M-14</sub> and *bla*<sub>CTX-M-55</sub> genes. Interestingly, a higher  
344 number of acquired resistance genes has been found in seven *E. coli* ST10 isolates (mean 11.71,  
345 ranging from 10 to 14), compared to other STs groups with average 9.67 ranging from 2 to  
346 11(**Fig.1**), further supporting the previously findings that ST10-like *E. coli* strains are linked  
347 to *mcr-1* gene [50].

348

349 Compared with the prevalence of MCRPEC, the reported incidence of MCRPKP is  
350 comparatively rare. In a recent study, *mcr-1*-positive *E.coli*, *Providencia* spp and *Enterobacter*  
351 *cloacae* strains were recovered from blowflies in China, but no *mcr-1*-carrying *K. pneumoniae*



352 strain was identified [12]. Sporadic *mcr-1*-positive isolates of *K. pneumoniae* have been  
353 identified from patients [60-62], animal samples [63] and environmental sector (sewage water)  
354 [64]. However, a recent outbreak of *bla*<sub>KPC</sub> positive MCRPKP in Portugal as further raised the  
355 seriousness of MCRPKP [65]. In this study 17 MCRPKP isolates were recovered from 300  
356 blowflies (5.67%, 17/300) and all belonged to ST43 which has been reported globally in  
357 clinical bacteria associated with abdominal infections [66,67], bacteremia [68] and intensive  
358 care unit infections [69,70]. In addition, ST43 *K. pneumoniae* strains can carry clinically  
359 relevant  $\beta$ -lactamases including NDM-1, CTX-M-15, VIM-5, and OXA-181 [66,70]  
360 (supplementary Table S2). Furthermore, the pairwise analysis of SNPs data (no more than 15  
361 SNPs, Table S2) further suggest ST43 MCRPKP clonality. This scenario is worrying, as  
362 blowflies can act as an efficient and “unseen” environmental vectors of virulent bacteria, and  
363 are associated with outbreaks of enteric pathogens in rural areas in low- and middle income  
364 countries where sanitation and hygiene infrastructure is poor [6,71]. Additionally, the ST43  
365 MCRPKP isolates described in our study also contain at least four major virulence  
366 determinants responsible for disease progression: capsular synthesis loci KL61;  
367 lipopolysaccharide; siderophores enterobactin and (mobilizable) aerobactin, iron acquisition  
368 *kfuABC* that are responsible for binding ferric iron in the host cell; and adherence factors  
369 (fimbria type I and III) that allow bacteria to attach to the host cell surface [24,36,72]. The  
370 virulence potential of these isolates performed in a *G. mellonella* model (**Fig. 4**), suggest that  
371 ST43 MCRPKP are virulent clones circulating with blowflies in Thailand. Thus, MCRPE  
372 strains obtained from blowflies present a global public health problem owing to: i) common  
373 blowflies inhabit the environment and global communities, and act as bacterial environmental  
374 reservoirs via animals/humans and waste; ii) most of MCRPE strains in these study are  
375 multidrug resistant, especially the virulent MCRPKP strains that can be transmitted and cause  
376 infections in humans via contact with blowflies e.g post-surgical wounds, and iii) *mcr-1*

377 detected from blowflies is located on transferable plasmids increasing the possibility of  
378 horizontal transfer of *mcr-I* gene between bacteria as part of the blowflies microbiota, thereby  
379 increasing the environmental gene pool and posing a greater public health risk.

380

381 **Data availability:** Whole genomic sequences of 48 studied MCRPE strains have been  
382 deposited in the NCBI database (BioProject accession No. PRJNA503337 and BioSample  
383 accession No. SAMN10358806 to SAMN10358853). Genomic data of additional 17 MCRPEC  
384 strains showed in Fig.3B were also submitted to the NCBI database (BioProject accession No.  
385 PRJNA504530 and BioSample accession No. SAMN10394864 to SAMN10394880).

386

387

### 388 **Acknowledgement**

389 We thank Ana Cristina Gales (Escola de Medicina, Universidade Federal Paulista, Brazil),  
390 for providing *K. pneumoniae* A58300 strain. This work was supported by MRC grant DETER-  
391 XDR-CHINA (MR/P007295/1). Qiu E Yang is funded by a Chinese Scholarship Council  
392 (CSC). Diego O. Andrey benefits a Geneva University Hospitals (HUG) and Swiss National  
393 Science Foundation (P300PB\_171601) overseas fellowship. This work was partly funded by  
394 Naresuan University (R2562B090). Uttapoln Tansawai (PHD/0054/2555) and Anong Kiddee  
395 (PHD/0181/2557) were supported by the Royal Golden Jubilee-PhD program from Thailand  
396 Research Fund, Rajamangala University of Technology Lanna and Naresuan University.

397

398

399 **All authors have no conflict of interest to declare in the article.**

400

401

402 Reference

- 403 1. Czekalski N, Diez EG, Burgmann H (2014) Wastewater as a point source of antibiotic-  
404 resistance genes in the sediment of a freshwater lake. *Isme Journal* 8: 1381-1390.
- 405 2. Zhu YG, Gillings M, Simonet P, Stekel D, Banwart S, et al. (2017) Microbial mass  
406 movements. *Science* 357: 1099-1100.
- 407 3. Carroll D, Wang J, Fanning S, McMahon BJ (2015) Antimicrobial Resistance in Wildlife:  
408 Implications for Public Health. *Zoonoses Public Health* 62: 534-542.
- 409 4. Onwugamba FC, Fitzgerald JR, Rochon K, Guardabassi L, Alabi A, et al. (2018) The role of  
410 'filth flies' in the spread of antimicrobial resistance. *Travel Med Infect Dis*  
411 10.1016/j.tmaid.2018.02.007.
- 412 5. Graczyk TK, Knight R, Gilman RH, Cranfield MR (2001) The role of non-biting flies in the  
413 epidemiology of human infectious diseases. *Microbes Infect* 3: 231-235.
- 414 6. Echeverria P, Harrison BA, Tirapat C, Mcfarland A (1983) Flies as a Source of Enteric  
415 Pathogens in a Rural Village in Thailand. *Applied and environmental microbiology* 46:  
416 32-36.
- 417 7. Wang H, Sangwan N, Li HY, Su JQ, Oyang WY, et al. (2017) The antibiotic resistome of  
418 swine manure is significantly altered by association with the *Musca domestica* larvae  
419 gut microbiome. *Isme Journal* 11: 100-111.
- 420 8. Liu Y-Y, Wang Y, Walsh TR, Yi L-X, Zhang R, et al. (2015) Emergence of plasmid-mediated  
421 colistin resistance mechanism MCR-1 in animals and human beings in China: a  
422 microbiological and molecular biological study. *The Lancet Infectious Diseases*.
- 423 9. Wang R, van Dorp L, Shaw LP, Bradley P, Wang Q, et al. (2018) The global distribution and  
424 spread of the mobilized colistin resistance gene *mcr-1*. *Nat Commun* 9: 1179.
- 425 10. Dobiasova H, Dolejska M (2016) Prevalence and diversity of IncX plasmids carrying  
426 fluoroquinolone and beta-lactam resistance genes in *Escherichia coli* originating  
427 from diverse sources and geographical areas. *J Antimicrob Chemother* 71: 2118-  
428 2124.
- 429 11. Schwarz S, Johnson AP (2016) Transferable resistance to colistin: a new but old threat. *J*  
430 *Antimicrob Chemother* 71: 2066-2070.
- 431 12. Zhang JL, Wang JW, Chen L, Yassin AK, Kelly P, et al. (2018) Housefly (*Musca domestica*)  
432 and Blow Fly (*Protophormia terraenovae*) as Vectors of Bacteria Carrying Colistin  
433 Resistance Genes. *Applied and environmental microbiology* 84.
- 434 13. Guenther S, Falgenhauer L, Semmler T, Imirzalioglu C, Chakraborty T, et al. (2017)  
435 Environmental emission of multiresistant *Escherichia coli* carrying the colistin  
436 resistance gene *mcr-1* from German swine farms. *Journal of Antimicrobial*  
437 *Chemotherapy* 72: 1289-1292.
- 438 14. Wang R, Liu Y, Zhang Q, Jin L, Wang Q, et al. (2018) The prevalence of colistin resistance  
439 in *Escherichia coli* and *Klebsiella pneumoniae* isolated from food animals in China:  
440 coexistence of *mcr-1* and *bla*<sub>NDM</sub> with low fitness cost. *International journal of*  
441 *antimicrobial agents* 10.1016/j.ijantimicag.2018.01.023.
- 442 15. Li Y, Sun QL, Shen Y, Zhang Y, Yang JW, et al. (2018) Rapid increase in the prevalence of  
443 carbapenem-resistant Enterobacteriaceae (CRE) and emergence of colistin resistance  
444 gene *mcr-1* in CRE in a hospital in Henan, China. *Journal of clinical microbiology*  
445 10.1128/JCM.01932-17.
- 446 16. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, et al. (2015) Genomic analysis  
447 of diversity, population structure, virulence, and antimicrobial resistance in *Klebsiella*

448 pneumoniae, an urgent threat to public health. Proceedings of the National  
449 Academy of Sciences of the United States of America 112: E3574-3581.

450 17. KURAHASHI H, BUNCHU N (2011) The Blow flies recorded from Thailand, with the  
451 Description of  
452 a New Species of *Isomyia* WALKER (Diptera, Calliphoridae). Japanese Journal of Systematic  
453 Entomology 17: .

454 18. Bunchu N, Sukontason K, Sanit S, Chidburee P, Kurahashi H, et al. (2012) Occurrence of  
455 Blow Fly Species (Diptera: Calliphoridae) in Phitsanulok Province, Northern Thailand.  
456 Trop Biomed 29: 532-543.

457 19. Toleman MA (2018) Direct in Gel Genomic Detection of Antibiotic Resistance Genes in S1  
458 Pulsed Field Electrophoresis Gels. Methods Mol Biol 1736: 129-136.

459 20. Yu WL, Ko WC, Cheng KC, Lee CC, Lai CC, et al. (2008) Comparison of prevalence of  
460 virulence factors for *Klebsiella pneumoniae* liver abscesses between isolates with  
461 capsular K1/K2 and non-K1/K2 serotypes. Diagnostic microbiology and infectious  
462 disease 62: 1-6.

463 21. Lin TL, Yang FL, Yang AS, Peng HP, Li TL, et al. (2012) Amino Acid Substitutions of MagA  
464 in *Klebsiella pneumoniae* Affect the Biosynthesis of the Capsular Polysaccharide. PloS  
465 one 7.

466 22. Cheng HY, Chen YS, Wu CY, Chang HY, Lai YC, et al. (2010) RmpA regulation of capsular  
467 polysaccharide biosynthesis in *Klebsiella pneumoniae* CG43. Journal of bacteriology  
468 192: 3144-3158.

469 23. Chou HC, Lee CZ, Ma LC, Fang CT, Chang SC, et al. (2004) Isolation of a chromosomal  
470 region of *Klebsiella pneumoniae* associated with allantoin metabolism and liver  
471 infection. Infect Immun 72: 3783-3792.

472 24. Ma LC, Fang CT, Lee CZ, Shun CT, Wang JT (2005) Genomic heterogeneity in *Klebsiella*  
473 *pneumoniae* strains is associated with primary pyogenic liver abscess and metastatic  
474 infection. The Journal of infectious diseases 192: 117-128.

475 25. Lai YC, Lin GT, Yang SL, Chang HY, Peng HL (2003) Identification and characterization of  
476 KvgAS, a two-component system in *Klebsiella pneumoniae* CG43. FEMS microbiology  
477 letters 218: 121-126.

478 26. Hsieh PF, Lin TL, Lee CZ, Tsai SF, Wang JT (2008) Serum-induced iron-acquisition systems  
479 and TonB contribute to virulence in *Klebsiella pneumoniae* causing primary pyogenic  
480 liver abscess. The Journal of infectious diseases 197: 1717-1727.

481 27. Struve C, Bojer M, Krogfelt KA (2008) Characterization of *Klebsiella pneumoniae* type 1  
482 fimbriae by detection of phase variation during colonization and infection and  
483 impact on virulence. Infect Immun 76: 4055-4065.

484 28. Di Martino P, Cafferini N, Joly B, Darfeuille-Michaud A (2003) *Klebsiella pneumoniae*  
485 type 3 pili facilitate adherence and biofilm formation on abiotic surfaces. Res  
486 Microbiol 154: 9-16.

487 29. El Fertas-Aissani R, Messai Y, Alouache S, Bakour R (2013) Virulence profiles and  
488 antibiotic susceptibility patterns of *Klebsiella pneumoniae* strains isolated from  
489 different clinical specimens. Pathol Biol (Paris) 61: 209-216.

490 30. Thomas S, Holland IB, Schmitt L (2014) The Type 1 secretion pathway - the hemolysin  
491 system and beyond. Biochim Biophys Acta 1843: 1629-1641.

492 31. Liu Q, Bender RA (2007) Complex regulation of urease formation from the two  
493 promoters of the ure operon of *Klebsiella pneumoniae*. Journal of bacteriology 189:  
494 7593-7599.

- 495 32. Juhas M, Crook DW, Hood DW (2008) Type IV secretion systems: tools of bacterial  
496 horizontal gene transfer and virulence. *Cell Microbiol* 10: 2377-2386.
- 497 33. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, et al. (2016) Identification of  
498 *Klebsiella* capsule synthesis loci from whole genome data. *Microb Genom* 2:  
499 e000102.
- 500 34. Yang Q, Li M, Spiller OB, Andrey DO, Hinchliffe P, et al. (2017) Balancing *mcr-1*  
501 expression and bacterial survival is a delicate equilibrium between essential cellular  
502 defence mechanisms. *Nat Commun* 8: 2054.
- 503 35. Coutinho RL, Visconde MF, Descio FJ, Nicoletti AG, Pinto FC, et al. (2014) Community-  
504 acquired invasive liver abscess syndrome caused by a K1 serotype *Klebsiella*  
505 *pneumoniae* isolate in Brazil: a case report of hypervirulent ST23. *Mem Inst Oswaldo*  
506 *Cruz* 109: 970-971.
- 507 36. Podschun R, Ullmann U (1998) *Klebsiella* spp. as nosocomial pathogens: Epidemiology,  
508 taxonomy, typing methods, and pathogenicity factors. *Clinical microbiology reviews*  
509 11: 589-+.
- 510 37. Skov RL, Monnet DL (2016) Plasmid-mediated colistin resistance (*mcr-1* gene): three  
511 months later, the story unfolds. *Eurosurveillance* 21: 2-7.
- 512 38. Walsh TR (2018) A one-health approach to antimicrobial resistance. *Nat Microbiol* 3:  
513 854-855.
- 514 39. Wang Y, Zhang R, Li J, Wu Z, Yin W, et al. (2017) Comprehensive resistome analysis  
515 reveals the prevalence of NDM and MCR-1 in Chinese poultry production. *Nature*  
516 *Microbiology* 2: 16260.
- 517 40. Quan J, Li X, Chen Y, Jiang Y, Zhou Z, et al. (2017) Prevalence of *mcr-1* in *Escherichia coli*  
518 and *Klebsiella pneumoniae* recovered from bloodstream infections in China: a  
519 multicentre longitudinal study. *The Lancet Infectious Diseases* 17: 400-410.
- 520 41. Shen Y, Zhou H, Xu J, Wang Y, Zhang Q, et al. (2018) Anthropogenic and environmental  
521 factors associated with high incidence of *mcr-1* carriage in humans across China. *Nat*  
522 *Microbiol* 10.1038/s41564-018-0205-8.
- 523 42. Blaak H, Hamidjaja RA, van Hoek AH, de Heer L, de Roda Husman AM, et al. (2014)  
524 Detection of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli*  
525 on flies at poultry farms. *Applied and environmental microbiology* 80: 239-246.
- 526 43. Ahmad A, Ghosh A, Schal C, Zurek L (2011) Insects in confined swine operations carry a  
527 large antibiotic resistant and potentially virulent enterococcal community. *BMC*  
528 *microbiology* 11: 23.
- 529 44. Ahmad A, Nagaraja TG, Zurek L (2007) Transmission of *Escherichia coli* O157:H7 to cattle  
530 by house flies. *Prev Vet Med* 80: 74-81.
- 531 45. Matamoros S, van Hattem JM, Arcilla MS, Willemse N, Melles DC, et al. (2017) Global  
532 phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene  
533 indicates bacterial diversity but plasmid restriction. *Scientific reports* 7.
- 534 46. Partridge SR, Kwong SM, Firth N, Jensen SO (2018) Mobile Genetic Elements Associated  
535 with Antimicrobial Resistance. *Clin Microbiol Rev* 31.
- 536 47. Snesrud E, He S, Chandler M, Dekker JP, Hickman AB, et al. (2016) A Model for  
537 Transposition of the Colistin Resistance Gene *mcr-1* by IS*Apl1*. *Antimicrob Agents*  
538 *Chemother* 60: 6973-6976.
- 539 48. Snesrud E, McGann P, Chandler M (2018) The Birth and Demise of the IS*Apl1*-*mcr-1*-  
540 IS*Apl1* Composite Transposon: the Vehicle for Transferable Colistin Resistance. *Mbio*  
541 9.

542 49. Poirel L, Kieffer N, Nordmann P (2017) In Vitro Study of IS*Apl1*-Mediated Mobilization of  
543 the Colistin Resistance Gene *mcr-1*. *Antimicrobial agents and chemotherapy* 61.

544 50. Matamoros S, van Hattem JM, Arcilla MS, Willemse N, Melles DC, et al. (2017) Global  
545 phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene  
546 indicates bacterial diversity but plasmid restriction. *Sci Rep* 7: 15364.

547 51. Shen Y, Wu Z, Wang Y, Zhang R, Zhou HW, et al. (2018) Heterogeneous and Flexible  
548 Transmission of *mcr-1* in Hospital-Associated *Escherichia coli*. *MBio* 9.

549 52. Cohen Stuart J, van den Munckhof T, Voets G, Scharringa J, Fluit A, et al. (2012)  
550 Comparison of ESBL contamination in organic and conventional retail chicken meat.  
551 *International journal of food microbiology* 154: 212-214.

552 53. Cortes P, Blanc V, Mora A, Dahbi G, Blanco JE, et al. (2010) Isolation and Characterization  
553 of Potentially Pathogenic Antimicrobial-Resistant *Escherichia coli* Strains from  
554 Chicken and Pig Farms in Spain. *Applied and environmental microbiology* 76: 2799-  
555 2805.

556 54. Oteo J, Diestra K, Juan C, Bautista V, Novais A, et al. (2010) Extended-spectrum beta-  
557 lactamase-producing *Escherichia coli* in Spain belong to a large variety of multilocus  
558 sequence typing types, including ST10 complex/A, ST23 complex/A and ST131/B2  
559 (vol 34, pg 173, 2009). *International journal of antimicrobial agents* 36: 483-483.

560 55. Manges AR, Johnson JR (2012) Food-Borne Origins of *Escherichia coli* Causing  
561 Extraintestinal Infections. *Clinical Infectious Diseases* 55: 712-719.

562 56. Manges AR, Mende K, Murray CK, Johnston BD, Sokurenko EV, et al. (2017) Clonal  
563 distribution and associated characteristics of *Escherichia coli* clinical and surveillance  
564 isolates from a military medical center. *Diagnostic microbiology and infectious  
565 disease* 87: 382-385.

566 57. Bernasconi OJ, Kuenzli E, Pires J, Tinguely R, Carattoli A, et al. (2016) Travelers Can  
567 Import Colistin-Resistant Enterobacteriaceae, Including Those Possessing the  
568 Plasmid-Mediated *mcr-1* Gene. *Antimicrobial agents and chemotherapy* 60: 5080-  
569 5084.

570 58. Xavier BB, Lammens C, Butaye P, Goossens H, Malhotra-Kumar S (2016) Complete  
571 sequence of an IncFII plasmid harbouring the colistin resistance gene *mcr-1* isolated  
572 from Belgian pig farms. *Journal of Antimicrobial Chemotherapy* 71: 2342-2344.

573 59. Sun P, Bi Z, Nilsson M, Zheng B, Berglund B, et al. (2017) Occurrence of *blaKPC-2*, *blaCTX-*  
574 *M*, and *mcr-1* in Enterobacteriaceae from Well Water in Rural China. *Antimicrob  
575 Agents Chemother* 61.

576 60. Li A, Yang Y, Miao M, Chavda KD, Mediavilla JR, et al. (2016) Complete Sequences of  
577 *mcr-1*-Harboring Plasmids from Extended-Spectrum-beta-Lactamase- and  
578 Carbapenemase-Producing Enterobacteriaceae. *Antimicrob Agents Chemother* 60:  
579 4351-4354.

580 61. Rolain JM, Kempf M, Leangapichart T, Chabou S, Olaitan AO, et al. (2016) Plasmid-  
581 Mediated *mcr-1* Gene in Colistin-Resistant Clinical Isolates of *Klebsiella pneumoniae*  
582 in France and Laos. *Antimicrob Agents Chemother* 60: 6994-6995.

583 62. Srijan A, Margulieux KR, Ruekit S, Snesrud E, Maybank R, et al. (2018) Genomic  
584 Characterization of Nonclonal *mcr-1*-Positive Multidrug-Resistant *Klebsiella  
585 pneumoniae* from Clinical Samples in Thailand. *Microbial drug resistance* 24: 403-  
586 410.

- 587 63. Kieffer N, Aires-de-Sousa M, Nordmann P, Poirel L (2017) High Rate of MCR-1-Producing  
588 Escherichia coli and Klebsiella pneumoniae among Pigs, Portugal. Emerging  
589 infectious diseases 23: 2023-2029.
- 590 64. Ovejero CM, Delgado-Blas JF, Calero-Caceres W, Muniesa M, Gonzalez-Zorn B (2017)  
591 Spread of mcr-1-carrying Enterobacteriaceae in sewage water from Spain. J  
592 Antimicrob Chemother 72: 1050-1053.
- 593 65. Mendes AC, Novais A, Campos J, Rodrigues C, Santos C, et al. (2018) mcr-1 in  
594 Carbapenemase-Producing Klebsiella pneumoniae with Hospitalized Patients,  
595 Portugal, 2016-2017. Emerging infectious diseases 24: 762-766.
- 596 66. Lascols C, Peirano G, Hackel M, Laupland KB, Pitout JDD (2013) Surveillance and  
597 Molecular Epidemiology of Klebsiella pneumoniae Isolates That Produce  
598 Carbapenemases: First Report of OXA-48-Like Enzymes in North America.  
599 Antimicrobial agents and chemotherapy 57: 130-136.
- 600 67. Peirano G, Lascols C, Hackel M, Hoban DJ, Pitout JD (2014) Molecular epidemiology of  
601 Enterobacteriaceae that produce VIMs and IMPs from the SMART surveillance  
602 program. Diagnostic microbiology and infectious disease 78: 277-281.
- 603 68. Shankar C, Nabarro LE, Devanga Ragupathi NK, Muthurandhi Sethuvel DP, Daniel JL, et  
604 al. (2016) Draft Genome Sequences of Three Hypervirulent Carbapenem-Resistant  
605 Klebsiella pneumoniae Isolates from Bacteremia. Genome announcements 4.
- 606 69. Halaby T, Kucukkose E, Janssen AB, Rogers MR, Doorduyn DJ, et al. (2016) Genomic  
607 Characterization of Colistin Heteroresistance in Klebsiella pneumoniae during a  
608 Nosocomial Outbreak. Antimicrob Agents Chemother 60: 6837-6843.
- 609 70. Kayama S, Koba Y, Shigemoto N, Kuwahara R, Kakuham T, et al. (2015) Imipenem-  
610 Susceptible, Meropenem-Resistant Klebsiella pneumoniae Producing OXA-181 in  
611 Japan. Antimicrobial agents and chemotherapy 59: 1379-1380.
- 612 71. Olsen AR (1998) Regulatory action criteria for filth and other extraneous materials III.  
613 Review of flies and foodborne enteric disease. Regul Toxicol Pharm 28: 199-211.
- 614 72. Paczosa MK, Meccas J (2016) Klebsiella pneumoniae: Going on the Offense with a Strong  
615 Defense. Microbiology and Molecular Biology Reviews 80: 629-661.

617

618

619

620

621

622

623

624 **Table 1** characteristics of 48 MCRPE strains recovered from blowflies in Northern Thailand

Strain code	Species	Isolated area	colistin MIC(mg/L)	MLST	mcr-1-bearing Inc plasmid type	Conjugation frequency
-------------	---------	---------------	--------------------	------	--------------------------------	-----------------------

PN100	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	2.1x10 <sup>-5</sup>
PN104	<i>K. pneumoniae</i>	Rural area	16	43	IncX4	1.5x10 <sup>-6</sup>
PN105	<i>K. pneumoniae</i>	Suburb area	8	43	IncX4	1.5x10 <sup>-5</sup>
PN106	<i>K. pneumoniae</i>	Suburb area	8	43	IncX4	2.1x10 <sup>-5</sup>
PN107	<i>K. pneumoniae</i>	Rural area	16	43	IncX4	9.45x10 <sup>-7</sup>
PN110	<i>K. pneumoniae</i>	Rural area	16	43	IncX4	2.08x10 <sup>-6</sup>
PN114	<i>K. pneumoniae</i>	Suburb area	8	43	IncX4	7.63x10 <sup>-6</sup>
PN118	<i>K. pneumoniae</i>	Suburb area	8	43	IncX4	2.44x10 <sup>-5</sup>
PN120	<i>K. pneumoniae</i>	Rural area	4	43	IncX4	2.0x10 <sup>-5</sup>
PN77	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	1.5x10 <sup>-6</sup>
PN79	<i>K. pneumoniae</i>	Rural area	4	43	IncX4	3.13x10 <sup>-5</sup>
PN81	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	2.86x10 <sup>-5</sup>
PN84	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	3.0x10 <sup>-5</sup>
PN95	<i>K. pneumoniae</i>	Suburb area	8	43	IncX4	1.0x10 <sup>-5</sup>
PN96	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	7.11x10 <sup>-7</sup>
PN97	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	2.31x10 <sup>-6</sup>
PN98	<i>K. pneumoniae</i>	Suburb area	4	43	IncX4	8.03x10 <sup>-7</sup>
PN123	<i>E.coli</i>	Local market in urban community	4	2345	IncHI1A	1.44X10 <sup>-7</sup>
PN33	<i>E.coli</i>	Local market in urban community	4	10	NA	1.33x10 <sup>-3</sup>
PN93	<i>E.coli</i>	Local market in urban community	4	162	IncX4	NA
PN103	<i>E.coli</i>	Rural area	8	10	IncHI1A	3.1x10 <sup>-7</sup>
PN109	<i>E.coli</i>	Rural area	16	1244	IncX4	7.6x10 <sup>-4</sup>
PN111	<i>E.coli</i>	Rural area	8	457	NA	NA
PN116	<i>E.coli</i>	Rural area	4	648	NA	NA
PN119	<i>E.coli</i>	Rural area	8	549	NA	NA
PN74	<i>E.coli</i>	Rural area	16	10	IncX4	NA
PN75	<i>E.coli</i>	Rural area	4	58	IncHI1B	NA
PN87	<i>E.coli</i>	Rural area	4	549	NA	NA
PN88	<i>E.coli</i>	Rural area	4	10	IncX4	NA
PN91	<i>E.coli</i>	Rural area	4	181	IncX4	6.67x10 <sup>-5</sup>
PN101	<i>E.coli</i>	Suburb area	4	58	NA	NA
PN102	<i>E.coli</i>	Suburb area	4	648	NA	NA
PN108	<i>E.coli</i>	Suburb area	8	549	IncX4	NA
PN112	<i>E.coli</i>	Suburb area	8	181	IncX4	NA
PN117	<i>E.coli</i>	Suburb area	8	201	IncX4	5.94x10 <sup>-3</sup>
PN121	<i>E.coli</i>	Suburb area	8	5487	NA	NA
PN122	<i>E.coli</i>	Suburb area	8	549	NA	NA
PN124	<i>E.coli</i>	Suburb area	8	181	IncX4	5.33x10 <sup>-4</sup>
PN126	<i>E.coli</i>	Suburb area	4	648	IncHI1B	N.S
PN127	<i>E.coli</i>	Suburb area	4	648	IncHI1B	1.85x10 <sup>-7</sup>
PN73	<i>E.coli</i>	Suburb area	4	648	IncHI1B	N.S
PN76	<i>E.coli</i>	Suburb area	4	58	IncHI1A_IncHI1B	8.70x10 <sup>-8</sup>
PN78	<i>E.coli</i>	Suburb area	4	10	IncX4	NA
PN80	<i>E.coli</i>	Suburb area	8	218	NA	1.14x10 <sup>-4</sup>
PN83	<i>E.coli</i>	Suburb area	8	10	IncX4	6.27x10 <sup>-7</sup>
PN85	<i>E.coli</i>	Suburb area	4	2705	NA	N.S
PN86	<i>E.coli</i>	Suburb area	8	218	IncX4	NA
PN92	<i>E.coli</i>	Suburb area	4	10	NA	NA

625 MLST is analysed by seven allele sequence using MLST2.0 (See methods). NA, not available; N.S, not successful

626



627 **Fig.1** Phylogenetic trees of 31 MCRPEC isolates recovered from blowflies, were analysed by Parsnp  
628 in the Harvest package, and visualized by iTOL (<https://itol.embl.de/>). The pink circles indicate  
629 the presence of *mcr-1* gene. The presence or lack of AMR genes is colored in red or light yellow,  
630 respectively.

631

632

633 **Fig.2** PFGE analysis of MCR-1-producing strains digested with S1 nuclease (right) and hybridization  
634 with *mcr-1* gene probe (right). White arrows showed the location and size of *mcr-1*-carrying plasmids.

635

636

637 **Fig.3** (A). Alignment of 26 *mcr-1*-complete plasmids and visualized using BLAST Ring Image  
638 Generator (BRIG v0.9555). First inner ring is the plasmid obtained from PN105, used as reference for  
639 the alignment. (B) Alignment of 29 *mcr-1*-complete plasmids and visualized using BRIG  
640 v0.9555. First inner ring is the plasmid pMCR0617mcr used as a reference for the alignment,  
641 GenBank accession number and size of the reference plasmid indicated in the middle of rings.  
642 These *mcr-1*-carrying plasmids are recovered from different sources, namely, 10 representative  
643 *mcr-1*-linked plasmids from blowflies (PN78, PN83, PN86, PN88, PN91, PN93, PN112,  
644 PN117, PN109 and PN120), 2 from companion animals (PN10 and PN11), 4 from poultry  
645 (PN23, 24, 25 and 29), 11 from human feces (PN45, 42, 57, 41, 46, 51, 58, 60, 47, 49 and  
646 71). Besides those strains from blowflies fully described in Table 1, whole genomic sequence  
647 of all these MCRPE strains have been deposited in the NCBI database (see data availability).

648

649

650 **Fig.4** A. The image of *G. mellonella* over 12 h post-infection with ST43 MCRPKP strains and a clinical  
651 reference strain K1. B and C, Kaplan-Meier plots showing the percent survival of *G. mellonella* over

652 24 h post-infection with the  $10^4$  CFU/ml (B) and  $10^5$  CFU/ml (C) inoculum of MCRPKP and strain K1.  
653 Survival curves were plotted using the Kaplan-Meier method (GraphPad Software).

654

655 **Fig.5** A and B, Kaplan-Meier plots showing the percent survival of *G. mellonella* over 72 h post-  
656 infection with the  $10^5$  CFU/ml inoculum of clinical susceptible *K. pneumoniae* ff101 and clinical KPC-  
657 positive *K. pneumoniae* p35, with or without *mcr-1*-carrying plasmid. Survival curves were plotted  
658 using the Kaplan-Meier method (GraphPad Software).