

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/121525/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Wood, Ashley and Guggenheim, Jeremy 2019. Refractive error has minimal influence on the risk of age-related macular degeneration: A Mendelian randomization study. *American Journal of Ophthalmology* 206 , pp. 87-93. 10.1016/j.ajo.2019.03.018 Item availability restricted. file

Publishers page: <http://dx.doi.org/10.1016/j.ajo.2019.03.018>  
<<http://dx.doi.org/10.1016/j.ajo.2019.03.018>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## Abstract

**Purpose:** To test the hypothesis that refractive errors such as myopia and hyperopia cause an increased risk of age-related macular degeneration (AMD), and to quantify the degree of risk.

**Design:** Two-sample Mendelian randomization analysis of data from a genome-wide association study.

**Participants:** As instrumental variables for refractive error, 126 genome-wide significant genetic variants identified by the CREAM consortium and 23andMe Inc. were chosen. The association with refractive error for the 126 variants was obtained from a published study for a sample of n=95,505 European ancestry participants from UK Biobank. Association with AMD for the 126 genetic variants was determined from a genome-wide association study (GWAS) published by the International Age-related Macular Degeneration Genomics consortium of n=33,526 (16,144 cases and 17,832 controls) European ancestry participants.

**Methods:** Two-sample Mendelian randomization analysis was used to assess the causal role of refractive error on AMD risk, using the 126 genetic variants associated with refractive error as instrumental variables, under the assumption that the relationship between refractive error and AMD risk is linear.

**Main Outcome Measures:** The risk AMD caused by a 1 diopter (D) change in refractive error.

**Results:** Mendelian randomization analysis suggested that refractive error had very limited influence on the risk of AMD. Specifically, a 1 D more hyperopic refractive error was associated with an OR=1.080 (95% CI: 1.021 to 1.142, P=0.007) increased risk of AMD. MR-Egger, MR-PRESSO, weighted median, and Phenoscanner-based sensitivity analyses detected minimal evidence to suggest that this result was biased by horizontal pleiotropy.

**Conclusions:** Under the assumption of a linear relationship between refractive error and the risk of AMD, myopia and hyperopia only minimally influence the causal risk for AMD. Thus, inconsistently-reported strong associations between refractive error and AMD are likely to be the result of non-causal factors, such as stochastic variation, confounding or selection bias.

1  
2  
3  
4 **Abstract**  
5

6 Purpose: To test the hypothesis that refractive errors such as myopia and hyperopia cause an  
7 increased risk of age-related macular degeneration (AMD), and to quantify the degree of risk.  
8

9  
10 Design: Two-sample Mendelian randomization analysis of data from a genome-wide association  
11 study  
12

13 Participants: As instrumental variables for refractive error, 126 genome-wide significant genetic  
14 variants identified by the CREAM consortium and 23andMe Inc. were chosen. The association  
15 with refractive error for the 126 variants was obtained from a published study for a sample of  
16 n=95,505 European ancestry participants from UK Biobank. Association with AMD for the 126  
17 genetic variants was determined from a genome-wide association study (GWAS) published by  
18 the International Age-related Macular Degeneration Genomics consortium of n=33,526 (16,144  
19 cases and 17,832 controls) European ancestry participants.  
20

21  
22 Methods: Two-sample Mendelian randomization analysis was used to assess the causal role of  
23 refractive error on AMD risk, using the 126 genetic variants associated with refractive error as  
24 instrumental variables, under the assumption that the relationship between refractive error and  
25 AMD risk is linear.  
26

27  
28 Main Outcome Measures: The risk AMD caused by a 1 diopter (D) change in refractive error.  
29

30 Results: Mendelian randomization analysis suggested that refractive error had very limited  
31 influence on the risk of AMD. Specifically, a 1 D more hyperopic refractive error was associated  
32 with an OR=1.080 (95% CI: 1.021 to 1.142, P=0.007) increased risk of AMD. MR-Egger, MR-  
33 PRESSO, weighted median, and Phenoscanner-based sensitivity analyses detected minimal  
34 evidence to suggest that this result was biased by horizontal pleiotropy.  
35

36  
37 Conclusions: Under the assumption of a linear relationship between refractive error and the risk  
38 of AMD, myopia and hyperopia only minimally influence the causal risk for AMD. Thus,  
39 inconsistently-reported strong associations between refractive error and AMD are likely to be  
40 the result of non-causal factors, such as stochastic variation, confounding or selection bias.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

Age-related macular degeneration (AMD) is amongst the leading causes of visual impairment worldwide, and the leading cause in economically developed countries such as the United Kingdom where it is responsible for over 50% of registered visual impairment.<sup>1</sup> The socio-economic burden due to AMD-related visual impairment is set to increase even further as the elderly population expands.<sup>2,3</sup> AMD is a progressive condition that affects the macular region of the retina. The early stage is characterised by increasing drusen (number, size & confluence) and pigmentary abnormalities with vision usually minimally affected. Advanced disease may manifest either via gradual, progressive atrophy of the macula, or rapid development of sub-retinal neovascularisation leading to oedema, haemorrhage and eventual scar formation. Both atrophic and neovascular forms are associated with contemporaneous, and often severe, adverse impacts on central vision. Currently, treatment is only available for the neovascular form of AMD; the most effective treatment involving the regular injection of Anti-VEGF drugs into the eye.<sup>4-6</sup> The development of new treatments and therapies requires a better understanding of the aetiology and pathogenesis of AMD, so that disease mechanisms can be effectively targeted. Recognising *causal* risk factors for AMD is therefore key to this work.

Whilst the aetiology of AMD is not fully understood, it is clear that the condition results from a combination of genetic and environmental factors, the identification of which has been important in informing the understanding of AMD pathogenesis.<sup>7,8</sup> The principle risk factor for AMD is age.<sup>9-12</sup> Of the additional known risk factors, genetic predisposition and smoking have been the most consistently and confidently identified – the latter having particular significance as the risk can be modified with intervention. A systematic review by Chakravarthy et al. identified 73 potential risk factors for advanced AMD,<sup>9</sup> covering environmental (e.g. UV/sun exposure), demographic (e.g. age, race/ethnicity), genetic (e.g. Complement factor H), lifestyle (e.g. smoking, dietary fats & anti-oxidants), general health (e.g. cardiovascular disease) and ocular comorbidity (e.g. refractive error) factors. Nevertheless, few of these potential risk factors for AMD have been evaluated as part of a randomised control trial (RCT) designed to test for a causal role. Whilst RCTs are the gold standard for demonstrating causation, they are resource intensive and are not always practical; e.g. if a long-term intervention is required, an RCT may not be feasible, or if exposure to a risk factor poses a potential health risk, an RCT may be unethical.

Refractive error, in particular hyperopia, has been suspected to increase the risk of AMD since at least the late 1970's.<sup>13</sup> However, despite research over many decades, the evidence from cross-sectional and cohort studies has been inconsistent.<sup>14,15</sup> Standard cross-sectional 'observational studies' are susceptible to confounding and therefore are unable to establish a causal association between an exposure (e.g. degree of refractive error) and an outcome (e.g. patient has AMD).<sup>16,17</sup> Mendelian Randomization (MR) analyses have been proposed as a research method for drawing valid causal inferences using existing, cross-sectional datasets,<sup>16,18</sup> either as an alternative to RCTs in circumstances when an RCT is not practical, or as a precursor before embarking on a lengthy and costly RCT.<sup>19</sup> MR utilizes genetic variants that explain variation in an exposure as 'instrumental variables'<sup>20</sup> in order to quantify the effect of the exposure on the outcome independent of confounding factors that influence the exposure-outcome relationship. The ability to attribute a causal effect in MR depends primarily on three key assumptions: (1) that the genetic variants are robustly associated with the exposure, (2) the genetic variants only affect the outcome via the exposure, and (3) the genetic variants do not exert effects on confounders of the exposure-outcome relationship. Here, we sought to test the hypothesis that refractive error has a causal impact on the risk of AMD by applying Mendelian randomization, using as instrumental variables genetic variants associated with refractive error in a recent, large-scale GWAS.

## Methods

The study design is a two sample Mendelian randomization utilising GWAS summary statistics from CREAM, 23andMe, UK Biobank and IAMDGC. Participants provided informed consent to take part in the respective studies, which adhered to the principles of the Declaration of Helsinki.<sup>21,22</sup> The IRB (School of Optometry and Vision Sciences Research Audit Ethics Committee) provided a waiver confirming approval was not required as this was a retrospective analysis of data already in the public domain.

### ***Instrumental variables for refractive error***

Tedja et al.<sup>21</sup> reported the largest GWAS meta-analysis for refractive error to date. The GWAS meta-analysis included data for n=160,420 participants of European ancestry from the CREAM consortium and the 23andMe personal genomics company. The CREAM consortium GWAS was performed for the trait 'spherical equivalent refractive error' (SER) whereas the trait analysed in the 23andMe GWAS was 'age of diagnosis of myopia' (AODM). Tedja et al. identified 161 lead variants, where a lead variant was defined as 'the variant with the lowest p-value in a 100-kb window of the outermost genome-wide-significant variant of that same region'.<sup>21</sup> These 161 variants with  $P < 5 \times 10^{-8}$  in the CREAM/23andMe GWAS were tested for replication in an independent sample of n=95,505 participants of European ancestry aged 37–73 years old from UK Biobank who had no history of eye disorders<sup>23</sup> All of the UK Biobank participants had phenotype information for SER measured using non-cycloplegic autorefractometry in diopters (D). The mean  $\pm$  standard deviation age of the UK Biobank GWAS sample was 57.7  $\pm$  7.9 years, and 53.1% were female. A total of 149/161 variants provided independent evidence of replication in the UK Biobank sample ( $P < 0.05$ ), after excluding one tri-allelic variant.<sup>21</sup> From amongst these 149 variants, 18 (rs10003846, rs11723482, rs1207782, rs13069734, rs1550094, rs17400325, rs17837871, rs1994840, rs2276560, rs2326823, rs2573081, rs2823097, rs6903823, rs72621438, rs745480, rs7925340, rs79266634 and rs79953651) were excluded as potential instrumental variables for having a pairwise linkage disequilibrium (LD)  $r^2 > 0.05$  with another variant in the set and a further 2 markers (rs1983554 and rs931302) were excluded for having a Hardy Weinberg equilibrium test  $P < 0.05$ , leaving 129 independent variants. There were 24 palindromic single nucleotide polymorphisms (SNPs), i.e. SNPs with alleles A/T or C/G, amongst these 129 variants. To avoid the chance of incorrectly harmonizing alleles in the 2-sample MR analysis<sup>24</sup> palindromic SNPs were replaced by proxies in high LD ( $r^2 > 0.8$ ); proxies were available for 21 of the 24 palindromic SNPs, leaving a final set of 126 variants which were used as instrumental variables for refractive error in the current study (no proxy was available for rs2908972, rs74764079 or rs807037). The degree of association with refractive error for the variants was taken from the UK Biobank replication sample, rather than the larger CREAM/23andMe discovery sample, since the trait analysed in the UK Biobank GWAS was SER for all participants. The summary statistics for the 126 refractive error instrumental variables are shown in Supplementary Table 1 (available at [www.aaojournal.org](http://www.aaojournal.org)).

### ***Association of instrumental variables with age-related macular degeneration***

Fritsche et al.<sup>22</sup> reported a GWAS meta-analysis for AMD carried out by the International Age-related Macular Degeneration Genomics Consortium (IAMDGC). This meta-analysis examined a 'discovery sample' of n=33,526 individuals (16,144 cases and 17,832 controls) recruited across 26 studies. AMD cases were defined as individuals with either (i) geographic atrophy and/or choroidal neovascularization in at least one eye and an age at first diagnosis  $\geq 50$  years ('advanced AMD'), or (ii) pigmentary changes in the RPE or more than five macular drusen of diameter  $\geq 63\mu\text{m}$  or greater and age at first diagnosis  $\geq 50$  years ('intermediate AMD'). Control participants had a mean  $\pm$  standard deviation age of 70.7  $\pm$  9.7 years and were free from advanced or intermediate AMD. The GWAS meta-analysis summary statistics reported by the

1  
2  
3  
4 IAMDGC (<http://csg.sph.umich.edu/abecasis/public/amd2015/>) included the effect allele,  
5 reference allele, direction of effect (i.e. increased or decreased risk of AMD) and *P*-value. For  
6 MR analysis, the log(odds ratio) (log(OR)) and corresponding standard errors for this dataset  
7 were calculated as described by Davey Smith and Burgess.<sup>25</sup> Of the 126 genetic variants  
8 selected as instrumental variables for refractive error, summary data for all 126 were available  
9 in the AMD GWAS dataset.  
10

### 11 **Statistical analysis**

12 All analyses were carried out with R.<sup>26</sup> Inverse variance-weighted (IVW) MR,<sup>27</sup> under a  
13 multiplicative random-effects model, weighted median MR<sup>28</sup> and MR-Egger analyses<sup>29</sup> were  
14 carried out with the *MendelianRandomisation* package.<sup>30</sup> MR-PRESSO analysis was performed  
15 with the *MRPRESSO* package, with *K*=100,000 simulations.<sup>31</sup> Scatter plots were generated with  
16 the *ggplot2* package. I<sup>2</sup> heterogeneity statistics were calculated using the *metafor* package.<sup>32</sup>  
17 From amongst the 126 variants associated with refractive error selected as instrumental  
18 variables, those with known pleiotropic effects on additional traits were identified using  
19 Phenoscanner<sup>33</sup> with the settings  $P < 5 \times 10^{-8}$  and inclusion of proxy variants in LD ( $r^2 > 0.8$ ) in  
20 Europeans. The Phenoscanner analysis identified 31 variants with known pleiotropic effects  
21 (Supplementary Table 1; available at [www.aaojournal.org](http://www.aaojournal.org)). The variance in refractive error  
22 explained by the 126 instrumental variables was assessed as described by Ghorbani Mojarrad  
23 et al.<sup>34</sup> in a sample of *n*=1,516 unrelated female adults from the UK (mean  $\pm$  standard deviation  
24 age=44.6  $\pm$  4.4 years) whose refractive error was assessed by non-cycloplegic autorefractometry.  
25 Statistical power was assessed as described by Brion et al.,<sup>35</sup> using the online tool:  
26 <http://cns.genomics.com/shiny/mRnd/>.  
27  
28  
29  
30

### 31 **Results**

32 One hundred and twenty-six genetic variants associated with refractive error were chosen as  
33 instrumental variables for refractive error. The 126 genetic variants explained 4.4% of the  
34 variance in refractive error in an independent sample of UK adults, and the F-statistic from the  
35 first stage of the MR analysis was 1544.0. Both of these findings suggested there was minimal  
36 risk of 'weak instrument bias'.<sup>34</sup> The study had 45% power to detect an OR=1.10 increased risk  
37 of AMD per 1 D more hyperopic refractive error, and 80% power to detect an OR=1.16  
38 increased risk. Hence, the study was well-powered to detect risks below OR=0.85 or above  
39 OR=1.15, but was not well powered to detect very small risks close to the null value.  
40  
41

42 A quantile-quantile (QQ) plot for the 126 refractive error instrumental variables, designed to  
43 illustrate their association with AMD, demonstrated an excess of low *P*-values compared to that  
44 expected under the null hypothesis (Figure 1). In particular, two of the refractive error variants  
45 were very strongly associated with AMD risk: rs10760673, an intronic variant within the  
46 *TGFBR1* gene (AMD risk  $P=4.59\text{e-}09$ ), and rs6420484, a missense variant in the *TSPAN10*  
47 gene (AMD risk  $P=4.11\text{e-}11$ ). Both genetic loci were already well-known to be associated with  
48 AMD.<sup>22</sup>  
49  
50

51 <Figure 1 near here>  
52

53 A standard, inverse variance-weighted (IVW) MR analysis using all 126 genetic variants  
54 suggested that each 1.00 D change in refractive error in the direction of more hyperopia was  
55 associated with an approximately 8% increase in the risk of AMD (OR=1.080, 95% CI: 1.021 to  
56 1.142,  $P=0.007$ ; Table 1 and Figure 2).  
57  
58

59 <Figure 2 near here>  
60 <Table 1 near here>  
61  
62

1  
2  
3  
4  
5  
6  
7 The standard IVW MR analysis including all 126 genetic variants exhibited strong evidence of  
8 heterogeneity (Cochran  $Q=326.1$ ,  $P=7.1e-20$ ;  $I^2=61.6\%$ ), implying that the SNP-exposure vs.  
9 SNP-outcome relationship varied widely across variants. The most likely reason for such  
10 heterogeneity in an MR analysis is 'horizontal' pleiotropy.<sup>18</sup> Hence, a series of sensitivity  
11 analyses were undertaken to examine the robustness of the IVW MR causal effect estimate  
12 (Tables 1-3). A weighted median-based MR analysis, which provides a valid causal effect  
13 estimate if up to half of the information is from invalid instrumental variables, produced a similar  
14 estimate to that obtained from the IVW MR but with weaker statistical support. Specifically, while  
15 the median-based MR estimate was consistent with the IVW MR estimate, the 95% confidence  
16 interval included the null value (OR=1.045, 95% CI: 0.982 to 1.111,  $P=0.164$ ; Table 1),  
17 commensurate with the reduced statistical power of median-based MR compared to IVW MR.<sup>28</sup>  
18 An MR-Egger analysis found negligible evidence of directional pleiotropy: MR-Egger  
19 intercept=1.003 (95% CI: 0.991 to 1.016). Repeating the above analyses after excluding the 2  
20 variants (rs10760673 and rs6420484) very strongly associated with AMD risk had minimal  
21 impact on the results (Table 2). In addition to the 2 variants associated with AMD, a  
22 PhenoScanner analysis<sup>33</sup> identified a further 29 genetic variants with known effects on traits  
23 unrelated to refractive error. MR analysis after exclusion of these 31 variants from the original  
24 set of 126 variants produced comparable findings to the original analyses (Table 3). For  
25 example, the IVW MR causal effect estimate was OR=1.069 (95% CI: 1.016 to 1.124,  $P=0.010$ ).  
26 Heterogeneity was partially reduced in this analysis, compared to the original analysis with all  
27 126 variants (Cochran  $Q=161.6$ ,  $P=1.81e-05$ ;  $I^2=39.1\%$ ). Finally, an MR-PRESSO analysis  
28 identified 6 variants as pleiotropic outliers (Supplementary Table 1; available at  
29 [www.aaojournal.org](http://www.aaojournal.org)). After removal of these 6 variants, the MR-PRESSO causal effect estimate  
30 was OR=1.104 (95% CI: 1.054 to 1.157,  $P=8.12e-05$ ).  
31  
32

33  
34 <Table 2 near here>

35 <Table 3 near here>

36  
37  
38 In summary, a wide range of MR analysis models all produced causal effect estimates close to  
39 the null, with the most highly statistically powered analysis estimating an approximately 8-10%  
40 increased risk of AMD per 1 D more hyperopic refractive error.  
41

## 42 Discussion

43 We performed a two-sample MR analysis to determine if a causal relationship exists between  
44 refractive error and the risk of age-related macular degeneration, based on genetic data from  
45 populations with European ancestry. The results suggested that the SER does indeed have a  
46 causal relationship with AMD risk, albeit with the risk of AMD increasing by less than 10% per  
47 1 D increase in hyperopia. This study provides genetic evidence to support a causal link  
48 between refractive error and AMD risk, which while not as definitive as that obtained from an  
49 RCT, provides greater freedom from confounder bias than the risks reported in the  
50 observational epidemiological studies conducted to date.  
51  
52

53 A meta-analysis carried out by Pan et al.<sup>14</sup> suggested that each 1 D increase in hyperopia is  
54 associated with a 6–9% increased risk of AMD, based on pooled responses across 2 cohort and  
55 5 cross-sectional studies. A further meta-analysis by Li et al.<sup>15</sup>, using largely overlapping study  
56 samples as Pan et al.,<sup>14</sup> perhaps unsurprisingly found a similarly increased risk of 6–10% per  
57 diopter increase in hyperopia. Both figures are consistent with the upper estimate found in our  
58 analyses, of an approximately 8-10% increased risk of AMD per diopter of hyperopia. However,  
59 when making this comparison, it is important to note Pan et al.<sup>14</sup> observed that – other than age  
60  
61

1  
2  
3  
4 – many important confounders, such as smoking status, education level and socio-economic  
5 status were not accounted for across all the meta-analysed studies. Furthermore, neither meta-  
6 analysis<sup>14,15</sup> limited inclusion to individuals of European ancestry, in contrast to our MR  
7 analysis.  
8

9  
10 Various biologically plausible explanations have been proposed for the relationship between  
11 refractive error and AMD risk. Explanations previously explored include posterior vitreous  
12 detachment (PVD), the prevalence of which is greater in myopic eyes,<sup>36</sup> which has been  
13 suggested to reduce the likelihood of neovascularisation<sup>14,37</sup> (with the PVD hypothesized as  
14 removing a barrier to diffusion of VEGF away from the macular). Also, the VEGF concentration  
15 in the retina have been reported to be lower in myopic vs. hyperopic eyes, leading Jonas et al.<sup>38</sup>  
16 to propose that this reduced VEGF concentration may influence the risk of AMD. Yet these  
17 explanations are not consistent with the frequently-observed association between refractive  
18 error and early AMD as opposed to advanced AMD.<sup>9,14,15</sup>  
19

20  
21 Others, such as Pan et al.<sup>39</sup> have discussed possible links between light exposure and refractive  
22 error, in this case speculating that spectacle lens wear for refractive error may reduce UV  
23 exposure, although they noted that only a small proportion of their study sample classified as  
24 having either myopia or hyperopia did not wear spectacles. Furthermore, Quigley et al.<sup>40</sup>  
25 contended that spectacle lenses would have negligible filtering effect on the short wavelength  
26 visible light commonly linked to AMD risk. Quigley et al.<sup>40</sup> proposed that overall retinal light  
27 exposure, which using two separate methods was demonstrated to increase with hyperopic  
28 refractive error (by proxy of axial eye length), was a more likely potential mechanism by which  
29 refractive error may be linked to increased risk of AMD.  
30

31  
32 In addition to arguments centering on the retina, a greater rigidity of the sclera in shorter,  
33 hyperopic eyes has also been proposed,<sup>14,39,41,42</sup> since rigidity of the sclera has been identified  
34 as a risk factor in neovascular AMD.<sup>43</sup> Finally, myopic eyes characteristically have thinner  
35 choroids than hyperopic eyes.<sup>44</sup> While this might imply that the choroidal contribution to removal  
36 of photoreceptor phagocytosis breakdown products may be relatively impaired in myopic eyes,<sup>45</sup>  
37 this is not borne out by the smaller size of drusen in myopic eyes.<sup>46,47</sup>  
38

39  
40 This work had a number of limitations. Firstly, while we took steps to minimize the risk of bias  
41 from the potential use of invalid instrumental variables (due, for example, to horizontal  
42 pleiotropy) by performing MR-Egger, MR-PRESSO and weighted median-based MR sensitivity  
43 analyses, we cannot completely rule out such bias. Secondly, we made the assumption of a  
44 linear relationship between refractive error and the risk of AMD. Again, while observational  
45 studies of axial eye length (a surrogate for refractive error) and AMD risk suggest that this  
46 assumption was reasonable, data from a very large case-control sample with information on  
47 both AMD status and refractive error would be required to test it formally.<sup>48</sup> Thirdly, with regards  
48 to the instrumental variable assumptions necessary for a valid MR analysis, the 126 genetic  
49 variants used all demonstrated an association with refractive error in an independent replication  
50 sample (UK Biobank) distinct from the discovery sample (CREAM and 23andMe). The variants  
51 together explained 4.4% of the variance in refractive error in an additional, independent sample  
52 of participants. Thus, the variants satisfied the first MR assumption of displaying robust  
53 association with the exposure. There was evidence of heterogeneity in the genetic variant-  
54 exposure vs. genetic variant-outcome relationship ( $Q=326.1, I^2=61.6\%$ ) in the analysis using all  
55 126 variants, suggesting that either the second and/or third MR assumptions were not met. The  
56 most likely reason for this was horizontal pleiotropy, i.e. variants conferring a risk of AMD  
57 through a pathway other than via a direct effect on refractive error. While the sensitivity  
58 analyses reduced the level of residual heterogeneity, they did not exclude it completely.  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

However, the similarity in the causal effect estimates obtained with the original IVW MR analysis and the various sensitivity analyses provides reassurance that much of the original heterogeneity did not appreciably bias the results. However, as mentioned above, we cannot completely rule out bias due to horizontal pleiotropy that was still present in all of our sensitivity analyses. Fourthly, theoretically a 2-sample MR study design is at risk of confounding due to population stratification (if an instrumental variable tags groups of individuals with different ancestries and these groups differ in the prevalence of the outcome<sup>16</sup>). Since all 126 instrumental variables in the current study were associated with refractive error in UK Biobank participants carefully selected as having homogeneous European ancestry, and association with AMD was also assessed in a sample restricted to those of European ancestry, the risk of confounding due to population stratification is extremely low.

In summary, Mendelian randomization analysis provided evidence that hyperopic eyes have an increased risk of AMD, but that the causal effect size is modest (OR=1.08 per diopter,  $P=0.007$ ). This degree of protection is consistent with that estimated in two large meta-analyses of cross-sectional and longitudinal observational studies (OR=1.06–1.10 per diopter), which implies that confounder bias did not strongly impact previous risk estimates in these two studies. The increasing prevalence of myopia<sup>49</sup> and consequent reduction in the prevalence of hyperopia<sup>50</sup> may serve to partially counter the higher incidence of AMD due to increased life expectancy across most of the world. Nevertheless, this work emphasizes that risk factors other than refractive error, such as smoking, have a much greater impact on AMD risk.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Acknowledgments**

a) Funding/Support: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

b) Financial Disclosures: Dr. Wood has nothing to disclose; Dr. Guggenheim has nothing to disclose.

c) Other: This research has been conducted using the UK Biobank Resource (application #17351).

## References

1. Quartilho A, Simkiss P, Zekite A, Xing W, Wormald R, Bunce C. Leading causes of certifiable visual loss in England and Wales during the year ending 31 March 2013. *Eye*. 2016;30(4):602-607.
2. Pezzullo L, Streatfeild J, Simkiss P, Shickle D. The economic impact of sight loss and blindness in the UK adult population. *BMC Health Serv Res*. 2018;18:63.
3. Chamie J. World population prospects: The 2004 revision. *New York United Nations Dep Econ Soc Aff*. 2005.
4. Yonekawa Y, Miller J, Kim I, Yonekawa Y, Miller JW, Kim IK. Age-Related Macular Degeneration: Advances in Management and Diagnosis. *J Clin Med*. 2015;4(2):343-359.
5. Martin DF, Maguire MG, Fine SL, et al. Ranibizumab and bevacizumab for treatment of neovascular age-related macular degeneration: two-year results. *Ophthalmology*. 2012;119(7):1388-1398.
6. Rosenfeld PJ, Brown DM, Heier JS, et al. Ranibizumab for Neovascular Age-Related Macular Degeneration. *N Engl J Med*. 2006;355(14):1419-1431.
7. Ambati J, Fowler BJ. Mechanisms of age-related macular degeneration. *Neuron*. 2012;75(1):26-39.
8. Lim LS, Mitchell P, Seddon JM, Holz FG, Wong TY. Age-related macular degeneration. *Lancet*. 2012;379(9827):1728-1738.
9. Chakravarthy U, Wong TY, Fletcher A, et al. Clinical risk factors for age-related macular degeneration: a systematic review and meta-analysis. *BMC Ophthalmol*. 2010;10:31.
10. Cheung CMG, Tai ES, Kawasaki R, et al. Prevalence of and risk factors for age-related macular degeneration in a multiethnic Asian cohort. *Arch Ophthalmol*. 2012;130(4):480-486.
11. Clemons TE, Milton RC, Klein R, Seddon JM, Ferris FL. Risk factors for the incidence of Advanced Age-Related Macular Degeneration in the Age-Related Eye Disease Study (AREDS) AREDS report no. 19. *Ophthalmology*. 2005;112(4):533-539.
12. Lawrenson JG, Evans JR. Advice about diet and smoking for people with or at risk of age-related macular degeneration: a cross-sectional survey of eye care professionals in the UK. *BMC Public Health*. 2013;13:564.
13. Maltzman BA, Mulvihill MN, Greenbaum A. Senile macular degeneration and risk factors: a case-control study. *Ann Ophthalmol*. 1979;11(8):1197-1201.
14. Pan CW, Ikram MK, Cheung CY, et al. Refractive errors and age-related macular degeneration: A systematic review and meta-analysis. *Ophthalmology*. 2013;120(10):2058-2065.
15. Li Y, Wang J, Zhong X, et al. Refractive Error and Risk of Early or Late Age-Related Macular Degeneration: A Systematic Review and Meta-Analysis. *PLoS One*. 2014;9(3):e90897. doi:10.1371/journal.pone.0090897.
16. Lawlor DA, Harbord RM, Sterne JAC, Timpson N, Smith GD, Davey Smith G. Mendelian randomization: Using genes as instruments for making causal inferences in epidemiology. *Stat Med*. 2008;27(8):1133-1163.

17. Vansteelandt S, Dukes O, Martinussen T. Survivor bias in Mendelian randomization analysis. *Biostatistics*. 2017;19(4):426-443.
18. Burgess S, Foley CN, Zuber V. Inferring Causal Relationships Between Risk Factors and Outcomes from Genome-Wide Association Study Data. *Annu Rev Genom Hum Genet*. 2018;19:303-327.
19. Burgess S, Malarstig A. Using Mendelian randomization to assess and develop clinical interventions: limitations and benefits. *J Comp Eff Res*. 2013;2(3):209-212.
20. Swanson SA, Labrecque J, Hernán MA. Causal null hypotheses of sustained treatment strategies: What can be tested with an instrumental variable? *Eur J Epidemiol*. 2018;33(8):723-728.
21. Tedja MS, Wojciechowski R, Hysi PG, et al. Genome-wide association meta-analysis highlights light-induced signaling as a driver for refractive error. *Nat Genet*. 2018;50:834-848.
22. Fritsche LG, Igl W, Bailey JNC, et al. A large genome-wide association study of age-related macular degeneration highlights contributions of rare and common variants. *Nat Genet*. 2015;48(2):134-143.
23. Plotnikov D, Guggenheim J, Consortium TUBE and V. Is a large eye size a risk factor for myopia? A Mendelian randomization study. *bioRxiv*. December 2017:240283. doi:10.1101/240283.
24. Hartwig FP, Davies NM, Hemani G, Smith GD. Two-sample Mendelian randomization: avoiding the downsides of a powerful, widely applicable but potentially fallible technique. *Int J Epidemiol*. 2016;45(6):1717-1726.
25. Burgess S, Davey Smith G. Mendelian Randomization Implicates High-Density Lipoprotein Cholesterol–Associated Mechanisms in Etiology of Age-Related Macular Degeneration. *Ophthalmology*. 2017;124(8):1165-1174.
26. R Development Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria; 2008.
27. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol*. 2013;37(7):658-665.
28. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol*. 2016;40(4):304-314.
29. Bowden J, Smith GD, Burgess S. Mendelian randomization with invalid instruments: Effect estimation and bias detection through Egger regression. *Int J Epidemiol*. 2015;44(2):512-525.
30. Yavorska OO, Burgess S. MendelianRandomization: An R package for performing Mendelian randomization analyses using summarized data. *Int J Epidemiol*. 2017;46(6):1734-1739.
31. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet*. 2018;50(5):693-698.

- 1
- 2
- 3
- 4 32. Viechtbauer W. Conducting meta-analysis in R with metafor package. *J Stat Softw.* 2010;36(3):1-48.
- 5
- 6
- 7 33. Staley JR, Blackshaw J, Kamat MA, et al. PhenoScanner: A database of human
- 8 genotype-phenotype associations. *Bioinformatics.* 2016;32(20):3207-3209.
- 9
- 10 34. Ghorbani Mojarrad N, Williams C, Guggenheim JA. A genetic risk score and number of
- 11 myopic parents independently predict myopia. *Ophthalmic Physiol Opt.* 2018;38(5):492-
- 12 502.
- 13
- 14 35. Brion MJA, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian
- 15 randomization studies. *Int J Epidemiol.* 2013;42(5):1497-1501.
- 16
- 17 36. Saw SM, Gazzard G, Shin-Yen EC, Chua WH. Myopia and associated pathological
- 18 complications. *Ophthalmic Physiol Opt.* 2005;25(5):381-391.
- 19
- 20 37. Simpson ARH, Petrarca R, Jackson TL. Vitreomacular Adhesion and Neovascular Age-
- 21 Related Macular Degeneration. *Surv Ophthalmol.* 2012;57(6):498-509.
- 22
- 23 38. Jonas JB, Tao Y, Neumaier M, Findeisen P. VEGF and refractive error. *Ophthalmology.*
- 24 2010;117(11):2234.
- 25
- 26 39. Pan CW, Cheung CY, Aung T, et al. Differential associations of myopia with major age-
- 27 related eye diseases: The Singapore Indian eye study. *Ophthalmology.* 2013;120(2):284-
- 28 291.
- 29
- 30 40. Quigley MG, Powell I, Wittich W. Increased axial length corresponds to decreased retinal
- 31 light dose: A parsimonious explanation for decreasing AMD risk in myopia. *Invest*
- 32 *Ophthalmol Vis Sci.* 2018;59(10):3852-3857.
- 33
- 34 41. Fraser-Bell S, Choudhury F, Klein R, Azen S, Varma R. Ocular Risk Factors for Age-
- 35 Related Macular Degeneration: The Los Angeles Latino Eye Study. *Am J Ophthalmol.*
- 36 2010;149(5):735-740.
- 37
- 38 42. Ikram MK, Van Leeuwen R, Vingerling JR, Hofman A, De Jong PTVM. Relationship
- 39 between refraction and prevalent as well as incident age-related maculopathy: The
- 40 Rotterdam study. *Invest Ophthalmol Vis Sci.* 2003;44(9):3778-3782.
- 41
- 42 43. Pallikaris IG, Kymionis GD, Ginis HS, Kounis GA, Christodoulakis E, Tsilimbaris MK.
- 43 Ocular rigidity in patients with age-related macular degeneration. *Am J Ophthalmol.*
- 44 2006;141(4):611-615.
- 45
- 46 44. Nickla DL, Wallman J. The multifunctional choroid. *Prog Retin Eye Res.* 2010;29(2):144-
- 47 168.
- 48
- 49 45. Lee JY, Lee DH, Lee JY, Yoon YH. Correlation between subfoveal choroidal thickness
- 50 and the severity or progression of nonexudative age-related macular degeneration. *Invest*
- 51 *Ophthalmol Vis Sci.* 2013;54(12):7812-7818.
- 52
- 53 46. Xu L, Li Y, Wang S, Wang Y, Wang Y, Jonas JB. Characteristics of Highly Myopic Eyes.
- 54 The Beijing Eye Study. *Ophthalmology.* 2007;114(1):121-126.
- 55
- 56 47. Anand R, Bressler SB, Davis MD, Ferris III FL, Klein R. Risk factors associated with age-
- 57 related macular degeneration. A case-control study in the age-related eye disease study:
- 58 Age-Related Eye Disease Study Report Number 3. *Ophthalmology.* 2000;107(12):2224-
- 59 2232.
- 60
- 61
- 62
- 63
- 64
- 65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

48. Staley JR, Burgess S. Semiparametric methods for estimation of a nonlinear exposure-outcome relationship using instrumental variables with application to Mendelian randomization. *Genet Epidemiol.* 2017;41(4):341-352.

49. Morgan IG, Ohno-Matsui K, Saw SM. Myopia. *Lancet.* 2012;379(9827):1739-1748.

50. Wang H, Naghavi M, Allen C, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet.* 2016;388(10053):1459-1544.

1  
2  
3  
4 **Figure captions**  
5

6 **Figure 1. Quantile-quantile plot illustrating observed level of association with AMD risk**  
7 **(negative log<sub>10</sub> p-value) for 126 genetic variants used as instrumental variables for**  
8 **refractive error.** Note the inflation of observed negative log<sub>10</sub> p-values compared to that  
9 expected under the null hypothesis of no association with AMD risk. The red line is the line of  
10 unity and the grey shaded region is the 95% confidence interval expected under the null  
11 hypothesis. Each point represents an individual genetic variant.  
12

13 **Figure 2. Graphical representation of 2-sample Mendelian randomization meta-analysis**  
14 **results.** Each point represents an individual genetic variant (instrumental variable) with error  
15 bars indicating the standard error. The association of each genetic variant with refractive error  
16 (in units of diopters per copy of the risk allele) is plotted on the x-axis, and the association with  
17 AMD risk (log odds ratio) is plotted on the y-axis. The solid red line shows the fit from an inverse  
18 variance weighted (IVW) MR meta-analysis model, while the dashed red line shows the fit for an  
19 MR-Egger model.  
20  
21

22 **Table 1.** Mendelian randomization causal effect estimate for the risk of AMD (odds ratio) per 1  
23 diopter more hyperopic refractive error.  
24

25 **Table 2.** Sensitivity analysis results for Mendelian randomization analyses after excluding 2  
26 variants (rs10760673 and rs6420484) strongly associated with AMD risk.  
27

28 **Table 3.** Sensitivity analysis results for Mendelian randomization analyses after excluding 31  
29 variants strongly associated with non-refractive error traits identified by Phenoscanner.  
30

31 **Supplementary Table 1.** is available in the Online Supplementary Material.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1. Mendelian randomization causal effect estimate for the risk of AMD (odds ratio) per 1 diopter more hyperopic refractive error.**

<b>Analysis</b>	<b>Causal effect (OR per D)</b>	<b>95% Confidence interval</b>	<b>P-value</b>
Inverse variance-weighted	1.080	1.021 to 1.142	0.007
Weighted median	1.045	0.982 to 1.111	0.164
MR-Egger	1.048	0.922 to 1.190	0.474
<b>Analysis</b>	<b>Intercept</b>	<b>95% Confidence interval</b>	<b>P-value</b>
MR-Egger	1.003	0.991 to 1.016	0.609



**Table 2. Sensitivity analysis results for Mendelian randomization analyses after excluding 2 variants (rs10760673 and rs6420484) strongly associated with AMD risk.**

<b>Analysis</b>	<b>Causal effect (OR per D)</b>	<b>95% Confidence interval</b>	<b>P-value</b>
Inverse variance-weighted	1.085	1.033 to 1.140	0.001
Weighted median	1.045	0.982 to 1.111	0.164
MR-Egger (slope)	1.057	0.945 to 1.182	0.329
<b>Analysis</b>	<b>Intercept</b>	<b>95% Confidence interval</b>	<b>P-value</b>
MR-Egger (intercept)	1.003	0.992 to 1.014	0.608

**Table 3. Sensitivity analysis results for Mendelian randomization analyses after excluding 31 variants strongly associated with non-refractive error traits identified by Phenoscanner.**

<b>Analysis</b>	<b>Causal effect (OR per D)</b>	<b>95% Confidence interval</b>	<b>P-value</b>
Inverse variance-weighted	1.069	1.016 to 1.124	0.010
Weighted median	1.046	0.980 to 1.117	0.175
MR-Egger (slope)	1.060	0.949 to 1.182	0.301
<b>Analysis</b>	<b>Intercept</b>	<b>95% Confidence interval</b>	<b>P-value</b>
MR-Egger (intercept)	1.001	0.990 to 1.012	0.865

Figure 1  
[Click here to download high resolution image](#)





Figure 2

[Click here to download high resolution image](#)



**TOC Statement** (75 words)

This manuscript describes a Mendelian randomization analysis of publicly available genome wide association study data to test the hypothesis that refractive error causes an increased risk of age-related macular degeneration. The analysis suggested that a one diopter more hyperopic refractive error caused an 8-10% increased risk of age-related macular degeneration, assuming a linear relationship. Thus, past studies reporting large risks due to refractive error may have been subject to confounding bias.

**Supplementary Table 1. GWAS summary statistics for 126 genetic variants identified by the CREAM consortium and 23andMe.** Association with refractive error in n=95,505 UK Biobank participants (Tedja et al. 2018<sup>21</sup>) and association with AMD risk in 16,144 cases and 17,832 controls from the IAMDCG (Fritsche et al.<sup>22</sup>).

Variant rsID	Effect allele	Non-effect allele	Effect allele frequency	Association with refractive error (D)			Association with AMD (logOR)			Phenoscanter outlier	MR-PRESSO outlier
				BETA	SE	P	BETA	SE	P		
rs10100333	T	C	0.612	-0.056	0.012	6.60E-06	-0.038	0.017	3.12E-02	Yes	---
rs10122788	G	A	0.432	-0.036	0.012	3.80E-04	-0.027	0.017	1.19E-01	---	---
rs10187371	T	C	0.166	-0.04	0.016	3.00E-03	0.022	0.023	3.30E-01	---	---
rs10458138	A	G	0.218	-0.067	0.014	6.30E-07	-0.004	0.021	8.37E-01	---	---
rs10511652	G	A	0.592	-0.096	0.012	3.50E-18	-0.002	0.017	9.15E-01	---	---
rs1064583	G	A	0.398	-0.088	0.012	2.90E-14	-0.067	0.017	1.26E-04	Yes	---
rs10760673	A	G	0.202	-0.072	0.014	3.40E-07	-0.124	0.021	4.59E-09	Yes	Yes
rs10853531	G	A	0.786	-0.09	0.014	2.60E-10	-0.027	0.021	1.91E-01	Yes	---
rs10880855	T	C	0.504	-0.062	0.012	4.80E-08	0.023	0.017	1.86E-01	---	---
rs10887262	C	T	0.292	-0.103	0.013	2.80E-15	0.003	0.019	8.93E-01	---	---
rs10919908	A	G	0.387	-0.094	0.012	4.50E-17	0.003	0.018	8.52E-01	---	---
rs10948572	C	T	0.655	-0.067	0.012	2.90E-08	0.012	0.018	5.11E-01	---	---
rs11088317	T	C	0.295	-0.06	0.013	6.50E-06	0.01	0.019	5.82E-01	Yes	---
rs11101263	T	C	0.269	-0.094	0.013	2.20E-13	-0.063	0.019	1.09E-03	---	---
rs11118367	T	C	0.45	-0.087	0.012	1.20E-13	0.018	0.017	3.03E-01	Yes	---
rs11145465	A	C	0.217	-0.09	0.014	1.00E-10	0.016	0.021	4.34E-01	---	---
rs11178469	T	C	0.773	-0.049	0.014	2.60E-04	0.005	0.02	8.07E-01	---	---
rs11210537	G	A	0.683	-0.107	0.012	1.40E-17	-0.006	0.018	7.29E-01	---	---
rs1150687	C	T	0.395	-0.072	0.012	3.10E-10	0.021	0.017	2.19E-01	Yes	---
rs11589487	G	A	0.557	-0.075	0.012	2.20E-10	0.013	0.017	4.49E-01	---	---
rs11654644	T	C	0.188	-0.077	0.015	9.50E-08	0.025	0.022	2.59E-01	---	---
rs117735470	A	G	0.092	-0.092	0.02	5.20E-06	-0.06	0.029	4.00E-02	---	---
rs11802995	C	A	0.227	-0.086	0.014	4.40E-11	-0.015	0.02	4.52E-01	---	---
rs11952819	T	C	0.275	-0.032	0.013	4.00E-02	0.014	0.019	4.78E-01	---	---

rs12193446	A	G	0.904	-0.423	0.02	4.6E-106	-0.015	0.029	6.04E-01	---	---
rs1237670	G	A	0.224	-0.089	0.014	1.10E-10	-0.014	0.02	4.92E-01	---	---
rs12451582	G	A	0.64	-0.106	0.012	8.80E-18	-0.033	0.018	6.60E-02	---	---
rs12883788	C	T	0.539	-0.078	0.012	4.90E-12	-0.015	0.017	3.65E-01	Yes	---
rs12898755	G	A	0.791	-0.114	0.014	1.40E-16	-0.043	0.021	3.93E-02	---	---
rs12965607	G	T	0.152	-0.125	0.016	8.10E-16	-0.031	0.024	1.87E-01	---	---
rs1313240	C	T	0.289	-0.098	0.013	2.00E-15	-0.068	0.019	2.84E-04	Yes	---
rs1317537	T	C	0.516	-0.087	0.012	4.00E-14	0.024	0.017	1.65E-01	---	---
rs1358684	C	T	0.27	-0.04	0.013	5.50E-04	-0.01	0.019	6.19E-01	---	---
rs1359543	A	G	0.648	-0.058	0.012	2.10E-06	0.002	0.018	9.25E-01	---	---
rs1381566	G	T	0.19	-0.191	0.015	7.70E-40	-0.024	0.022	2.73E-01	---	---
rs14165	G	A	0.692	-0.067	0.013	4.80E-08	-0.01	0.018	5.94E-01	---	---
rs1454776	G	T	0.475	-0.058	0.012	2.00E-07	-0.032	0.017	6.27E-02	---	---
rs1532278	T	C	0.4	-0.048	0.012	3.10E-05	0.036	0.017	3.89E-02	Yes	---
rs1555075	C	T	0.649	-0.077	0.012	4.10E-11	-0.013	0.018	4.81E-01	Yes	---
rs1556867	T	C	0.239	-0.109	0.014	4.20E-17	-0.02	0.02	3.18E-01	---	---
rs1649068	A	C	0.449	-0.074	0.012	7.50E-11	-0.002	0.017	8.90E-01	Yes	---
rs17032696	A	C	0.814	-0.063	0.015	9.20E-05	-0.003	0.022	8.99E-01	---	---
rs17125093	A	G	0.198	-0.076	0.015	1.30E-08	-0.029	0.021	1.78E-01	---	---
rs17274750	C	A	0.098	-0.108	0.02	1.90E-08	-0.057	0.029	4.56E-02	---	---
rs17382981	T	C	0.424	-0.063	0.012	4.10E-07	-0.019	0.017	2.81E-01	Yes	---
rs1790165	C	A	0.576	-0.074	0.012	1.80E-10	-0.003	0.017	8.40E-01	---	---
rs1928175	A	G	0.561	-0.087	0.012	5.90E-15	-0.017	0.017	3.23E-01	Yes	---
rs1954761	T	C	0.371	-0.097	0.012	1.20E-16	0.018	0.018	3.13E-01	---	---
rs1969091	C	A	0.71	-0.073	0.013	3.10E-09	0.009	0.019	6.18E-01	---	---
rs2150458	G	A	0.575	-0.084	0.012	1.80E-13	-0.005	0.017	7.72E-01	---	---
rs2155413	A	C	0.47	-0.097	0.012	1.10E-17	0.028	0.017	1.07E-01	---	---
rs2166181	G	A	0.47	-0.1	0.012	5.80E-18	0.003	0.017	8.82E-01	---	---
rs2312538	G	A	0.712	-0.057	0.013	1.20E-06	-0.048	0.019	1.12E-02	---	---
rs235770	T	C	0.389	-0.074	0.012	4.80E-11	0.013	0.017	4.74E-01	Yes	---

rs2466574	G	A	0.292	-0.069	0.013	9.70E-09	0.035	0.019	6.19E-02	Yes	---
rs2573210	G	A	0.195	-0.194	0.015	7.90E-43	0.061	0.022	4.65E-03	---	Yes
rs2573232	T	C	0.908	-0.159	0.02	1.20E-16	0.015	0.03	6.07E-01	---	---
rs2622646	C	A	0.65	-0.083	0.012	2.30E-12	-0.019	0.018	2.77E-01	---	---
rs2761884	G	T	0.546	-0.101	0.012	2.00E-18	-0.003	0.017	8.48E-01	Yes	---
rs28471081	A	G	0.792	-0.112	0.014	3.70E-14	0.008	0.021	7.10E-01	---	---
rs284818	T	C	0.132	-0.093	0.017	1.60E-08	0.016	0.025	5.19E-01	---	---
rs28658452	A	G	0.916	-0.085	0.021	5.80E-05	0.034	0.031	2.64E-01	---	---
rs297593	T	C	0.288	-0.084	0.013	7.80E-11	-0.015	0.019	4.37E-01	---	---
rs3110134	G	A	0.687	-0.066	0.013	5.00E-08	-0.023	0.018	2.19E-01	---	---
rs3138137	C	A	0.514	-0.116	0.012	6.60E-25	-0.044	0.017	1.05E-02	---	---
rs34898258	T	C	0.44	-0.076	0.012	2.40E-10	-0.034	0.017	4.84E-02	---	---
rs35337422	C	A	0.146	-0.066	0.016	3.50E-05	0.066	0.024	6.45E-03	---	---
rs36024104	G	A	0.191	-0.098	0.015	2.20E-12	0.024	0.022	2.66E-01	---	---
rs41393947	A	G	0.141	-0.116	0.017	1.00E-12	-0.002	0.024	9.35E-01	Yes	---
rs4237285	C	T	0.535	-0.054	0.012	3.50E-06	-0.007	0.017	6.62E-01	---	---
rs4260345	C	T	0.373	-0.055	0.012	2.20E-06	-0.008	0.018	6.65E-01	---	---
rs4764038	T	G	0.267	-0.081	0.013	1.30E-09	-0.003	0.019	8.80E-01	---	---
rs4793501	T	C	0.579	-0.081	0.012	3.70E-12	-0.003	0.017	8.64E-01	---	---
rs4795364	G	A	0.252	-0.068	0.013	8.50E-08	-0.035	0.02	7.20E-02	Yes	---
rs4808962	G	A	0.169	-0.082	0.016	1.40E-08	0.032	0.023	1.64E-01	Yes	---
rs4894529	G	A	0.516	-0.037	0.012	2.00E-03	-0.032	0.017	6.47E-02	Yes	---
rs522774	A	C	0.741	-0.112	0.013	2.70E-18	0.009	0.019	6.42E-01	---	---
rs5442	A	G	0.07	-0.267	0.023	1.20E-33	-0.087	0.033	8.86E-03	---	---
rs55885222	A	C	0.365	-0.037	0.012	2.60E-03	0	0.018	9.99E-01	---	---
rs56014528	G	T	0.846	-0.093	0.016	9.10E-10	-0.006	0.024	8.06E-01	---	---
rs56055503	A	G	0.778	-0.061	0.014	8.00E-06	0.009	0.021	6.64E-01	---	---
rs56075542	T	G	0.549	-0.103	0.012	1.30E-18	-0.005	0.017	7.72E-01	---	---
rs61875120	C	T	0.22	-0.11	0.014	2.60E-16	-0.06	0.021	3.57E-03	Yes	---
rs62070229	G	A	0.187	-0.129	0.015	1.30E-18	0.01	0.022	6.53E-01	Yes	---



rs634990	C	T	0.488	-0.245	0.012	1.1E-103	-0.054	0.017	1.59E-03	---	---
rs6420484	A	G	0.356	-0.091	0.012	4.30E-15	0.117	0.018	4.11E-11	Yes	Yes
rs6433704	G	T	0.412	-0.121	0.012	5.60E-24	-0.028	0.017	1.03E-01	---	---
rs6495367	A	G	0.418	-0.149	0.012	7.20E-37	0.006	0.017	7.29E-01	Yes	---
rs6753137	T	C	0.432	-0.057	0.012	1.40E-06	-0.029	0.017	9.43E-02	---	---
rs7042950	G	A	0.224	-0.116	0.014	2.90E-18	-0.021	0.02	3.05E-01	---	---
rs7107014	C	A	0.516	-0.047	0.012	1.10E-04	0.004	0.017	8.21E-01	---	---
rs7122817	G	A	0.518	-0.072	0.012	1.10E-10	-0.009	0.017	5.81E-01	---	---
rs7143516	T	G	0.256	-0.101	0.013	3.70E-14	0.025	0.02	2.02E-01	---	---
rs7188859	C	T	0.366	-0.175	0.012	3.00E-49	-0.026	0.018	1.44E-01	---	---
rs7207217	A	G	0.382	-0.043	0.012	1.20E-03	0.008	0.018	6.48E-01	---	---
rs72655575	C	A	0.789	-0.07	0.014	7.10E-07	-0.028	0.021	1.82E-01	---	---
rs7337610	T	C	0.373	-0.071	0.012	7.70E-09	-0.045	0.018	1.01E-02	---	---
rs73730144	C	A	0.012	-0.311	0.053	7.00E-10	0.145	0.077	6.04E-02	---	---
rs7449443	T	G	0.602	-0.037	0.012	2.70E-04	0.001	0.017	9.37E-01	---	---
rs7554219	C	T	0.342	-0.097	0.012	1.10E-16	0.06	0.018	8.16E-04	---	Yes
rs7584258	A	G	0.752	-0.07	0.013	2.00E-07	-0.009	0.02	6.63E-01	Yes	---
rs7624084	T	C	0.557	-0.098	0.012	6.50E-17	-0.06	0.017	4.68E-04	Yes	---
rs7662551	G	A	0.255	-0.096	0.013	6.00E-12	-0.002	0.02	9.28E-01	---	---
rs7667446	C	T	0.184	-0.104	0.015	7.50E-13	-0.012	0.022	5.73E-01	---	---
rs7737179	A	G	0.224	-0.059	0.014	1.00E-05	-0.06	0.02	3.43E-03	Yes	---
rs7744813	A	C	0.586	-0.216	0.012	1.00E-75	0.004	0.017	8.26E-01	---	---
rs7747	C	T	0.802	-0.115	0.015	7.70E-16	0.012	0.021	5.90E-01	---	---
rs7829127	A	G	0.794	-0.135	0.014	3.10E-22	-0.011	0.021	6.13E-01	---	---
rs7895108	T	G	0.372	-0.126	0.012	1.10E-27	-0.003	0.018	8.48E-01	---	---
rs7933504	G	A	0.292	-0.05	0.013	1.40E-05	-0.035	0.019	6.25E-02	---	---
rs7941828	C	T	0.64	-0.044	0.012	1.90E-04	-0.015	0.018	3.86E-01	Yes	---
rs7968679	G	A	0.316	-0.076	0.013	4.20E-10	0.003	0.018	8.87E-01	---	---
rs7971334	T	G	0.303	-0.029	0.013	3.30E-02	-0.044	0.019	1.67E-02	---	---
rs8032307	C	T	0.874	-0.128	0.018	1.50E-11	-0.04	0.026	1.23E-01	---	---

rs8073754	T	C	0.174	-0.105	0.015	5.30E-13	0.002	0.023	9.12E-01	---	---
rs8076642	G	A	0.596	-0.055	0.012	8.60E-06	-0.022	0.017	2.10E-01	Yes	---
rs8137714	G	T	0.214	-0.04	0.014	2.20E-02	0.069	0.021	9.78E-04	---	---
rs837323	C	T	0.525	-0.091	0.012	5.30E-16	0.011	0.017	5.32E-01	---	---
rs9295499	C	A	0.68	-0.071	0.012	6.20E-09	-0.017	0.018	3.42E-01	---	---
rs9388766	C	T	0.697	-0.048	0.013	4.10E-05	0.038	0.019	3.81E-02	Yes	---
rs9416017	T	C	0.362	-0.057	0.012	6.00E-06	-0.013	0.018	4.69E-01	---	---
rs9516194	G	A	0.5	-0.044	0.012	2.60E-05	-0.017	0.017	3.19E-01	---	---
rs9517964	C	T	0.427	-0.108	0.012	3.40E-20	-0.009	0.017	5.89E-01	---	---
rs9547035	G	T	0.267	-0.095	0.013	1.60E-13	0.011	0.019	5.54E-01	---	---
rs9680365	A	G	0.034	-0.078	0.032	1.70E-02	-0.082	0.047	8.08E-02	---	---
rs9681162	C	T	0.289	-0.088	0.013	6.30E-13	-0.004	0.019	8.37E-01	---	---

