

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/124116/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Cardoso-Moreira, Margarida, Halbert, Jean, Valloton, Delphine, Velten, Britta, Chen, Chunyan, Shao, Yi, Liechti, Angélica, Ascensão, Kelly, Rummel, Coralie, Ovchinnikova, Svetlana, Mazin, Pavel V., Xenarios, Ioannis, Harshman, Keith, Mort, Matthew, Cooper, David N., Sandi, Carmen, Soares, Michael J., Ferreira, Paula G., Afonso, Sandra, Carneiro, Miguel, Turner, James M. A., VandeBerg, John L., Fallahshahroudi, Amir, Jensen, Per, Behr, Rüdiger, Lisgo, Steven, Lindsay, Susan, Khaitovich, Philipp, Huber, Wolfgang, Baker, Julie, Anders, Simon, Zhang, Yong E. and Kaessmann, Henrik 2019. Gene expression across mammalian organ development. *Nature* 571 , pp. 505-509. 10.1038/s41586-019-1338-5 file

Publishers page: <http://dx.doi.org/10.1038/s41586-019-1338-5> <<http://dx.doi.org/10.1038/s41586-019-1338-5>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Gene expression across mammalian organ development

Margarida Cardoso-Moreira^{1,2*}, Jean Halbert², Delphine Valloton², Britta Velten³, Chunyan Chen^{4,5,6}, Yi Shao^{4,5,6}, Angélica Liechti², Kelly Ascenção², Coralie Rummel², Svetlana Ovchinnikova¹, Pavel V. Mazin^{7,8,9}, Ioannis Xenarios², Keith Harshman², Matthew Mort¹⁰, David N. Cooper¹⁰, Carmen Sandi¹¹, Michael J. Soares^{12,13}, Paula G. Ferreira^{14,15}, Sandra Afonso¹⁶, Miguel Carneiro^{16,17}, James M. A. Turner¹⁸, John L. VandeBerg^{19,20}, Amir Fallahshahroudi²¹, Per Jensen²¹, Rüdiger Behr^{22,23}, Steven Lisgo²⁴, Susan Lindsay²⁴, Philipp Khaitovich^{7,25,26}, Wolfgang Huber³, Julie Baker²⁷, Simon Anders¹, Yong E. Zhang^{4,5,25} & Henrik Kaessmann^{1*}

- 1) Center for Molecular Biology of Heidelberg University (ZMBH), Heidelberg, Germany.
- 2) Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland.
- 3) Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany.
- 4) Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing, China.
- 5) State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing, China.
- 6) University of Chinese Academy of Sciences, Beijing, China.
- 7) Center for Neurobiology and Brain Restoration, Skolkovo Institute of Science and Technology, Moscow, Russia.
- 8) Institute for Information Transmission Problems (Kharkevich Institute) RAS, Moscow, Russia.
- 9) Faculty of Computer Science, HSE University, Moscow, Russia.
- 10) Institute of Medical Genetics, Cardiff University, Cardiff, UK.
- 11) Laboratory of Behavioral Genetics, Brain Mind Institute, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland.
- 12) Institute for Reproduction and Perinatal Research, Departments of Pathology and Laboratory Medicine and Pediatrics, University of Kansas Medical Center, Kansas City, MO, USA.
- 13) Center for Perinatal Research, Children's Research Institute, Children's Mercy, Kansas City, MO, USA.
- 14) Departamento de Anatomia, Universidade do Porto, Porto, Portugal.
- 15) ICBAS (Instituto de Ciências Biomédicas Abel Salazar), UMIB (Unidade Multidisciplinar de Investigação Biomédica), Universidade do Porto, Porto, Portugal.
- 16) CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Porto, Portugal.
- 17) Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, Porto, Portugal.
- 18) Sex Chromosome Biology Laboratory, The Francis Crick Institute, London, UK.
- 19) South Texas Diabetes and Obesity Institute, School of Medicine, The University of Texas Rio Grande Valley, Brownsville, Harlingen and Edinburg, TX, USA.
- 20) The Department of Human Genetics, School of Medicine, The University of Texas Rio Grande Valley, Brownsville, Harlingen and Edinburg, TX, USA.
- 21) AVIAN Behavioural Genomics and Physiology Group, IFM Biology, Linköping University, Linköping, Sweden.
- 22) Platform Degenerative Diseases, German Primate Center, Leibniz Institute for Primate Research (DPZ), Göttingen, Germany.
- 23) DZHK (German Center for Cardiovascular Research), Partner Site Göttingen, Göttingen, Germany.
- 24) Human Developmental Biology Resource, Institute of Genetic Medicine, Newcastle University, Newcastle, UK.
- 25) Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming, China.
- 26) CAS Key Laboratory of Computational Biology, CAS-MPG Partner Institute for Computational Biology, Shanghai Institutes for Biological Sciences, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai, China.
- 27) Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.

50 **The evolution of gene expression in mammalian organ development remains largely uncharacterized. Here**
51 **we report the transcriptomes of seven organs (cerebrum, cerebellum, heart, kidney, liver, ovary and testis)**
52 **across developmental time points from early organogenesis to adulthood for human, macaque, mouse,**
53 **rat, rabbit, opossum and chicken. Comparisons of gene expression patterns identified developmental stage**
54 **correspondences across species, and differences in the timing of key events during the development of the**
55 **gonads. We found that the breadth of gene expression and the extent of purifying selection gradually**
56 **decrease during development, whereas the amount of positive selection and expression of new genes**
57 **increase. We identified differences in the temporal trajectories of expression of individual genes across**
58 **species, with brain tissues showing the smallest percentage of trajectory changes, and the liver and testis**
59 **showing the largest. Our work provides a resource of developmental transcriptomes of seven organs**
60 **across seven species, and comparative analyses that characterize the development and evolution of**
61 **mammalian organs.**

62
63 Understanding the molecular evolution of mammalian phenotypic traits is a fundamental biological goal. To
64 achieve it, evolutionary studies need to be conducted within a developmental framework, both because adult
65 phenotypes are defined during development¹⁻³ and because evolutionary and developmental processes are
66 intertwined. von Baer noted in the 19th century that morphological differences between species increase as
67 development advances⁴ and evidence for it has accumulated^{4,5}. Understanding the molecular foundations of
68 these patterns will facilitate the identification of general principles underlying phenotypic evolution.

69
70 Here we provide a resource of bulk transcriptomes across developmental stages, covering multiple organs
71 from early organogenesis to adulthood in seven species, enabling direct comparisons of expression patterns
72 in organ development within and across mammals (<http://evodevoapp.kaessmannlab.org>). This resource
73 enabled us to analyse the evolution of gene expression within mammalian organs across developmental
74 stages.

75
76 **Organ developmental transcriptomes**
77 We generated gene expression time series (RNA-seq) for six mammals (human, rhesus macaque, mouse, rat,
78 rabbit, opossum) and a bird (red junglefowl, henceforth “chicken”). We sampled seven organs representing
79 the three germ layers: brain (forebrain/cerebrum) and cerebellum (hindbrain/cerebellum) (ectoderm); heart,
80 kidney, ovary and testis (mesoderm); and liver (endoderm) (Fig. 1a). The time series span from early
81 organogenesis to adulthood, plus senescence in primates. We sampled prenatal development at regular
82 intervals (e.g., daily in rodents, weekly in humans), and postnatally we sampled neonates, “infants”,
83 juveniles, and adults (Fig. 1). There are exceptions: we lack early prenatal data for rhesus and chicken, and
84 ovary data for rhesus and postnatal human development (Supplementary Table 1). Because marsupial organ
85 development occurs predominantly postnatally⁶, all sampled stages except for one were collected
86 postnatally. The dataset consists of 1,893 libraries (Supplementary Table 2).

87
88 The global relationships among all samples can be explored through a principle component analysis (PCA)
89 (Fig. 1b). The first principal component (PC1), explaining most variance in gene expression, separates the
90 samples by the germ layer from which organs originate. PC2 separates the samples by developmental stage
91 (from early to late development). PC3 and PC4 separate the samples by species (Extended Data Fig. 1a). In
92 PCAs of individual organs, PC1 and PC2 order the samples by developmental stage and separate them by
93 species (Extended Data Fig. 1b). In the global PCA (Fig. 1b), the earliest samples from different organs cluster
94 together, suggesting strong commonalities. We hypothesized that developmental programs are still largely
95 shared across organs at these early stages and found that organ transcriptomes are indeed most similar at
96 these stages (Extended Data Fig. 1c). Throughout development, the expression of transcription factors (TFs)

97 differs more between organs than that of the whole transcriptome (Extended Data Fig. 1c), consistent with
98 TFs directing organogenesis.

99
100 Next, we identified genes with significant temporal expression changes in each organ, termed
101 developmentally dynamic genes (DDGs; Extended Data Fig. 2a; Supplementary Tables 3-9; Methods). DDGs
102 reflect changes during development in gene regulation, cell type abundance, and/or the proportion of cells
103 undergoing division⁴. Consistently, between 79-91% of protein-coding genes in each species are DDGs
104 (Extended Data Fig. 2b), including genes with housekeeping functions. DDGs are enriched with phenotypes
105 associated with the development, anatomy and physiology of each organ, plus general growth and patterning
106 (FDR < 0.01, hypergeometric test; Supplementary Tables 10-11). DDGs are under stronger functional
107 constraints⁷⁻¹⁰ than non-DDGs, and the constraints increase with the number of organs in which genes show
108 temporal dynamic expression (Extended Data Fig. 2c). The increased constraints extend to dosage changes,
109 with DDGs being less tolerant to duplication and deletion variants¹¹ (Extended Data Fig. 2d).

110
111 In each species, 6-15% of the genes (Extended Data Fig. 2b) are DDGs in only one organ, and are consistently
112 enriched with organ-specific phenotypes (FDR < 0.01, hypergeometric test; Supplementary Table 12). The
113 fraction of expressed organ-specific DDGs increases during development (Extended Data Fig. 2e), correlating
114 with organ differentiation and maturation. The opposite is observed for TFs, whose contribution is highest
115 earlier in development (Extended Data Fig. 2f).

116

117 **Developmental correspondences and heterochrony**

118 Embryonic development is divided into 23 Carnegie stages, which are matched across species¹²⁻¹⁵ (Extended
119 Data Fig. 3a). However, cross-species correspondences during fetal and postnatal development are unknown.
120 Therefore, we used the developmental transcriptomes to establish stage correspondences across species
121 throughout the entire development (Methods; Fig. 2a; Extended Data Fig. 3b). We recapitulated the Carnegie
122 stage correspondences (rabbit is shifted 1-2 days; Methods; Extended Data Fig. 3a) and found that gene
123 expression in a newborn opossum is closest to a mouse at e11.5, matching previous estimates¹⁶.

124

125 Organ development includes periods of greater transcriptional change¹⁷. We identified and characterized
126 these periods across species using our stage correspondences. These periods occur at similar stages across
127 species and are enriched with orthologous genes ($P \leq 0.001$, hypergeometric test; Fig. 2b; Extended Data
128 Figs. 4, 5). In somatic organs, there are two main periods of transcriptional change. The first occurs during
129 embryonic development and is defined by an increase in expression of genes with early organ-specific
130 functions and a decrease in expression of genes involved in cell division and general morphogenesis (Fig. 2b;
131 Extended Data Figs. 4, 5; Supplementary Table 13). The second occurs either postnatally or around birth,
132 depending on the maturity of the newborns of the different species. Mammals exhibit great variability in
133 their level of independence at birth, being classified as altricial (born less mature) or precocial (more
134 mature)⁶. These classifications are recapitulated by the developmental transcriptomes, with the altricial
135 newborn opossum at one end of the maturation spectrum and the precocial rhesus macaque at the other
136 (Fig. 2a). This second period of greater transcriptional change is defined by an increase in expression of genes
137 with late organ-specific functions and, again, by a decrease in expression of genes involved in cell division
138 and morphogenesis (Fig. 2b; Extended Data Figs. 4, 5; Supplementary Table 13). Thus, whereas in altricial
139 species this period of intense organ maturation occurs postnatally, in precocial species it overlaps with birth.

140

141 Developmental programs can change through shifts in the timing of events, i.e. "heterochrony"¹¹. If the
142 development of an organ were to be shifted in one species, the developmental correspondences for that

143 organ would be different from the global correspondences. Overall, organ-specific correspondences are
144 consistent with the global correspondences, except for early heart development in opossum and early ovary
145 development in human and rabbit (Extended Data Fig. 6; Methods).

146
147 However, heterochronies do not have to involve whole organs, they can occur in developmental modules
148 within organs. Indeed, heterochronies occur during the production of gametes¹⁸, a process dependent on
149 meiosis. *Stra8* is the gatekeeper for germ cell entry into meiosis and its role is conserved across
150 vertebrates^{3,19}. Therefore, we analyzed the expression of *Stra8* and other meiotic genes to identify the start
151 of meiosis in each species, and identify differences in its timing across species (Fig. 2a; Extended Data Figs.
152 7a-d). During oogenesis, meiotic genes are expressed as early as during embryonic development (mouse), at
153 the boundary between embryonic and fetal development (rat and human), or during late fetal development
154 (rabbit) (Fig. 2a; Extended Data Figs. 6, 7a-b). Although less frequent, we also identified heterochronies in
155 the onset of meiosis during spermatogenesis (Fig. 2a; Extended Data Fig. 7c-d). In spermatogenesis the onset
156 of meiosis marks the beginning of dramatic changes in cellular composition²⁰, which sharply change testis
157 transcriptomes (Extended Data Fig. 7e). Starting at birth in rodents and later in rabbit there are also profound
158 changes in ovary transcriptomes (Extended Data Fig. 7e), coinciding with the breakdown of germ cell nests
159 and follicle assembly²¹. Heterochronies are therefore abundant during mammalian gonadal development,
160 representing another mechanism underlying the extreme variability of gonadal morphogenesis³.

161

162 **Relationships between evolution and development**

163 After the phylotypic period, the most conserved embryonic stage, morphological differences between
164 species increase as development progresses — von Baer's divergence^{4,5} (Extended Data Fig. 8a). Previous
165 studies assessed the relationship between development and molecular divergence for whole embryos and
166 found that molecular divergence increases as development progresses²²⁻²⁵. We recapitulated this
167 observation for individual organs, consistently finding transcriptome correlations between species to decline
168 with developmental time (Extended Data Fig. 8b).

169

170 Two non-mutually-exclusive hypotheses can account for the increasing molecular and morphological
171 divergence during development²⁶. First, early development could be under greater functional constraints and
172 be more refractory to change. Second, divergence could be driven by adaptive changes, which are more likely
173 to occur later in development, when the influence of the environment is stronger²⁶. To assess potential
174 differences in functional constraints across development, we compared the tolerance to functional
175 mutations of genes employed in early versus late development. Across all organs, genes employed early in
176 development are less tolerant to loss-of-function mutations ($P < 10^{-10}$, Wilcoxon rank sum test, two-sided;
177 Fig. 3a; Extended Data Fig. 8c). Consistently, using a set of neutrally ascertained mouse knockouts²⁷, we
178 observed for all organs that the percentage of expressed genes associated with lethality decreases during
179 development (Fig. 3b; Extended Data Fig. 8d). Both observations suggest that early development is subject
180 to stronger functional constraints. Next, we evaluated the relationship between adaptation and development
181 by examining the temporal expression of genes identified as carrying signatures of positive selection²⁸.
182 Organs differ in the proportion of positively selected genes: it is highest in liver and testis and lowest in brain
183 tissues (Fig. 3c). However, across all organs, the fraction of expressed positively selected genes increases
184 during development (Fig. 3c). Thus, an increase in positive selection likely also contributes to the progressive
185 molecular and morphological divergence of organs during development.

186

187 Organ transcriptomes can also diverge between species due to species-specific genes²⁹. Therefore, we
188 investigated how the contribution of recent gene duplications changes throughout development. For each
189 stage, we calculated an index combining the phylogenetic age of genes with their expression, where higher

190 values correspond to younger transcriptomes (Methods). We identified differences between organs similar
191 to those observed for positively selected genes: liver has the youngest developmental transcriptomes, brain
192 tissues the oldest (Fig. 3d). However, across organs, transcriptomes become younger during development,
193 indicating that gene duplications play increasingly more prominent roles (Fig. 3d).

194

195 Together, these analyses suggest that the increase in morphological and molecular divergence observed
196 between species during development is driven by a decrease in functional constraints as development
197 advances (Figs. 3a-b), and by a concurrent increase in positive selection (Fig. 3c) and addition of new genes
198 (Fig. 3d).

199

200 **Pleiotropy and the evolution of development**

201 The breadth of expression across tissues and timepoints (which we refer to here as pleiotropy) has an
202 influence on gene evolution^{30,31}. Therefore, we calculated the tissue- and time-specificity of each gene across
203 development (Extended Data Fig. 9a; Methods; Supplementary Tables 3-9). Time- and tissue-specificity are
204 highly correlated: tissue-specific genes are more likely to be expressed in narrower time windows and *vice*
205 *versa* (Pearson correlation coefficients, r : 0.63-0.89, $P < 10^{-15}$). Genes also tend to have similar temporal
206 breadths across organs (r : 0.48-0.92, $P < 10^{-15}$). As described^{32,33}, pleiotropy correlates with levels of
207 functional constraint: the more broadly expressed, the more intolerant genes are to functional variation (r =
208 0.29, $P < 10^{-15}$; Extended Data Fig. 9b). Consistently, genes associated with lethality²⁷ are more broadly
209 expressed than genes associated with subviability, which in turn are more broadly expressed than genes
210 associated with viability (all $P \leq 2 \times 10^{-8}$, Wilcoxon rank sum test, two-sided; Fig. 3e; Extended Data Fig. 9c).
211 This contrast with measures of tolerance to functional mutations, which distinguish genes associated with
212 lethality or subviability from viability ($P = 2 \times 10^{-12}$), but do not differentiate between lethality and subviability
213 (Fig. 3e; Extended Data Fig. 9d). Expression pleiotropy is therefore uniquely correlated with phenotypic
214 impact.

215

216 Pleiotropy has been forwarded as an explanation for the conservation of the phylotypic period^{24,34} and is a
217 determinant of the types of mutations that are permissible under selection^{30,31}. Therefore, we tested for
218 differences in pleiotropy between genes employed early versus late in development and found that genes
219 employed earlier have broader spatial and temporal expression than genes employed later (all $P < 10^{-6}$,
220 Wilcoxon rank sum test, two-sided; Fig. 3f; Extended Data Fig. 9e). Because a decrease in pleiotropy can
221 explain both a decrease in functional constraints and an increase in adaptation^{30,31}, we suggest that it may
222 be a major contributor to the increase in morphological and molecular divergence observed between species
223 during development.

224

225 **Evolution of developmental trajectories**

226 Differences between species in organ development are often correlated with changes in gene expression.
227 Consequently, we sought to identify genes that evolved new developmental trajectories. Hence, we
228 compared, within a phylogenetic framework, the temporal profiles of orthologous DDGs and identified those
229 with trajectory changes between species (Fig. 4a, b; Supplementary Tables 14-18; Methods).

230

231 Brain exhibits the smallest percentage of trajectory changes (11% DDGs), liver and testis the highest (23%
232 and 27%, respectively; Extended Data Fig. 10a). These organ differences are consistent with those observed
233 for positively selected genes and for gene duplications. Thus, across all evaluated mechanisms of
234 evolutionary change, the brain is the slowest evolving organ, whereas liver and testis are the fastest.

235

236 In mouse, rat and rabbit the distribution of trajectory changes among organs is similar (Extended Data Fig.
237 10b). Compared to these species, humans display an excess of trajectory changes in brain (20% changes in
238 human vs. 12-13% in others; $P = 1 \times 10^{-5}$, binomial test) and cerebellum (26% in human vs. 21-22% in others;
239 $P = 0.02$), and a paucity in testis (21% in human vs. 34-37% in others; $P = 1 \times 10^{-10}$) (Extended Data Fig. 10b).
240 Although it is tempting to invoke adaptation, the excess of changes in the human brain tissues could partly
241 stem from differences in sampling (Methods). Overall, rodents evolved a higher number of trajectory changes
242 when compared to human and rabbit ($P < 10^{-10}$; Fig. 4b).

243
244 Orthologs tested for trajectory changes are more pleiotropic than the full set of genes in each species, which
245 also includes gene duplications/losses (all $P < 10^{-12}$, Wilcoxon rank sum test, two-sided; Extended Data Fig.
246 10c). However, among those tested, genes with new trajectories are as pleiotropic as genes with conserved
247 trajectories (Extended Data Fig. 10d). Importantly, while genes with trajectory changes are broadly
248 expressed, the changes themselves are organ-specific (Extended Data Fig. 10e). Trajectory changes are
249 restricted to one organ in 93-96% of the cases. This is consistent with the underlying mutations affecting
250 regulatory elements, which control a subset of the total spatiotemporal profile of each gene, and with
251 evolutionary theory, as mutations affecting multiple organs are less likely to fix in populations^{30,31}. However,
252 not all trajectory changes are directly due to regulatory mutations; they can also be caused by changes in
253 cellular composition.

254

255 Discussion

256 We profiled the development of seven major organs, from early organogenesis to adulthood, across multiple
257 mammals, to create an extensive resource (evodevoapp.kaessmannlab.org). We used developmental
258 transcriptomes to match stages across species and identified extensive heterochronies during gonadal
259 development. We found the evolution of mammalian organs to be inextricably linked to their development.
260 Organs are most similar between species early in development and then become increasingly more distinct,
261 which is likely explained by changes in pleiotropy during development. Early in development, active genes
262 tend to function in multiple organs and stages, rendering them more refractory to change. As organs
263 differentiate and mature, active genes have more restricted spatiotemporal profiles, which may reduce
264 functional constraints and facilitate evolutionary change. The increase in species divergence as development
265 progresses has also been described for mammalian limb development³⁵ and whole embryos⁴⁻⁵ and we
266 suggest it occurs in developmental systems where pleiotropy decreases as a function of time.

267

268 A next challenge will be to identify the molecular drivers of the new developmental trajectories, which may
269 underlie the evolution of organ phenotypes. It will be important to assess the extent to which these trajectory
270 changes are caused by changes in gene regulation and/or cellular composition. This can be accomplished by
271 complementing the data and results of this study with single-cell transcriptomic and epigenomics datasets
272 and bioinformatic deconvolution procedures. Such endeavors will further advance our understanding of the
273 genetic and developmental foundations of the evolution of mammalian phenotypes.

274

275 Acknowledgments

276 We thank R. Arguello, F. Lamanna, I. Sarropoulos, M. Sepp, and members of the Kaessmann group for
277 discussion; I. Moreira for developing the Shiny application; M. Sanchez-Delgado and N. Trost for figure
278 assistance; P. Dhakal for assistance collecting rat samples; S. Nef for assistance collecting gonadal samples;
279 the Lausanne Genomics Technology Facility for high-throughput sequencing; the Joint MRC/Wellcome
280 (MR/R006237/1) Human Developmental Biology Resource (HDBR), the Maryland Brain Collection Center and
281 the Chinese Brain Bank Center for providing human samples; and the Suzhou Experimental Animal Center for
282 providing rhesus samples. Computations were performed at the Vital-IT Center of the SIB Swiss Institute of

283 Bioinformatics and at the bwForCluster from the Heidelberg University Computational Center (supported by
284 the state of Baden-Württemberg through bwHPC and the German Research Foundation - INST 35/1134-1
285 FUGG). M.J.S. was supported by the National Institutes of Health (HD020676); P.J. by the European Research
286 Council (322206, Genewell); Y.E.Z. by the National Natural Science Foundation of China (31771410,
287 91731302); M.C. by Fundação para a Ciência e Tecnologia through POPH-QREN funds from the European
288 Social Fund and Portuguese MCTES (IF/00283/2014/CP1256/CT0012). This research was supported by grants
289 from the European Research Council (615253, OntoTransEvol) and Swiss National Science Foundation
290 (146474) to H.K. and by the Marie Curie FP7-PEOPLE-2012-IIF to M.C.M. (329902).

291

292 **Author contributions**

293 M.C.M. and H.K. conceived and organized the study based on H.K.'s original design. M.C.M. performed all
294 analyses. M.C.M. and H.K. wrote the manuscript, with input from all authors. J.H., D.V., C.S., M.J.S., P.G.F.,
295 S.Afonso, M.C., J.M.T, J.L.V., A.F., P.J., R.B., S.Lisgo, S.Lindsay, P.K. and J.B. collected samples. J.H., D.V., A.L.,
296 K.A. and C.R performed RNA-seq experiments. B.V. and W.H. contributed to analyses on trajectory changes.
297 S.O., S.Anders and P.V.M. contributed to analyses on developmental correspondences. C.C., Y.S. and Y.E.Z.
298 contributed to analyses on transcriptome age. M.M. and D.N.C. contributed to data interpretation. I.X.
299 organized high-performance computational resources. K.H. organized high-throughput sequencing.

300

301 **Competing interests**

302 The authors declare no competing financial interests.

303

304 **Author information**

305 Correspondence and requests for materials should be addressed to M.C.M. ([m.moreira@zmbh.uni-
306 heidelberg.de](mailto:m.moreira@zmbh.uni-heidelberg.de)) or H.K. (h.kaessmann@zmbh.uni-heidelberg.de).

307

308 **References**

- 309 1. Pantalacci, S. & Semon, M. Transcriptomics of Developing Embryos and Organs: A Raising Tool for
310 Evo–Devo. *J. Exp. Zool.* **00B**, 1-9 (2014).
- 311 2. Silbereis, J. C., Pochareddy, S., Zhu, Y., Li, M. & Sestan, N. The Cellular and Molecular Landscapes of
312 the Developing Human Central Nervous System. *Neuron* **89**, 248-268 (2016).
- 313 3. DeFalco, T. & Capel, B. Gonad morphogenesis in vertebrates: divergent means to a convergent end.
314 *Annu. Rev. Cell. Dev. Biol.* **25**, 457-482 (2009).
- 315 4. Abzhanov, A. von Baer's law for the ages: lost and found principles of developmental evolution.
316 *Trends Genet.* **29**, 712-722 (2013).
- 317 5. Kalinka, A. T. & Tomancak, P. The evolution of early animal embryos: conservation or divergence?
318 *Trends Ecol. Evol.* **27**, 385-393 (2012).
- 319 6. Ferner, K., Schultz, J. A. & Zeller, U. Comparative anatomy of neonates of the three major mammalian
320 groups (monotremes, marsupials, placentals) and implications for the ancestral mammalian neonate
321 morphotype. *J. Anat.* **231**, 798-822 (2017).
- 322 7. Dickinson, M. E. et al. High-throughput discovery of novel developmental phenotypes. *Nature* **537**,
323 508-514 (2016).
- 324 8. Petrovski, S., Wang, Q., Heinzen, E.L., Allen, A. S. & Goldstein, D. B. Genic intolerance to functional
325 variation and the interpretation of personal genomes. *PLoS Genet.* **9**, e1003709 (2013).
- 326 9. Lek, M. et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285-91
327 (2016).
- 328 10. Cassa, C. A. et al. Estimating the selective effects of heterozygous protein-truncating variants from
329 human exome data. *Nat. Genet.* **49**, 806-810 (2017).
- 330 11. Ruderfer, D.M. et al. Patterns of genic intolerance of rare copy number variation in 59,898 human
331 exomes. *Nat. Genet.* **48**, 1107-1111 (2016).

- 332 12. Hill, M.A. Embryology Carnegie Stage Comparison. https://jeltsch.org/carnegie_stage_comparison
333 (2017).
- 334 13. de Bakker, B.S. et al. An interactive three-dimensional digital atlas and quantitative database of
335 human development. *Science* **354** (2016).
- 336 14. Kerwin, J. et al. The HUDSEN Atlas: a three-dimensional (3D) spatial framework for studying gene
337 expression in the developing human brain. *J. Anat.* **217**, 289-299 (2010).
- 338 15. Butler, H. & Juurlink, B. H. J. *An Atlas for Staging Mammalian and Chick Embryos*. (CRC Press, Boca
339 Raton, Fla., 1987).
- 340 16. Smith, K. K. Early development of the neural plate, neural crest and facial region of marsupials. *J.*
341 *Anat.* **199**, 121-131 (2001).
- 342 17. Dillman, A. A. et al. mRNA expression, splicing and editing in the embryonic and adult mouse cerebral
343 cortex. *Nat. Neurosci.* **16**, 499-506 (2013).
- 344 18. Glucksmann, A. Sexual dimorphism in mammals. *Biol. Rev.* **49**, 423-475 (1974).
- 345 19. Feng, C. W., Bowles, J. & Koopman, P. Control of mammalian germ cell entry into meiosis. *Mol. Cell.*
346 *Endocrinol.* **382**, 488-497 (2014).
- 347 20. Soumillon, M. et al. Cellular source and mechanisms of high transcriptome complexity in the
348 mammalian testis. *Cell Rep.* **3**, 2179-2190 (2013).
- 349 21. Ungewitter, E. K. & Yao, H. H. How to make a gonad: cellular mechanisms governing formation of the
350 testes and ovaries. *Sex. Dev.* **7**, 7-20 (2013).
- 351 22. Roux, J. & Robinson-Rechavi, M. Developmental Constraints on Vertebrate Genome Evolution. *PLoS*
352 *Genet.* **4**, 10 (2008).
- 353 23. Kalinka, A. T. et al. Gene expression divergence recapitulates the developmental hourglass model.
354 *Nature* **468**, 811-814 (2010).
- 355 24. Hu, H. et al. Constrained vertebrate evolution by pleiotropic genes. *Nat. Ecol. Evol.* **1**, 1722-1730
356 (2017).
- 357 25. Hazkani-Covo, E., Wool, D. & Graur, D. In search of the vertebrate phylotypic stage: a molecular
358 examination of the developmental hourglass model and von Baer's third law. *J. Exp. Zool. B. Mol.*
359 *Dev. Evol.* **304**, 150-8 (2005).
- 360 26. Garfield, D. A. & Wray, G. A. Comparative embryology without a microscope: using genomic
361 approaches to understand the evolution of development. *J. Biol.* **8**, 65 (2009).
- 362 27. Koscielny, G. et al. The International Mouse Phenotyping Consortium Web Portal, a unified point of
363 access for knockout mice and related phenotyping data. *Nucleic Acids Res.* **42**, D802-9 (2014).
- 364 28. Kosiol, C. et al. Patterns of Positive Selection in Six Mammalian Genomes. *PLoS Genet.* **4**, e1000144
365 (2008).
- 366 29. Kaessmann, H. Origins, evolution, and phenotypic impact of new genes. *Genome Res.* **20**, 1313-1326
367 (2010).
- 368 30. Stern, D. L. Evolutionary developmental biology and the problem of variation. *Evolution* **54**, 1079-91
369 (2000).
- 370 31. Carroll, S. B. Evolution at two levels: on genes and form. *PLoS Biol* **3**, e245 (2005).
- 371 32. Duret, L. & Mouchiroud, D. Determinants of substitution rates in mammalian genes: expression
372 pattern affects selection intensity but not mutation rate. *Mol. Biol. Evol.* **17**, 68-74 (2000).
- 373 33. Winter, E., Goodstadt, L. & Ponting, C. P. Elevated Rates of Protein Secretion, Evolution, and Disease
374 Among Tissue-Specific Genes. *Genome Res.* **14**, 54-61 (2004).
- 375 34. Galis, F. & Metz, J. Testing the Vulnerability of the Phylotypic Stage: On Modularity and Evolutionary
376 Conservation. *Journal of Experimental Zoology (Mol Dev Evol)* **291**, 195-204 (2001).
- 377 35. Sears, K., Maier, J. A., Sadier, A., Sorensen, D. & Urban, D. J. Timing the developmental origins of
378 mammalian limb diversity. *Genesis* **56** (2018).
- 379 36. Plant, T. M., Ramaswamy, S., Simorangkir, D. & Marshall, G. R. Postnatal and pubertal development
380 of the rhesus monkey (*Macaca mulatta*) testis. *Ann. N. Y. Acad. Sci.* **1061**, 149-162 (2005).
- 381
- 382

Figure Legends

383

384 **Figure 1 | Organ developmental transcriptomes. a**, Species, organs, and stages sampled. Red slashes
385 highlight two sampling gaps: human development is not covered between 20 and 38 weeks post-conception
386 (wpc) and rhesus development between embryonic (e) day 130 and e160. **b**, PCA based on 7,696 1:1
387 orthologs across all species. Each dot represents the median across replicates (~2-4).

388

389 **Figure 2 | Developmental correspondences. a**, Stage correspondences across species. Grey bars represent
390 additionally sampled stages. Red shading highlights sampling gaps. In rhesus, male meiosis starts at 3-4
391 years³⁶. Because we did not detect expression of meiotic genes in the 3-year-olds, we placed meiosis' onset
392 between 3 and 9 years. **c**, Number of genes differentially expressed between adjacent, species-matched,
393 stages for brain and liver (\log_2 fold-change ≥ 0.5 ; other organs in Extended Data Fig. 5). Solid lines mark genes
394 that increase in expression and dashed lines genes that decrease. Vertical dotted line marks birth.

395

396 **Figure 3 | Relationships between evolution and development. a**, Intolerance to functional mutations (pLI)
397 for human genes whose expression decreases (blue) or increases (orange) during development (4,589/4,478
398 genes decrease/increase in brain, 2,673/3,442 in heart, and 2,290/3,794 in testis; all $P < 10^{-10}$, Wilcoxon rank
399 sum test, two-sided). **b**, Percentage of lethal genes expressed at each stage (out of 2,676 knockouts).
400 Weighted average Spearman correlation coefficient is -0.89 ($P = 1 \times 10^{-12}$); all organ-specific Spearman
401 correlations are significant ($P \leq 0.04$). **c**, Percentage of positively-selected genes expressed at each stage (out
402 of 13,362 genes tested for positive selection). Ovary excluded due to lack of postnatal data. Weighted
403 average Spearman correlation coefficient is $+0.57$ ($P = 5 \times 10^{-11}$); all organ-specific correlations are significant
404 ($P \leq 0.05$). **d**, Phylogenetic age of organs' transcriptomes throughout development for rat ($n = 18,695$ genes),
405 human ($n = 18,820$) and chicken ($n = 15,155$). Higher values indicate larger contributions of lineage-specific
406 genes (i.e. younger transcriptomes). Weighted average Spearman correlation coefficients are $+0.87$ ($P = 1 \times$
407 10^{-12}) for rat, $+0.77$ ($P = 1 \times 10^{-12}$) for human and $+0.96$ ($P = 1 \times 10^{-12}$) for chicken. All Spearman correlations
408 are significant except for rat brain and cerebellum (ρ : $+0.53$ to $+0.99$, P : 0.03 to 10^{-15}). Testis plotted
409 separately because of the young age of sexually mature transcriptomes. **e**, Tissue-specificity, time-specificity
410 (median across organs) and intolerance to functional mutations (pLI) of human orthologs of mouse genes
411 identified as lethal, subviable and viable ($n = 2,686$; Wilcoxon rank sum test, two sided; 'N.S.' means non-
412 significant). Box plots depict the median \pm 25th and 75th percentiles, whiskers at 1.5 times the interquartile
413 range. **f**, Tissue-specificity for mouse genes whose expression decreases (blue) or increases (orange) during
414 development (3,981/5,164 genes decrease/increase in brain, 4,631/5,051 in kidney, and 4,270/4,026 in liver;
415 all $P < 10^{-15}$, Wilcoxon rank sum test, two-sided). In **b-d**, the x-axes show samples ordered by stage (Fig. 1a).
416 In **b-c**, lines were estimated through linear regression; in **d** through loess. In **b-d** the 95% confidence interval
417 is shown in grey.

418

419 **Figure 4 | Evolution of developmental trajectories. a**, Example of two genes that evolved new trajectories
420 in the human cerebellum. *GRIA3* is a glutamate receptor associated with mental retardation. *MDGA1*
421 encodes a cell surface glycoprotein important for the developing nervous system. **b**, Pie charts depict the
422 number of genes in each organ that evolved new trajectories in each phylogenetic branch (3,980 genes tested
423 in brain, 3,064 in cerebellum, 1,871 in heart, 2,284 in liver and 3,191 in testis). Bar charts depict the total
424 number of trajectory changes in each species. For mouse, that meant adding the changes that occurred at
425 the base of the glires (I), those shared by mouse and rat (II) and those that are mouse specific (III). $**P < 10^{-10}$,
426 binomial test.

427

428 **Online Methods**

429

430 **Ethics statement**

431 The human prenatal samples were provided by the MRC-Wellcome Trust Human Developmental Biology
432 Resource (HDBR) and are from elective abortions with normal karyotypes. The tissue donations were made
433 entirely voluntarily by women undergoing termination of pregnancy. Donors were asked to give explicit
434 written consent for the fetal material to be collected, and only after they had been counselled about the
435 termination of their pregnancy. The human postnatal samples were provided by the NICHD Brain and Tissue
436 Bank for Developmental Disorders at the University of Maryland (USA) and by the Chinese Brain Bank Center
437 (CBBC) in Wuhan (China). They originated from individuals with diverse causes of death that, given the
438 information available, were not associated with the organ sampled. Written consent for the use of human
439 tissues for research was obtained from all donors or their next of kin by the respective tissue banks. The
440 rhesus macaque samples were provided by the Suzhou Experimental Animal Center (China). All rhesus
441 macaques used in this study suffered sudden deaths for reasons other than their participation in this study
442 and without any relation to the organ sampled. The use of all samples for the work described in this study
443 was approved by an ERC Ethics Screening panel (associated with H.K.'s ERC Consolidator Grant 615253,
444 OntoTransEvol) and local ethics committees in Lausanne (authorization 504/12) and Heidelberg
445 (authorization S-220/2017).

446

447 **Human developmental samples**

448 We started sampling human prenatal development at 4 weeks post-conception (wpc) and then sampled the
449 developing organs each week until 20 wpc (except for 14, 15 and 17 wpc). There are no samples available
450 between 20 and 38 wpc. Postnatally we sampled neonates, “infants” (6-9 months), “toddlers” (2-4 years),
451 “school” (7-9 years), “teenagers” (13-19 years), and then adults from each decade until 63 years of age
452 (Supplementary Tables 1-2). Human ovary development was only sampled prenatally (until 18wpc) and
453 kidney development was sampled up until (and including) 8 years of age (“school”). Prenatally, we considered
454 as biological replicates individuals from the same developmental week. Hence, for example, individuals from
455 Carnegie stages 13 and 14 were considered to be replicates (i.e. 4 wpc) even though they were not at the
456 same developmental stage according to phenotypic milestones. Supplementary Table 2 provides the precise
457 age of the donors. The number of biological replicates ranges from 1 to 4 (median of 2), for a total of 297
458 RNA-seq libraries.

459

460 **Other species developmental samples**

461 In mouse (*Mus musculus*, outbred strain CD-1 - RjOrl:SWISS), we started sampling prenatal development at
462 e10.5 and then collected samples daily until birth (i.e. until e18.5). Postnatally we sampled individuals at 5
463 stages: P0, P3, P14, P28 and P63. There are 4 replicates (2 males and 2 females) per stage, except for ovary
464 and testis where we aimed for 2 replicates, for a total of 316 RNA-seq libraries.

465

466 In rat (*Rattus norvegicus*, outbred strain Holtzman SD), the time-series start at e11 and cover prenatal
467 development daily until birth (i.e. until e20). Postnatally we sampled individuals at 6 stages: P0, P3, P7, P14,
468 P42 and P112. We generated replicates as described for mouse, for a total of 350 RNA-seq libraries.

469

470 In rabbit (*Oryctolagus cuniculus*, outbred New Zealand breed), we started sampling prenatal development at
471 e12 and then sampled 11 time points up until (and including) e27 (gestation length is ~29-32 days).
472 Postnatally we sampled individuals at 4 stages: P0, P14, P84 and between P186-P548. The number of
473 replicates is the same as described for the rodents, for a total of 315 RNA-seq libraries.

474

475 For rhesus macaque (*Macaca mulatta*), the sample collection started at a fetal stage (e93) and we sampled
476 5 stages before birth (until e130; gestation lasts ~167 days). Postnatally we sampled 8 stages selected to

477 match the human time-series (Supplementary Tables 1-2). The number of replicates ranges from 1 to 4
478 (median of 2), for a total of 168 RNA-seq libraries. Ovary development was not sampled.

479
480 Because opossums (*Monodelphis domestica*) are born in a very immature state⁶, the stages of organ
481 development sampled in the other species during prenatal development occur postnatally in the marsupial.
482 Accordingly, we sampled one prenatal stage (e13.5; gestation length is ~14-15 days) and then sampled 14
483 postnatal stages, more densely right after birth and then at increasingly longer intervals (Supplementary
484 Table 1). There is a median of 3 biological replicates per stage for somatic organs and 2 for the gonads, for a
485 total of 232 RNA-seq libraries.

486
487 In chicken (*Gallus gallus*, the red junglefowl, progenitor of domestic chicken), we started sampling organ
488 development at a fetal stage (e10) and sampled 3 additional stages until e17 (egg incubation lasts ~21 days).
489 We then sampled postnatal development at 5 stages: P0, P7, P35, P70 and P155. There is a median of 4
490 biological replicates per stage for somatic organs (2 males and 2 females) and 2 for the gonads, for a total of
491 215 RNA-seq libraries.

492
493 This resource consists of 1,893 libraries, covering the development of 7 organs, 9-23 developmental stages
494 (depending on the species) and a median of 2-4 replicates per stage (full details in Supplementary Tables 1-
495 2).

496 497 **Organ dissections**

498 Mammalian embryos are morphologically similar⁴, and this similarity extends to the internal organs. Early in
499 development, the only clear morphological difference between the organs of the different species, when
500 present, is size. We started collecting samples when the organs could be dissected and isolated from nearby
501 tissues. For the brain this was possible across the entire time series. Human and rhesus prenatal brain was
502 divided into two regions: forebrain together with the midbrain (referred to as the 'brain') and hindbrain
503 (referred to as the 'cerebellum'). Human and rhesus postnatal 'brain' and 'cerebellum' samples comprise the
504 dorsolateral prefrontal region of the cerebral cortex and lateral cerebellar cortex, respectively. For the other
505 species the dissected 'brain' samples correspond to the cerebral hemispheres (without the olfactory bulbs).
506 The early 'cerebellum' samples correspond to the prepontine hindbrain-enriched brain region (until the
507 period matching a mouse e15.5) and from e16.5 onwards only to the cerebellum. For mouse, rat and rabbit,
508 the earliest developmental samples consist of whole brains, which were analysed as part of the brain time
509 series (Supplementary Table 1). We dissected heart samples across the entire time series. At the earliest
510 stage sampled, the heart is beating and the four chambers are already present³⁷. For most species, the liver
511 could also be individually dissected at the start of the time series. The developing gonads are visible as a
512 paired structure on the ventromedial surface of the mesonephros before the start of the time series.
513 However, depending on the species, we were only able to completely isolate the developing gonads at later
514 stages. The same was true for the developing kidneys. In chicken only the left ovary develops and this was
515 the one dissected. The dissections include the main organ structures/cell types in all species but, with the
516 exception of the early samples, they did not include the whole organ.

517 518 **RNA extraction and sequencing**

519 RNA was extracted using the RNeasy protocol from QIAGEN. RNeasy Micro columns were used to extract
520 RNA from small (< 5 mg) or fibrous samples and RNeasy Mini columns were used to extract RNA from larger
521 samples. The tissues were homogenized in RLT buffer supplemented with 40 mM dithiothreitol (DTT) or
522 QIAzol. In order to make sure that we were not introducing technical biases by using two different
523 homogenization procedures, we generated libraries for four samples (two adult rat brains and two adult rat

524 testis) using RLT buffer with dithiothreitol or QIAzol (with two technical replicates). All libraries from the
525 same organ showed a Pearson correlation coefficient ≥ 0.99 irrespective of the homogenization procedure
526 (the median correlation between replicates in the rat dataset was 0.99). RNA quality was assessed using the
527 Fragment Analyzer (Advanced Analytical). The RNA-seq libraries were created using the TruSeq Stranded
528 mRNA LT Sample Prep Kit (Illumina) and sequenced on the HiSeq 2500 platform (multiplexed in sets of 6 or
529 8). All libraries are strand-specific, 100 nucleotides single-end, and were sequenced to a median depth of 33
530 million reads at the Lausanne Genomic Technologies Facility (Supplementary Table 2). The sequencing depth
531 was uniform across the libraries (5% and 95% quantiles of 20 and 54 million reads, respectively). A subset of
532 the adult libraries was used in previous publications^{38,39}.

533

534 **QC of the libraries and estimation of expression levels**

535 We mapped the reads from each library against the species reference genome (Supplementary Table 19)
536 using GSNAP (22-10-2014)⁴⁰. The alignments were guided by the known gene annotations and the discovery
537 of novel splice sites was enabled (Supplementary Tables 19-20). We used HTSeq (0.6.1)⁴¹ to generate read
538 counts for the set of protein-coding genes (Supplementary Tables 19-20). Only uniquely mapping reads were
539 allowed. We normalized the count data using the method TMM as implemented in the package EdgeR
540 (3.14.0)⁴². EdgeR was also used to generate the expression tables used in the study. Expression levels were
541 calculated as cpm (counts per million) or in RPKM (reads per kilobase of exon model per million mapped
542 reads). The alignment files were manipulated using samtools (0.1.18)⁴³ and general alignment statistics
543 created using Picard (1.86)⁴⁴ (Supplementary Table 20).

544

545 We used Fragment Analyzer's RQN values to evaluate the quality of the samples and generally generated
546 sequencing data for those with high values (≥ 7). However, because we also sequenced libraries with lower
547 RQN values, we performed an additional check on RNA integrity after sequencing. We used Picard's
548 "CollectRnaSeqMetrics" tool to calculate the distribution of read coverage along transcripts. RNA
549 degradation leads to a bias in read coverage by favoring the 3' end of genes and this can be identified by
550 calculating the median 3' bias of transcript coverage. We excluded from our dataset all libraries that showed
551 a significant 3' bias in read coverage.

552

553 We evaluated the quality of the sequenced libraries using unsupervised hierarchical clustering (hclust) and
554 PCA (FactoMineR 1.34⁴⁵) as implemented in R⁴⁶. In a PCA the developmental samples from an organ of a
555 given species should be ordered by developmental time in a characteristic U or V-shape⁴⁷. Samples with low
556 RNA quality, insufficient sequencing depth, or showing potential contamination with other tissues appeared
557 as outliers in the organ PCAs and were excluded (the outlier status was confirmed using hierarchical
558 clustering). The global and organ-specific PCAs used as input the read counts after applying the variance
559 stabilizing transformation (vst) implemented in DESeq2 (1.12.4)⁴⁸. The sex of the samples was confirmed
560 using the female-specific genes *Xist* (eutherians), *Rsx* (opossum) and *CDC34* (chicken) (and for eutherians
561 with available Y chromosomes also with the Y-linked gene *Ddx3y*) using Bedtools (2.18)⁴⁹. Finally, we removed
562 from the dataset libraries where the correlation among replicates (Spearman's ρ) was lower than 0.90. We
563 are making available the libraries that passed the general QC but had correlations with their replicates < 0.90 ,
564 but they were not used in this study and are marked as such in Supplementary Table 2.

565

566 **Developmentally Dynamic Genes (DDGs)**

567 In each organ, we identified the genes with dynamic temporal profiles (DDGs) using maSigPro, an R package
568 designed for transcriptomic time-courses^{50,51}. We used as input the count tables from EdgeR (in cpm), and
569 only excluded genes that did not reach a minimum of 10 reads in at least 3 libraries. We ran maSigPro on the
570 log-transformed time (measured in days post-conception) with a degree = 3 (polynomial). We considered

571 genes as DDGs in an organ when the goodness-of-fit (R^2) was at least 0.3 and the maximum RPKM in that
572 organ was at least 1. The lists of DDGs in each organ and species are provided in Supplementary Tables 3-9.

573

574 We identified differences between species and organs in the number of DDGs (Extended Data Fig. 2a).
575 However, technical aspects of the datasets can explain these differences, particularly those between species.
576 First, due to the nature of the statistical test used, the power to call differential temporal expression
577 depended on the magnitude of the expression change and on the agreement between the biological
578 replicates. Smaller expression changes could only be detected if there was strong agreement between the
579 biological replicates. There are differences between species in the median correlation across replicates
580 (Spearman's ρ : 0.94 – 0.99) and these are strongly correlated with the number of DDGs detected ($\rho = 0.66$, P
581 $< 10^{-6}$). Two factors contribute to the species differences in the correlations among replicates. One is the
582 amount of genetic diversity (e.g. lower in mouse than human); the other is how close biological replicates
583 are in terms of development. In rodents the biological replicates are from identical developmental stages
584 (sometimes even the same litter) but in primates the biological replicates span developmental periods.
585 Second, there are differences between species and organs in the length of the time series (Supplementary
586 Table 1). Notably, in chicken and rhesus we are missing the earliest developmental stages, when key
587 developmental processes occur. Finally, some differences could also derive from differences in genome
588 annotation.

589

590 We characterized DDGs using 3 different metrics of functional constraint: 1) the residual variation intolerance
591 score (RVIS); 2) the probability of being loss-of-function intolerant (pLI); and 3) the selection against
592 heterozygous loss of gene function (s_{het}). All metrics were applied to data from the Exome Aggregation
593 Consortium (ExAC)⁹. We obtained the pLI and RVIS scores from ref. ⁷ and s_{het} from ref. ¹⁰. We also used the
594 CNV intolerance score as applied to the ExAC data from ref. ¹¹. The lists of TFs were from the animalTFDB
595 (version 2.0)⁵².

596

597 **Stage correspondences across species**

598 We identified stage correspondences across species using the set of 1:1 DDGs in all species. Because of the
599 shorter time-series we did not require genes to be DDGs in rhesus. We used the combined information from
600 the somatic organ DDGs to calculate the Spearman correlations between all stages in mouse and all stages
601 in each of the other species (using for each stage the median across replicates). We then ran the dynamic
602 time warping algorithm implemented in the R package 'dtw' (1.18-1)⁵³ to identify the optimal alignment
603 between each of the two time series. We ran dtw using as step pattern 'symetricP05' (except for rhesus and
604 chicken where the late fetal start required us to use 'asymmetric' with 'begin.open=T'). When a stage in a
605 given species matched two or more stages in mouse we kept the one with the highest correlation (Extended
606 Data Figs. 3a-b). Our cross-species stage correspondences recapitulated the stage correspondences based on
607 the Carnegie staging for all species except rabbit (shifted 1-2 days; Extended Data Fig. 3a). An independent,
608 neural development-based stage assignment across mammals⁵⁴, suggested an even more pronounced shift
609 (3-4 days) forward for rabbit.

610

611 We then evaluated whether the stage correspondences based on the combined information from the
612 somatic organs were consistent with the information available for each individual organ. For each organ and
613 stage in mouse, we selected in the other species the stage with the maximum correlation plus all stages
614 within 1% of the maximum correlation. We then fitted a local polynomial regression ('loess') to identify the
615 organ-specific correspondences (Extended Data Fig. 6). Overall, the global stage correspondences are within
616 the 98% confidence interval of organ-specific correspondences, suggesting that a single stage
617 correspondence can be used for all organs. But there are exceptions. The heart-specific correspondence

618 between mouse and opossum differs from the global correspondence early in development, suggesting that
619 in relation to the other organs, heart development in opossum could be shifted forward, i.e. be in a more
620 advanced developmental stage. Early opossum development is characterized by heterochronies in the
621 craniofacial, axial and limb skeleton that allow the neonates to crawl without their mother's help to the teat
622 immediately after birth^{16,55}. It is possible that heart development is also shifted forward to accommodate the
623 greater demands of what is postnatal life in opossum, and still prenatal life in the other species. The other
624 potential exception applies to early ovary development in human and rabbit, where we observe development
625 to be shifted forward in the two species. Using the ovary-specific correspondences, the heterochronies
626 associated with the onset of meiosis during oogenesis in these species are even more pronounced than when
627 using the global stage correspondences (Extended Data Fig. 6).

628
629 We were underpowered to detect shifts in individual organs that encompass a small number of adjacent time
630 points. Throughout organ development, the correlation between adjacent stages is, as expected, high, and
631 we would only be able to detect small shifts if they led to a high discordance between species (i.e. significantly
632 lower correlations for a short interval when compared to the rest of the time-series). The only instance of
633 this in our dataset was during testis development, in association with the onset of meiosis (inset in Extended
634 Data Fig. 6). The onset of meiosis marks the beginning of dramatic changes in cell composition in the testis²⁰,
635 which make the transcriptomes that flank this event distinct from each other (Extended Data Fig. 7e), thereby
636 allowing the detection of significant differences between species between adjacent stages.

637

638 **Periods of greater transcriptional change**

639 For each species, we identified the genes that are differentially expressed between adjacent time points
640 (based on the cross-species stage correspondences) using DESeq2 (1.12.4)⁴⁸. We required the adjusted *P*-
641 value to be ≤ 0.05 and the \log_2 fold-change to be ≥ 0.5 . Differences between species in the number of
642 replicates and in the correlation among the replicates (above) impacted our power to call differential
643 expression. Both factors led to lower power to detect differential expression in primates than in mouse, rat
644 and rabbit. Therefore, we are likely underestimating the amount of transcriptional change in humans.

645

646 **Relationships between evolution and development**

647 In Figs. 3a and 3f (and in Extended Data Figs. 8c and 9e) we compare the tolerance to functional mutations
648 and the time- and tissue-specificity of genes employed early vs. late in development in human and mouse.
649 For each species we identified these genes in the following way. First, we identified the most common
650 profiles in each organ using the soft-clustering approach (c-means) implemented in the R package mFuzz
651 (2.32.0)^{56,57}. The clustering was restricted to DDGs and we used as input the read counts after applying the
652 variance stabilizing transformation (vst) to the raw counts implemented in DESeq2 (1.12.4)⁴⁸. The number of
653 clusters was set to 6-8 depending on the organ. For each organ, we settled on a cluster number when
654 increasing it would not add a new cluster but instead split a previous cluster in two. We considered that a
655 cluster was split into two when the median profile of the genes in the two new clusters was similar and when
656 functional enrichment analyses were also similar between the clusters. Between 86-92% of genes in mouse
657 and 89-93% of genes in human were clearly assigned to one of the clusters (cluster membership ≥ 0.7).
658 Among these genes, those assigned to clusters characterized by a decrease in expression during development
659 were classified as genes employed early in development and those assigned to clusters with the opposite
660 profile were classified as genes employed late in development. Genes assigned to clusters with other profiles
661 were classified as other. The classification of each gene in each organ as "Early", "Late", "Other" or "NA" (if
662 a gene is not DDG in the organ or if it has a membership < 0.7) is provided in Supplementary Tables 3 (mouse)
663 and 6 (human).

664

665 In Fig. 3b (and Extended Data Fig. 8d), we used a set of neutrally ascertained mouse knockouts that consists
666 of 2676 protein-coding genes: 646 are classified as lethal, 257 as subviable (less than 12.5% of expected pups)
667 and 1773 as viable. These were the data on viability available for download on June 7, 2017 from the
668 International Mouse Phenotyping Consortium (IMPC)²⁷. For each developmental stage, the denominator is
669 the number of genes expressed that were tested for lethality and the numerator the genes among those that
670 resulted in a lethal phenotype. In Extended Data Fig. 8d we also include in the numerator the genes that
671 resulted in a subviable phenotype (top) and exclude from the analysis a set of housekeeping genes identified
672 by Eisenberg and Levanon⁵⁸ (bottom). We excluded housekeeping genes because they are typically most
673 highly expressed early in development and are enriched among lethals²².

674

675 In Fig. 3c we used a set of genes with evidence for coding-sequence adaptation in mammals identified by
676 Kosiol and colleagues²⁸. For each developmental stage, the denominator is the number of expressed genes
677 that were tested for signatures of positive selection and the numerator is the number of genes among those
678 with evidence for positive selection.

679

680 In Fig. 3d we plotted the age of the transcriptome for each developmental stage. The “age of the
681 transcriptome” was inspired by the Transcriptome Age Index (TAI) developed by Domazet-Lošo and Tautz⁵⁹
682 but differs fundamentally from it in that we are dating the emergence of individual genes and not of gene
683 families (i.e. the emergence of the founder member of a gene family). The TAI measure is a weighing
684 procedure (weighted arithmetic mean) that gives greater weight to young duplicates. The age of the
685 duplicates was determined based on syntenic alignments across vertebrates and parsimony as previously
686 described⁶⁰. The pipeline was run for human, mouse, rat and chicken (based on Ensembl 69 annotations).
687 Most new genes emerged via SSDs in mammals⁶⁰. Genes predating the vertebrate split were given a score of
688 1, genes shared by amniotes were given a score of 2 and so on, until genes that are species-specific were
689 given the maximum score. The range of the score differed between species depending on the number of
690 outgroup lineages available (more lineages allowed for more details in the phylogeny) and therefore this
691 index cannot be compared across species, only within species (i.e. across organs). The score assigned to each
692 gene was multiplied by the gene’s expression (but only if RPKM > 1). The results reported used the log2
693 transformed RPKM values but similar trends were obtained using the raw RPKM values. Higher values
694 indicate larger contributions of recently duplicated genes (i.e. younger transcriptomes).

695

696 **Pleiotropy indexes**

697 The time- and tissue-specificity indexes are based on the Tau metric of tissue-specificity⁶¹. To calculate our
698 tissue-specificity index, we applied the Tau formulation to the maximum expression observed during
699 development in each organ. The time-specificity index uses the Tau formulation for time-points instead of
700 organs. Both indexes range from 0 (broad expression) to 1 (restricted expression). These indexes are provided
701 in Supplementary Tables 3-9.

702

703 **Comparing developmental trajectories**

704 We compared developmental trajectories between human, mouse, rat, rabbit and opossum. Rhesus and
705 chicken were not included because their time series start at a late fetal stage. We used GPclust, a method to
706 cluster time-series using Gaussian processes⁶²⁻⁶⁴, to identify the most common developmental trajectories in
707 each organ. We used the expression (vst-counts) of all available orthologous DDGs as the input (median
708 across replicates for matching stages only). We set the noise variance (k2.variance.fix) to 0.7. GPclust
709 assigned each gene the probability of belonging to each of the trajectories (clusters). We then inferred within
710 a phylogenetic framework the probability that there were changes in developmental trajectories, i.e. that

711 genes changed their cluster assignment in specific branches. We did this in a two-step approach. First, we
712 inferred ancestral cluster probabilities along the tree by calculating the weighted averages from the child-
713 nodes. The weights are given by the inverse branch lengths, which were retrieved from TimeTree⁶⁵, so that
714 closer child-nodes have more weight. To detect changes in the overall pattern at each branching in the tree
715 we calculated the probability that its two nodes are in the same cluster. If the probability was below 1% we
716 called the node as having changed. Second, after identifying all such nodes we mapped the change to one of
717 the two branches by comparing the two children of the node with the outgroup node. The results are
718 provided in Supplementary Tables 14-18.

719

720 It was not always possible to identify the specific branch where changes occurred. This was either because
721 changes were also detected at neighboring nodes (making unclear where the change occurred) or because
722 two nodes differed at the threshold used (1%) but they were both not different from their joint closest
723 relative (e.g. when a call was made for mouse vs. rat but neither for mouse vs. rabbit nor rat vs. rabbit). These
724 calls are classified as NA in Supplementary Tables 14-18. Finally, changes between opossum and the
725 eutherian species could not be polarized because of the lack of an outgroup (classified as Eutherian/opo in
726 Supplementary Tables 14-18). These changes were included in Extended Data Fig. 10. Fig. 4 and Extended
727 Data Fig. 10 summarize the results for genes that have one trajectory change across the phylogeny.

728

729 Differences in developmental trajectories between species can be created by changes in the expression levels
730 of genes in homologous cell populations, by expansions/contractions of homologous cell populations, or by
731 differences in the cell populations that express a given gene (all non-mutually exclusive possibilities). We
732 chose a conservative cutoff (1%) to identify trajectory changes because our aim was to identify those with
733 the largest biological effects. As a consequence we are likely enriching for differences between species
734 created by abrupt changes in the size of homologous cell populations, differences in the cell populations that
735 express a given gene, and/or by differences in expression levels of genes in homologous cell populations that
736 are time-specific (as opposed to being progressive during development).

737

738 **The impact of organ complexity on estimates of species divergence**

739 Organ complexity can impact estimates of gene expression. Expression changes in low abundant cell types
740 that can be detected in simpler organs can potentially go undetected in more complex organs¹. Because the
741 brain has a higher cellular complexity than the other organs studied⁶⁶, it may appear to be more conserved
742 between species than it truly is. Indeed, we found brain tissues to be consistently the slowest evolving,
743 irrespective of the variable being measured. Developmental datasets can help address the problem of
744 comparing organs with different levels of complexity. Organs are more homogeneous early in development
745 and then progressively increase in complexity (e.g. the number of distinct cell types increases during
746 development)¹. This means that when we analyze entire time series, we are comparing organs at different
747 levels of complexity, including early in development when organ complexity is lowest. Throughout the entire
748 times series we consistently observed more similarities between species' transcriptomes for the brain than
749 for the other organs (Extended Data Fig. 8b), including at the earliest stages. We also observed that overall
750 organs are most similar across species early in development (when the power to identify differences would
751 be greatest), and then progressively diverge through time (Extended Data Fig. 8b). Finally, the differences
752 between organs were also consistent throughout the entire development when evaluating the percentage
753 of expressed positively-selected genes (Fig. 3c) and the contribution of recent gene duplications (Fig. 3d).
754 Taken together, these observations suggest that the observed differences between organs in their
755 evolutionary rates are independent of organ complexity. We could, however, be underestimating the total
756 divergence of organs, particularly in adults.

757

758 **General statistics and plots**

759 Unless otherwise stated all statistical analyses and plots were done in R⁴⁶. Plots were created using the R
760 packages ggplot2 (2.2.1)⁶⁷, gridExtra (2.2.1)⁶⁸, reshape2 (1.4.2)⁶⁹, plyr (1.8.4)⁷⁰, and factoextra (1.0.4)⁷¹. All
761 functional enrichment analyses were done using the R implementation of WebGestalt (version 0.0.5)⁷². All
762 packages and versions used are described in Supplementary Table 20.

763

764 **Methods references**

- 765 37. Bruneau, B. G. Signaling and transcriptional networks in heart development and regeneration. *Cold*
766 *Spring Harb. Perspect. Biol.* **5**, a008292 (2013).
- 767 38. Carelli, F. N., Liechti, A., Halbert, J., Warnefors, M. & Kaessmann, H. Repurposing of promoters and
768 enhancers during mammalian evolution. *Nat. Commun.* **9**, 4066 (2018).
- 769 39. Marin, R. et al. Convergent origination of a Drosophila-like dosage compensation mechanism in a
770 reptile lineage. *Genome Res.* **27**, 1974-1987 (2017).
- 771 40. Wu, T. D. & Nacu, S. Fast and SNP-tolerant detection of complex variants and splicing in short reads.
772 *Bioinformatics* **26**, 873-881 (2010).
- 773 41. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput
774 sequencing data. *Bioinformatics* **31**, 166-169 (2015).
- 775 42. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
776 expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).
- 777 43. Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079
778 (2009).
- 779 44. Picard. <http://broadinstitute.github.io/picard> (2015)
- 780 45. Le, S., Josse, J. & Husson, F. FactoMineR: An R Package for Multivariate Analysis. *J. Stat. Softw.* **25**
781 (2008).
- 782 46. R Core Team. R: A Language and Environment for Statistical Computing. (2014).
- 783 47. Anavy, L. et al. BLIND ordering of large-scale transcriptomic developmental timecourses.
784 *Development* **141**, 1161-6 (2014).
- 785 48. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
786 data with DESeq2. *Genome Biol.* **15**, 550 (2014).
- 787 49. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
788 *Bioinformatics* **26**, 841-842 (2010).
- 789 50. Nueda, M. J., Tarazona, S. & Conesa, A. Next maSigPro: updating maSigPro bioconductor package for
790 RNA-seq time series. *Bioinformatics* **30**, 2598-2602 (2014).
- 791 51. Conesa, A., Nueda, M. J., Ferrer, A. & Talon, M. maSigPro: a method to identify significantly
792 differential expression profiles in time-course microarray experiments. *Bioinformatics* **22**, 1096-1102
793 (2006).
- 794 52. Zhang, H. M. et al. AnimalTFDB 2.0: a resource for expression, prediction and functional study of
795 animal transcription factors. *Nucleic Acids Res.* **43**, D76-81 (2015).
- 796 53. Giorgino, T. Computing and Visualizing Dynamic Time Warping Alignments in R: The dtw Package. *J.*
797 *Stat. Softw.* **31**, 1-24 (2009).
- 798 54. Clancy, B., Darlington, R. B. & Finlay, B. L. Translating developmental time across mammalian species.
799 *Neuroscience* **105**, 7-17 (2001).
- 800 55. Smith, K. K. Craniofacial development in marsupial mammals: developmental origins of evolutionary
801 change. *Dev. Dyn.* **235**, 1181-1193 (2006).
- 802 56. Futschik, M. E. & Carlisle, B. Noise-robust soft clustering of gene expression time-course data. *J.*
803 *Bioinform. Computat. Biol.* **3**, 965-988 (2005).
- 804 57. Kumar, L. & Futschik, M. Mfuzz: A software package for soft clustering of microarray data.
805 *Bioinformation* **2**, 5-7 (2007).
- 806 58. Eisenberg, E. & Levanon, E. Y. Human housekeeping genes, revisited. *Trends Genet.* **29**, 569-574
807 (2013).

- 808 59. Domazet-Loso, T. & Tautz, D. A phylogenetically based transcriptome age index mirrors ontogenetic
809 divergence patterns. *Nature* **468**, 815-818 (2010).
- 810 60. Zhang, Y. E., Vibranovski, M. D., Landback, P., Marais, G. A. & Long, M. Chromosomal redistribution
811 of male-biased genes in mammalian evolution with two bursts of gene gain on the X chromosome.
812 *PLoS Biol.* **8**, e1000494 (2010).
- 813 61. Yanai, I. et al. Genome-wide midrange transcription profiles reveal expression level relationships in
814 human tissue specification. *Bioinformatics* **21**, 650-9 (2005).
- 815 62. Hensman, J., Lawrence, N. D. & Rattray, M. Hierarchical Bayesian modelling of gene expression time
816 series across irregularly sampled replicates and clusters. *BMC Bioinformatics* **14**, 252 (2013).
- 817 63. Hensman, J., Rattray, M. & Lawrence, N. D. Fast Nonparametric Clustering of Structured Time-Series.
818 *IEEE Trans. Pattern. Anal. Mach. Intell.* **37**, 383-393 (2015).
- 819 64. Hensman, J., Rattray, M. & Lawrence, N. D. Fast Variational Inference in the Conjugate Exponential
820 Family. *Pattern Analysis and Machine Intelligence, IEEE Transactions on* (2014).
- 821 65. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines, Timetrees,
822 and Divergence Times. *Mol. Biol. Evol.* **34**, 1812-1819 (2017).
- 823 66. Vickaryous, M. K. & Hall, B. K. Human cell type diversity, evolution, development, and classification
824 with special reference to cells derived from the neural crest. *Biol. Rev. Camb. Philos. Soc.* **81**, 425-455
825 (2006).
- 826 67. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2009).
- 827 68. Auguie, B. gridExtra: Miscellaneous Functions for "Grid" Graphics. (2015).
- 828 69. Wickham, H. Reshaping Data with the reshape Package. *J. Stat. Softw.* **21** (2007).
- 829 70. Wickham, H. The Split-Apply-Combine Strategy for Data Analysis. *J. Stat. Softw.* **40** (2011).
- 830 71. Kassambara, A. & Mundt, F. factoextra: Extract and Visualize the Results of Multivariate Data
831 Analyses. (2017).
- 832 72. Wang, J., Vasaiakar, S., Shi, Z., Greer, M. & Zhang, B. WebGestalt 2017: a more comprehensive,
833 powerful, flexible and interactive gene set enrichment analysis toolkit. *Nucleic Acids Res* **45**, W130-
834 37 (2017).

835 **Data availability**

837 Raw and processed RNA-seq data have been deposited in ArrayExpress with the accession codes: E-MTAB-
838 6769 (chicken), E-MTAB-6782 (rabbit) E-MTAB-6798 (mouse), E-MTAB-6811 (rat), E-MTAB-6813 (rhesus), E-
839 MTAB-6814 (human) and E-MTAB-6833 (opossum) (<https://www.ebi.ac.uk/arrayexpress/>). The temporal
840 profiles of individual genes across organs and species can be visualized and downloaded using the web-based
841 application: <http://evodevoapp.kaessmannlab.org>.

842 **Extended Data Figure legends**

843

844 **Extended Data Figure 1 | Organ developmental transcriptomes.** **a**, PC3 and PC4 of the PCA based on 7,696
845 1:1 orthologs depicted in Fig. 1b (each dot represents the median across replicates), and scree plot describing
846 the amount of variance explained by the first 10 PCs. **b**, PCAs of individual organs (n = 7,696 1:1 orthologs).
847 **c**, Correlation of expression levels throughout development between (top) human brain and the other organs
848 (20,345 genes), (bottom) mouse liver and the other organs (21,798 genes). Similar patterns were observed
849 using other organs as the focal organ, and species. For human, the prenatal data are in weeks (w) and
850 postnatally 'new' means newborn, 'sch' school age (7-9 years), 'ya' young adult (25-32 years) and 'sen' senior
851 (58-63 years).
852
853

854

855 **Extended Data Figure 2 | DDGs.** **a**, Number of DDGs identified in each organ and species using (left) the set
856 of 7,696 1:1 orthologs and (right) the set of all protein-coding genes in each species. The horizontal bar

857 depicts the median. Br: Brain, Cr: Cerebellum, He: Heart, Ki: Kidney, Li: Liver, Ov: Ovary, Te: Testis, Te*: Testis
858 pre-sexual maturity. **b**, Number of DDGs per species, including number of organs where they show dynamic
859 expression. *In rhesus ovary development is not covered, hence there are only 6 organs in total. **c**,
860 Relationship between the number of organs in which genes show dynamic expression and the tolerance to
861 functional variants as measured by: the probability of being loss-of-function intolerant (pLI score), the
862 residual variation intolerance score (RVIS) and selection against heterozygous loss-of-function (s_{het} score) (n
863 = 13,160 genes; Wilcoxon rank sum test, two-sided). **d**, Relationship between the number of organs in which
864 genes show dynamic expression and intolerance to duplication and deletion variants (CNV intolerance score;
865 n = 15,728 genes; Wilcoxon rank sum test, two-sided). **e**, Percentage of organ-specific expressed DDGs at
866 each developmental stage. Bars indicate the range between the replicates. For the brain tissues, DDGs are
867 organ-specific in brain and/or cerebellum. Time-points on the x-axis described in Fig. 1a. **f**, Percentage of TFs
868 expressed at each developmental stage. Bars indicate the range between the replicates. Time-points on the
869 x-axis described in Fig. 1a. In **c-d**, box plots depict median \pm 25th and 75th percentiles, whiskers at 1.5 times
870 the interquartile range.

871

872 **Extended Data Figure 3 | Developmental correspondences across species.** **a**, Developmental stage
873 correspondences established in this study and correspondences based on the Carnegie staging (when
874 available). **b**, Using mouse as a reference, a dynamic time warping algorithm was used to select the best
875 alignment (pink line) between the time series based on stage transcriptome correlations combining all
876 somatic organs (n = 8,940 genes/organ combinations). In the human correspondence, “new” means
877 newborn, “tod” toddler (2-4 years), “teen” teenager (13-19 years), “yma” young middle age (39-41 years),
878 “sen” senior (58-63 years).

879

880 **Extended Data Figure 4 | Periods of greater transcriptional change in mouse.** Number of genes differentially
881 expressed between adjacent stages in each organ (\log_2 fold change ≥ 0.5). Solid lines refer to genes that
882 increase in expression and dashed lines to genes that decrease. The biological processes and phenotypes
883 enriched at the peaks of differential expression are detailed in Supplementary Table 13.

884

885 **Extended Data Figure 5 | Periods of greater transcriptional change across species.** Number of genes
886 differentially expressed between adjacent, species-matched, stages for each organ (\log_2 fold change ≥ 0.5).
887 Solid lines refer to genes that increase in expression and dashed lines to genes that decrease. The vertical
888 dotted line marks birth.

889

890 **Extended Data Figure 6 | Organ-specific stage correspondences.** Comparison of the global stage
891 correspondences (based on the combined expression of somatic organs; n = 8,940 genes/organ
892 combinations; black line) with organ-specific correspondences (based on 2,727 genes for brain, 2,146 for
893 cerebellum, 1,276 for heart, 1,486 for kidney, 1,305 for liver, 1,298 for ovary, and 2,153 for testis; colored
894 lines). With the exception of early heart development in opossum and early ovary development in rabbit and
895 human, the global correspondences are within the 98% confidence interval for predictions computed by the
896 loess function (local polynomial regression) for each of the organ-specific correspondences (shaded grey
897 area). The same applies to all organs in mouse-chicken and mouse-rhesus comparisons (data not shown).
898 The inset on the bottom right, shows the Spearman correlation between mouse and rabbit (top) and mouse
899 and human (bottom) for testis transcriptomes using the global stage correspondences (black line) or
900 adjusting for the different start of meiosis across species (orange line; i.e. matching a P14 mouse with a young
901 teenager in human and a P84 rabbit).

902

903 **Extended Data Figure 7 | Heterochronies in gonadal development.** **a**, Temporal dynamics of meiotic genes
904 during ovary development. *SYCP1* is not expressed in human ovary. The genes *SPO11* and *STAG3* are not
905 present in the chicken gene annotations used in this work. **b**, Expression of *Stra8* during ovary development.
906 The vertical bars show the range between the replicates and the horizontal dashed line marks 1 RPKM. **c**,
907 Temporal dynamics of meiotic genes during testis development. The profiles of *Stra8* and *Dmc1* are
908 represented not by their range of expression but by their highest peak of expression. In rhesus, meiosis is
909 known to start around 3-4 years³⁶; our data suggest it had not yet started in the 3-year-old individuals
910 examined. *STRA8* is lowly expressed in the human testis. **d**, Expression of *Stra8* during testis development.
911 The vertical bars show the range between the replicates and the horizontal dashed line marks 1 RPKM. **e**,
912 PCA of ovary and testis development for each species (n = 21,798 protein-coding genes in mouse, 19,390 in
913 rat, 19,271 in rabbit, 20,345 in human, 21,886 in rhesus and 15,481 in chicken).

914

915 **Extended Data Figure 8 | Relationships between evolution and development.** **a**, Observed relationship
916 between evolution and development. Divergence (horizontal distance) can be morphological or molecular.
917 **b**, Transcriptome similarity between three species-pairs throughout development (matched stages) using
918 11,439 1:1 orthologs. Similar trends were obtained using all species-pairs. The weighted average Spearman
919 correlation coefficients are -0.81 ($P = 1 \times 10^{-12}$) for the mouse-rat comparison, -0.69 ($P = 2 \times 10^{-11}$) for mouse-
920 human and -0.42 ($P = 0.0004$) for mouse-opossum. At the bottom are the Spearman correlations between
921 transcriptome correlation coefficients and matched developmental time for each organ and species-pair (** P
922 < 0.02 , * $P < 0.05$). Lines were estimated through linear regression and the 95% confidence interval is shown
923 in grey. **c**, Tolerance to loss-of-function variants (pLI score) for genes with different developmental
924 trajectories in human (top) and mouse (bottom). Lower values mean less tolerance. The pLI scores used for
925 mouse genes are from their human orthologs. The P -values refer to early vs. late comparisons, Wilcoxon rank
926 sum test, two-sided. Box plots depict the median \pm 25th and 75th percentiles, whiskers at 1.5 times the
927 interquartile range. **d**, Percentage of lethal and subviable genes expressed throughout development among
928 a set of 2,686 neutrally ascertained mouse knockouts (top) and the same after excluding housekeeping genes
929 (bottom). Spearman correlations at the bottom of each plot. Lines were estimated through linear regression
930 and the 95% confidence interval is shown in grey.

931

932 **Extended Data Figure 9 | Pleiotropy as a determinant of the evolution of development.** **a**, Relationship
933 between tissue- and time-specificity. Gene developmental profiles illustrate the extremes of the indexes,
934 which range from 0 (broad time/spatial expression) to 1 (specific time/spatial expression). In the gene plots,
935 the x-axis shows the samples ordered by stage and organ and the y-axis shows expression levels. **b**, Functional
936 constraints (measured by pLI) decrease with increasing time- and tissue-specificity (n = 9,965 genes). **All P
937 < 0.01 , Wilcoxon rank sum test, two-sided. **c**, Tissue- and time-specificity of mouse genes identified as lethal,
938 subviable, or viable (n = 2,686; Wilcoxon rank sum test, two-sided). **d**, Levels of functional constraint as
939 measured by RVIS, S_{het} , and pLI scores for the human orthologs of genes identified as lethal, subviable and
940 viable in mouse (n = 2,408; Wilcoxon rank sum test, two-sided). **e**, Tissue- and time-specificity of genes with
941 different developmental trajectories in human (top) and the same after excluding housekeeping genes
942 (bottom). The P -values refer to early vs. late comparisons, Wilcoxon rank sum test, two-sided. In **b-e**, the box
943 plots depict the median \pm 25th and 75th percentiles, whiskers at 1.5 times the interquartile range.

944

945 **Extended Data Figure 10 | Evolution of developmental trajectories.** **a**, Number of genes in each organ that
946 evolved new trajectories across the phylogeny. Includes genes that differ between opossum and eutherians,

947 for which the change cannot be polarized because of the lack of an outgroup. **b**, Distribution of trajectory
948 changes among organs for the different species. The number of genes that changed in each organ is depicted
949 in Fig. 4b. Humans show a relative excess of changes in brain tissues and a relative paucity in testis. $**P = 2$
950 $\times 10^{-5}$ for brain, $P = 0.02$ for cerebellum and $P = 1 \times 10^{-10}$ for testis (from binomial tests where the probability
951 of success is derived from what is observed in mouse, rat and rabbit). **c**, Genes tested for trajectory changes
952 (7,020 genes) in mouse (top) and human (bottom) have significantly lower tissue- and time-specificity than
953 genes not tested for trajectory changes (13,325 genes in mouse and 14,778 in human, Wilcoxon rank sum test,
954 two-sided). **d**, Genes with trajectory changes in mouse (top) and human (bottom) have similar or lower
955 tissue- and time-specificity than genes with conserved trajectories (Wilcoxon rank sum test, two-sided, 'N.S.'
956 means non-significant). **e**, Number of organs in which genes evolved new trajectories in the different species.
957 In **c-d**, the box plots depict the median \pm 25th and 75th percentiles, whiskers at 1.5 times the interquartile
958 range.