

**Review**

Integration of palliative, supportive, and nutritional care to alleviate eating-related distress among advanced cancer patients with cachexia and their family members

**Running Title**

Palliative, supportive, and nutritional care for advanced cancer

**Authors**

1. Koji Amano, M.D.

Department of Palliative Medicine, Osaka City General Hospital

2-13-22 Miyakojima-hondori, Miyakojima-ku, Osaka city, Osaka, Japan, 534-0021

[kojiamano4813@gmail.com](mailto:kojiamano4813@gmail.com)

2. Vickie E Baracos, Ph.D.

Division of Palliative Care Medicine, Department of Oncology, University of Alberta, Cross Cancer Institute

11560 University Avenue, Edmonton, Alberta, Canada, T6G1Z2

[vbaracos@ualberta.ca](mailto:vbaracos@ualberta.ca)

3. Jane B Hopkinson, Ph.D., R.N.

School of Healthcare Sciences, College of Biomedical and Life Sciences, Cardiff University

35-43 Eastgate House, Newport Road, Cardiff, Wales, UK, CF24 0AB

[HopkinsonJB@cardiff.ac.uk](mailto:HopkinsonJB@cardiff.ac.uk)

**Corresponding author**

Koji Amano, M.D.

Department of Palliative Medicine, Osaka City General Hospital

2-13-22 Miyakojima-hondori, Miyakojima-ku, Osaka city, Osaka 534-0021, Japan

Tel.: +81-6-6929-1221

Fax: +81-6-6929-1091

E-mail: [kojiamano4813@gmail.com](mailto:kojiamano4813@gmail.com)

Abstract: 150 words, Text: 4562 words, Tables: 1, Figures: 1, References: 62

## **Abstract**

Advanced cancer patients with cachexia and their families can suffer from *eating-related distress*. This complex entity encompasses patients' struggle to nourish themselves, emotional and social consequences of their inability to maintain food intake, and profound disturbance in family relationships. With evidence-based nutritional care, as well as symptom management to enable food intake, cachexia can be mitigated to some degree. In addition, patients and families require psychosocial support and education to understand and cope with this condition. Only by taking an integrated approach can health care teams alleviate eating-related distress, improve quality of life (QOL), reduce interpersonal conflicts, and alter perceptions of nutritional neglect for patients and families. However, few studies have investigated eating-related distress among patients and families. The aim of this narrative review is to describe what is known about eating-related distress and the roles of integrated palliative, supportive, and nutritional care in improving QOL of patients and families.

## **Key words**

Advanced cancer; palliative care; supportive care; nutritional support; nutrition impact symptoms; eating-related distress

## **1. Introduction**

## **2. Problems**

### **2.1. Nutrition impact symptoms**

### **2.2. Eating-related distress and perceived need for nutritional support in patients**

### **2.3. Eating-related distress and perceived need for nutritional support in family members**

### **2.4. Conflicts over food between patients and family members**

## **3. Treatment approaches**

### **3.1. Role of palliative and supportive care**

### **3.2. Role of nutritional care**

### **3.3. Integration of palliative, supportive, and nutritional care**

## **4. Conclusions**

## **5. Limitations**

## **Acknowledgements**

## **Conflict of interest**

## **Funding**

## **Contribution**

## 1. Introduction


Cachexia is particularly prevalent in patients with cancer of advanced stage and has been linked to deteriorations in quality of life (QOL), performance status, treatment outcomes, and survival [1-9]. Cancer cachexia is defined as a syndrome of body weight loss driven by “a variable combination of reduced food intake and altered metabolism” [1, 2]. Reduced food intake is an important and, in some cases, dominant driver of this body weight loss. A diminished central drive to eat (anorexia) is pervasive, and additionally accompanied by an obstacle course of symptoms that impede food intake (Table 1A). Owing to the importance of reduced food intake, cachexia is classified as a form of (disease-associated) malnutrition and is considered to be partially mitigated by nutritional support [1-4, 6, 8, 9].

Patients with cachexia and their family members may suffer from profound psychosocial distress. Prior qualitative research have identified *eating-related distress* as a significant contributor to overall psychosocial distress [10-17]. Eating-related distress encompasses a myriad of intersecting factors including patients’ and family members’ struggles to cope with cachexia, gross impairment of food intake, disturbance of relationships, and changing roles (Table 1A) [10-22]. The current management of cancer cachexia is primarily focused on pain and symptom management, nutrition interventions, as well as, development of novel drugs with appetite-enhancing actions. However, to adequately address eating-related distress, future clinical practice guidelines must consider holistic, multimodal care including education and psychosocial support (Table 1B, Figure 1) [10-22].

Few studies have investigated eating-related distress among patients and family members; furthermore, the segmentation of palliative, supportive, and nutritional care services complicates the matter [10-22]. Therefore, the aim of this narrative review is to describe eating-related distress and the roles of integrated palliative, supportive, and nutritional care in improving QOL of patients and family members. In particular, we will focus on nutrition impact symptoms, perceived need for nutritional support, and conflicts over food among patients and family members in palliative and supportive care settings.

**Table 1. Integration of palliative, supportive, and nutritional care for eating-related distress**

<b>A. Problem: Eating-related distress in patients and family members</b>
Nutrition impact symptoms impairing food intake [21-25, 28-32]
<ul style="list-style-type: none"><li>• Unpredictable occurrence and course</li><li>• Lack of appetite, early satiety, fatigue, pain, shortness of breath, nausea, vomiting, constipation, diarrhea, abnormal taste, abnormal smell, dry mouth, difficulty swallowing, dental problems, drowsiness, anxiety, depression</li><li>• Physical and psychosocial consequences: body weight loss, altered body image, impaired activities of daily living, reduced well-being</li></ul>
Unsuccessful coping strategies of patients and/or family members [10-22, 34-37]
<ul style="list-style-type: none"><li>• “Fighting back” against the cancer cachexia</li><li>• Gaps between reality and expectation</li><li>• Shortage of correct information about cancer cachexia and nutrition</li><li>• Insufficient nutritional support by health care teams</li><li>• Feelings of loss of control, helplessness, frustration</li></ul>
Conflicts over food between patients and family members [10-22, 34-37]
<ul style="list-style-type: none"><li>• Changes in both patients' food preferences and eating habits of whole families</li><li>• Challenges to spousal and family roles and traditions</li><li>• Family members' pressure on patients to eat</li><li>• Differences in beliefs and perceptions between patients and family members</li></ul>
<b>B. Solution: Integration of palliative, supportive, and nutritional care (Figure 1)</b>
Manage nutrition impact symptoms [21-25, 28-32, 39]
Provide psychosocial support for coping of patients and family members [10-22, 34-37]
Provide nutritional support according to evidence-based clinical practice guidelines [1, 2, 4, 6, 8, 9, 46, 47]
Provide evidence-based information and advice on nutritional care for patients by family members [10-22, 34-37]
Educate patients and family members about cancer cachexia [10-22, 34-37]
Initiate end-of-life discussion with patients and family members [13, 37]
<b>C. Outcome: Effectiveness of integration of palliative, supportive, and nutritional care</b>
Alleviating eating-related distress of patients and family members [18-22]
Improving quality of life in patients and family members [18-22]
Mitigating conflicts between patients and family members [18-22]
Promoting advanced care planning among patients and family members [13, 37]



**Figure 1. Integrated model of palliative, supportive, and nutritional care for eating-related distress**

Adequate management of nutrition impact symptoms, i.e., cancer-associated symptoms as well as side effects of cancer treatments, is essential to enable food intake. Psychosocial and nutritional support, provision of evidence-based information and advice, and education enable patients and family members to cope and alleviate eating-related distress. In addition, end-of-life discussion also contributes to optimizing coping strategies and advanced care planning. Interdisciplinary cooperation among oncologists, palliative care physicians, nurses, dietitians, and others is needed to provide the integrated palliative, supportive, and nutritional care.

## 2. Problems

### 2.1. Nutrition impact symptoms

Individuals with incurable cancers experience a diverse and complex burden of symptoms. Many of these are deemed *nutrition impact symptoms*, because they compromise food intake and in turn drive body weight loss (Table 1A). However, a consensus on the definition of nutrition impact symptoms in cancer patients does not exist. These symptoms may be owing to the cancer itself, cancer treatment, or co-morbidities [23, 24]. Although lack of a central drive to eat (anorexia) is among the most pervasive of cancer-associated symptoms [7, 25], many others are implicated in impaired food intake. Other symptoms implicated include early satiety, fatigue, pain, shortness of breath, nausea, vomiting, constipation, diarrhea, abnormal taste and smell, dry mouth, difficulty swallowing, dental problems, anxiety, depression, and drowsiness. Due to the complex symptom burden of advanced cancer, medical management of pain and symptoms is considered a principle component of cancer cachexia management [4, 6].

An important aspect of cancer cachexia management is the identification and evaluation of symptoms interfering with food intake. The Edmonton Symptom Assessment System-revised (ESAS-r) [26] and Patient-Generated Subjective Global Assessments (PG-SGA) [27] are widely used for symptom assessment, some of their components pertain specifically to reduced food intake [21]. While the ESAS-r is able to assess the severity of symptoms such as lack of appetite and nausea, other symptoms impeding food intake, such as early satiety and diarrhea, are not included. On the other hand, although the PG-SGA is a validated tool that can identify several nutrition impact symptoms, the binary outputs (yes/no) do not allow for assessment of symptom severity [28]. A comprehensive nutrition impact symptom tool specific to cancers of the head and neck has been developed [29], but this has not been validated in other cancer sites. Other tools used to capture reduced food intake include food/fluid records and Visual Analogue Scales [25], however, the nutritional impact symptoms are not captured by these tools. Given the weaknesses of the aforementioned assessment tools, further investigation is needed to develop a comprehensive tool that captures the both the breadth and severity of nutrition impact symptoms. Only then, can the association between impaired food intake and nutrition impact symptoms be adequately explored.

Despite the limitations described above, several previous studies have attempted to characterize symptom profiles of patients with advanced cancer. A study conducted at a cancer cachexia clinic indicated that most patients who experienced body weight loss had three or more uncontrolled symptoms. The most common symptoms were early satiety (62%), constipation (52%), nausea or vomiting (44%), and mood changes (42%) [30]. A study in a nutrition fatigue clinic reported that the five most frequent symptoms were abnormal taste and smell (27%), constipation (19%), abdominal pain (14%), difficulty swallowing (12%), and epigastric pain (10%) [31]. A large, European cohort study of patients from palliative care in- and out-patient units, hospices, and general oncology wards compared 9 symptoms of the ESAS-r between a non-

cachexia group and the cachexia group, categorized using cachexia criteria from the international consensus [2, 32]. Their analysis revealed that the frequency of 6 symptoms, i.e., lack of appetite, fatigue, pain, nausea, drowsiness, and reduced well-being, were significantly higher in the cachexia group [32]. Finally, a survey of advanced cancer patients in palliative and supportive care settings demonstrated that the severity of 16 symptoms, i.e., 9 symptoms of the ESAS-r and 7 of the PG-SGA, were rated moderate in severity with the top 3 being reduced well-being, lack of appetite, and fatigue. The severity of 8 symptoms, i.e., lack of appetite, early satiety, fatigue, diarrhea, abnormal taste, difficulty swallowing, drowsiness, and reduced well-being, were significantly greater in the cachexia group than in the non-cachexia group based on cachexia criteria from the international consensus [2, 21]. In summary, due to the heterogeneity in the patient populations, study settings, and assessment tools used in the studies above, direct comparison of the severity and frequency of nutrition impact symptoms would be challenging. Thus, development of a comprehensive assessment tool is urgently needed to identify and evaluate nutrition impact symptoms in palliative and supportive care settings.

## **2.2. Eating-related distress and perceived need for nutritional support in patients**

Eating-related distress experienced by the patient has been substantially informed by qualitative studies involving in depth interviews [10, 13, 14, 16], and this body of work has been systematically reviewed [11, 12, 15, 17]. Some questionnaires have been devised based on this information, however these require validation [21]. Patients experience many distressing symptoms that reduce food enjoyment and make eating unpleasant. Indeed, lack of appetite, reduced food intake, and ongoing body weight loss could lead patients to feel that their survival is threatened [10-22]. Nevertheless, patients often become resolute in their desire to maintain adequate food intake, forcing themselves to eat despite pain, nausea, and other symptoms. An ensuing challenge is the continued body weight loss in spite of varied efforts to maintain intake [1-4, 6, 8, 9]. The unsuccessful attempts by patients to increase their body weight widens the gap between reality and expectation resulting in escalation of patient distress [11-13]. This phenomenon is called the “Calman gap” [33], and it can escalate or de-escalate depending on patient’s knowledge of cancer cachexia and individual coping strategies [12]. For example, in a small survey of advanced cancer patients in a palliative care unit, the four most frequent eating-related distress items belonged to “coping strategies”. It also suggested that patients had strong concerns over “lack of appetite”, “inability to eat”, and “body weight loss” and that “hopelessness”, “fretting”, and shortage of “correct information” aggravated their distress [18]. Another potential exacerbating factor of eating-related distress may be the inability to halt the body weight loss, as surveys suggest that patients with cachexia had significantly greater eating-related distress than those without cachexia [21].

Lack of credible, proven strategies to manage body weight loss and impaired food intake is a concern expressed by patients [18, 20, 21]. In a survey conducted in an inpatient palliative care unit, 76% of patients

felt that they had unmet needs that could have been addressed by unit staff, e.g., physicians, nurses, and dietitians, with specific knowledge of nutritional support for cancer cachexia [18]. A survey in palliative and supportive care settings suggested that patients with advanced cancer wished to receive support from a health care team, including nutritional counselling (94%), ideas to improve food intake (88%), and oral nutritional supplements (83%). In the case that they became unable to take sufficient nourishment orally, patients expressed needs for parenteral nutrition and hydration (77%), and tube feeding (23%) [20]. Expressed needs for these forms of nutritional support were higher in patients with cachexia versus the non-cachexia group. Of note, tube feeding was regarded as generally undesirable, and almost all respondents to a survey did not wish to receive tube feeding even if they could not eat enough [20]. Finally, it has been shown that patient-perceived need for nutritional counselling is significantly associated with reduced QOL and eating-related distress, particularly in regard to “conflicts over food”, “concerns about food”, and “self-motivated effect related to nutrition” [22].

In conclusion, when patient attempts to combat cachexia, many are disappointed with gaps between reality and expectation. This disappointment can be intensified by a shortage of correct information, lack of provision of nutrition intervention by health care teams, poor coping strategies, and unmet need for nutritional support (Table 1A). Unmet, perceived need for nutritional support is significantly associated with cachexia and increased eating-related-distress and contributes to reduced QOL.

### **2.3. Eating-related distress and perceived need for nutritional support in family members**

Eating-related distress experienced by family members has been studied by interview-based qualitative approaches [10-17], and also by questionnaires-based quantitative approaches [19, 21]. It is known through previous studies that family members suffer when negative impact of cachexia becomes apparent in the patient [10-17, 19, 21]. A systematic review of qualitative studies reported 5 particular themes expressed by family members of advanced cancer patients: “impact on everyday life”, “taking charge”, “need for outside help”, “conflict with patient”, and “emotions” [17]. Additional topics that arose from previous studies included the role of female caregivers, sub-processes of “fighting back”, caregiver depression, and association between eating-related distress and progressive cachexia. A unique mixed methods study of male patients and their female partners also suggested that female caregivers may express feelings of deep concern, frustration, and insufficiency, as well as, making innovative efforts to prepare appealing food [10]. The sub-processes: “fighting back”, “letting nature take its course”, or “waffling” have been explored through previous qualitative studies [34-36], and further characterized in the most recent quantitative study [19, 21]. Specifically, in a 2016 survey of 702 bereaved family members of patients who had received care in palliative care units, the five most common items relating to eating-related distress were all categorized as “fighting back”, and the sixth item was classified as “letting nature take its course” [19]. The authors offered the following four domains for how the “fighting back” sub-process can cause eating-related



distress of family members: (factor 1) “feeling that family members forced the patient to eat to avoid death”, (factor 2) “feeling that family members made great efforts to help the patient eat”, (factor 3) “feeling that eating was a cause of conflict between the patient and family members”, and (factor 4) “feeling that correct information was insufficient”. Next, the authors also identified being a patient’s spouse, fair/poor mental status, factor 1, and factor 4 as independent prognostic factors of major depression in bereaved family members. Insufficient care for eating-related distress among family members, especially spouses with weakened mental status, might be associated with major depression among family members after the patient’s death [19]. Finally, a 2018 survey in palliative and supportive care settings suggested that family members of patients in a cachexia subgroup had significantly greater eating-related distress than those in the non-cachexia group. Additionally, the results of this study seem to show that cancer cachexia is more distressing for family members than for patients in some items, such as those relating to information about the patient’s diet. The questionnaire used in this study has not yet been validated [21].

Similar to patients, family members also express concerns of unmet needs for credible, proven strategies to manage body weight loss and meals. In an exploratory qualitative study, advanced cancer patients with prognosis of a few months were interviewed with their family members. The results revealed family members’ desire to maintain a nourishing role supported by professional advice on nutritional care [37]. Another quantitative study showed that 73% of bereaved family members believed that advanced cancer patients treated at palliative care units require nutritional support. Furthermore, 41% of bereaved family members would like to have received sufficient explanations about the reasons for the patient’s body weight loss [19].

In conclusion, eating-related distress in family members is multifaceted, comprising of themes of caregiver roles, attempts to fight back against the cancer, and caregiver mental health and well-being. Like patients, when family members fight back, many are disappointed by the futile attempts, which is compounded with lack of information, education as well as the perception of caregiver neglect of nutritional support. These experiences coalesce to generate and propagate family member’s eating-related distress (Table 1A).

#### **2.4. Conflicts over food between patients and family members**

Reduced food intake is frequently a source of conflict within families in palliative and supportive care settings [10-22]. Changes in patients' food preferences, changes in eating habits of whole families, challenges to spousal and family roles and traditions, family members' pressure on patients to eat, and differences in beliefs and perceptions between patients and family members can result in conflicts which worsen eating-related distress (Table 1A). A systematic review extracted “conflict with patient” as one of five themes of concern in family members [17]. Patients often feel pressured to eat by family members, and distress may be generated when patients eat solely to satisfy family members. This can make patients feel

dejected and harassed. As a result, patients may choose to isolate themselves or lie to family members to avoid such conflicts. Family members can experience feelings of rejection when food is refused, as well as, feelings of guilt when they argue with patients over patients' lack of dietary intake [10-22]. Family members' pressure may be applied unintentionally or due to a lack of insight into patient's evolving cachexia, symptom burden, food preferences, and eating habits [10-22]. In summary, patients' desire and ability to eat are sometimes incongruent with their family members' expectations. Thus, contrary to family members' intentions, their approach to diet and eating problems may become a barrier to increased food intake for patients, should the patient resist well-intentioned encouragement to eat [10-22].

Despite evidence offered by prior qualitative studies showing significant eating-related distress among patients and family members, quantitative studies suggest that overt conflicts over food between patients and family members occur less often than covert distress. In two earlier studies, the actual conflicts over food was reported in only 6-8% of patient and family members, whereas the perception of need for intervention in such conflicts was higher (18-28%) [18,19]. In a more recent study, patients and family members quoted eating-related distress originating from interpersonal conflict to range between 50-60%. However, overt conflicts over food occurred much less frequently, at 20-30% [21]. Although the proportion who experience such conflicts may be small, the impact is big in the family. Unfortunately, because the quantitative studies were carried out for palliative care patients in one cultural setting, the observed dichotomy between overt conflicts and covert distress may not be applicable to other cultures and traditions, as well as, patients across varying stages of cancer cachexia. Further research is required to establish universality of this observation.

### **3. Treatment approaches**

#### **3.1. Role of palliative and supportive care**

Palliative and supportive care is usually considered as an all-encompassing service providing care from cancer diagnosis to bereavement. The role of palliative and supportive care is for controlling symptoms and improving QOL for patients and their family members [38]. These services are invaluable to cancer patients as they are subjected to many nutrition impact symptoms, and potentially reversible causes of body weight loss. Some of these factors include intestinal motility disorders, esophageal/bowel obstruction, malabsorption, endocrine and metabolic disorders, and inability to sleep. Among them, pain (including oral pain due to mucositis), lack of appetite, early satiety, fatigue, drowsiness, nausea, vomiting, constipation, diarrhea, abnormal taste and smell, and difficulty swallowing may be particularly important [21-25, 28-32]. These symptoms need to be systematically assessed and treated, according to clinical practice guidelines, in order to minimize their impact on food intake. As evidence of this, a prospective trial in 2013 demonstrated that cancer patients receiving effective symptom management from a multidisciplinary palliative care service experienced concurrent improvement of appetite [39].

Although palliative and supportive care advocate for holistic, multimodal care for cancer patients, several limitations exist within this field. Despite clinicians' best efforts, up to 50% of patients with cancer have symptoms that have not been addressed [40]. Furthermore, eating-related distress has not been investigated in previous robust interventional studies. Once again, it is evident that a comprehensive assessment tool is needed to assess nutrition impact symptoms and related distress. Next, palliative and supportive care teams typically do not yet offer psychosocial interventions for cachexia-related distress. As such, communicative and educational interventions, tailored to patient's prognosis and severity of cachexia, need to be developed [10-22]. These interventions may help patients and family members to cope with refractory cachexia, defined by the presence of rapidly progressive cancer unresponsive to anticancer therapy, and cachexia, characterized by intense catabolism and the presence of factors that render active management of body weight loss no longer possible or appropriate. End-of-life discussion and advanced care planning are an integral part of this conversation (Table 1B, 1C, Figure 1).

### **3.2. Role of nutritional care**

Gaps in the availability or access to nutrition care is a major source of distress for patients and family members. Despite the best intentions, health care professionals may be unable to reliably identify the presence of insufficient dietary intake and implement active nutritional management [41]. Thus, evidence-based clinical practice guidelines for nutritional care in the setting of advanced cancer are established to serve as health care professionals' road maps for identification of malnutrition and the application of nutritional therapy. The 2017 European Society for Clinical Nutrition and Metabolism (ESPEN) guidelines are of high quality based on their scope and purpose, rigor of development, applicability, and editorial independence [6, 42]. Specifically, the ESPEN strongly recommends: 1) Screening/monitoring for nutrition risk in all patients with cancers of advanced stage, with in depth nutritional assessment for patients who screen positive. 2) Providing nutritional counselling and oral nutritional supplements as the 1<sup>st</sup> line approach, with escalation to artificial nutrition and hydration according to specified criteria.

There is some evidence that body weight loss can be stabilized or even reversed to some degree by nutritional intervention. A 2017 prospective randomized controlled trial demonstrated that nutritional support resulted in a trend towards body weight gain in adult females with cancer cachexia [43]. A 2018 retrospective study demonstrated that food, energy, and protein intake are associated with body weight gain in patients with advanced cancer referred to a multidisciplinary clinic for management of cachexia [44]. A 2018 meta-analysis demonstrated that intake of high energy, high protein supplements enriched with n-3 fatty acids resulted in body weight gain in patients undergoing chemotherapy or chemo-radiation with palliative intent [45]. In the very advanced cancer patients, only preliminary studies have shown that nutritional support may be associated with beneficial effects, i.e., prevented bedsores, improved

performance status [46, 47].

The use of artificial nutrition is controversial for a variety of reasons. Patient and family members' views on artificial nutrition are difficult to capture because views on artificial nutrition are often enmeshed with views on artificial hydration. It is also unclear whether patients and family members can distinguish between the two [20]. There are profound cultural and ethnic differences in perceptions of artificial hydration during the last weeks of life [48]. For example, in Japan, a large study of the general public and bereaved family members showed that 33-50% of the respondents thought that artificial hydration should be continued until death [49]. In Germany, a survey of cancer patients revealed that 23% wished for both artificial nutrition and hydration at the end-of-life. Increasing age was significantly correlated with a decision to forgo artificial nutrition and hydration [50]. Finally, many patients and family members may attach significant meaning, hope and dignity to artificial hydration which substantially increases the difficulty of discontinuation of this treatment [51]. In summary, these studies suggest that the practice of artificial nutrition is complicated by perceptions and beliefs, cultural background, and age of patients.

According to the ESPEN guideline on ethical aspects of artificial nutrition and hydration, decisions on artificial nutrition and hydration have to consider social, cultural, emotional, and existential aspects as well as the patients' spiritual and ethnic background [52]. However, the ESPEN guidelines on nutrition in cancer patients also suggest that artificial nutrition and hydration are unlikely to provide any benefit for most patients in very advanced terminal phase [6]. Thus, initiation or discontinuation of artificial nutrition must be accompanied by acceptable explanation of benefits versus risk to patients and family members, while taking into account patients' prognoses and goals of care.

### **3.3. Integration of palliative, supportive, and nutritional care**

Multimodal care has been recommended as a treatment approach for advanced cancer patients with cachexia [4, 6, 53-56]. A systemic review demonstrated that multimodal rehabilitation program incorporating exercise and nutritional support improved many outcomes, e.g., improvements in physical endurance and depression, in patients with incurable cancer. Although the authors slightly described symptom management, there were no description about effects of management of nutrition impact symptoms on such outcomes [56]. Effective multimodal care is predicated on a seamless integration of symptom management, psychosocial support, nutritional support, and rehabilitation. There is relatively little research evaluating the efficacy of these combined interventions and their ability to simultaneously improve symptoms, distress, food intake, body weight, physical function, and QOL [39, 56-60]. The most encouraging data comes from clinical observational studies [39, 44]. In these studies, a team-based approach, involving palliative care physicians, nurses, dietitians, and occupational therapists, is associated with significant improvement in food intake, nutritional indices, distress, concurrent with successful

management of pain and symptoms. These results suggest that multimodal care needs to be managed by an interdisciplinary team consisting of specialists e.g., trained physicians, nurses, dietitians, pharmacists, and physical/occupational therapists.

In spite of the merits of multimodal care in advanced cancer patients, several limitations are evident. Some patients with high symptom burden only achieved moderate improvement in their symptoms [39, 44]. Moreover, there often exists a gap between the ideal team environment and reality. Innovative local programs, such as a nurse-led walk-and-eat intervention, should not be discounted, as it could also improve patient outcomes, i.e., preserving functional walking capacity and nutritional status [61]. Finally, nutrition impact symptoms and eating-related distress are often not evaluated systematically. One recently developed instrument is the Quality of Life Questionnaire-Cachexia 24 (QLQ-CAX24), which offers a cachexia specific module for the assessment of QOL [62].

Adapting from previous successes in multimodal care research, we would like to propose an integrated model of palliative, supportive, and nutritional care to alleviate eating-related distress (Table 1, Figure1). Our strategy is predicated on the adequate management of nutrition impact symptoms, psychosocial and nutritional support, provision of evidence-based information and advice, education about cancer cachexia, and initiation of end-of-life discussions. Anticipated outcomes from our integrated strategy include improvement in QOL, mitigation of conflicts, and promotion of advanced care planning among patients and family members by supporting for adaptation and coping with nutritional impact symptoms and eating-related distress. Integral to the success of this strategy is the involvement and cooperation between oncologists, palliative care physicians, nurses, dietitians, and other healthcare professionals.

#### **4. Conclusions**

We summarized knowledge of eating-related distress in patients and family members and identified some potential priorities for future research. We found limited information about this important detriment to QOL for patients and family members, and there are effectively no well-developed educational tools or psychosocial interventions to alleviate this distress. There is a lack of validated tools for the clinical assessment of eating-related distress and its response to interventions. More robust evidence regarding breadth and depth of nutrition impact symptoms and their association with distress are needed. Palliative, supportive, and nutrition care have varying availability and in many contexts are implemented in isolation of one another. To inform and support patients and family members to cope effectively, health care professionals must appreciate the complexities and intricacies of eating-related distress. We propose that this will be facilitated by close coordination of palliative, supportive, and nutritional care.

#### **5. Limitations**

The scope of the review was confined to caring for nutrition impact symptoms, eating-related distress, need for nutritional support, and conflicts over food among patients and family members. There might be studies on the other aspects of cancer cachexia and the management of symptoms and malnutrition during/after cancer treatments, which were not included.

### **Acknowledgements**

The authors have read and complied with the principles of ethical authorship of the journal. We would like to thank Dr Catherine Kubrak, Dr Lisa Martin, and Dr Zhimeng Jia for review and discussion of the manuscript.

### **Conflict of interest**

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### **Funding**

The authors received no financial support for the research, authorship, and/or publication of this article.

### **Contribution**

Study concept and design: KA and VB

Data acquisition: KA, VB, and JH

Data interpretation: KA, VB, and JH

Manuscript preparation: KA, VB, and JH

Manuscript editing and review: KA, VB, and JH

## References

1. Radbruch L, Elsner F, Trottenberg P, et al. Clinical practice guidelines on cancer cachexia in advanced cancer patients with a focus on refractory cachexia: European clinical guidelines. Aachen, Department of Palliative Medicine/European Palliative Care Research Collaborative 2010.
2. Fearon KC, Strasser F, Anker SD, et al. Definition and classification of cancer cachexia: an international consensus. *Lancet Oncol* 2011; 12: 489–495.
3. Tisdale MJ. Mechanisms of cancer cachexia. *Physiol Rev* 2009; 89: 381–410.
4. Fearon K, Arends J, Baracos V. Understanding the mechanisms and treatment options in cancer cachexia. *Nat Rev Clin Oncol* 2013; 10(2): 90-99.
5. Amano k, Maeda I, Morita T, et al. Clinical implications of C-reactive protein as a prognostic marker in advanced cancer patients in palliative settings. *J Pain Symptom Manage* 2016; 51(5): 860-867.
6. Arends J, Bachmann P, Baracos V, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr* 2017; 36: 11-48.
7. Amano K, Maeda I, Morita T, et al. C-reactive protein, symptoms and activity of daily living in patients with advanced cancer receiving palliative care. *J Cachexia Sarcopenia Muscle* 2017; 8: 457-465.
8. Baracos VE, Martin L, Korc M, et al. Cancer-associated cachexia. *Nat Rev Dis Primers* 2018. doi: 10.1038/nrdp.2017.105.
9. Sadeghi M, Keshavarz-Fathi M, Baracos V, et al. Cancer cachexia: Diagnosis, assessment, and treatment. *Crit Rev Oncol Hematol* 2018; 127: 91-104.
10. Strasser F, Binswanger J, Cemy T, et al. Fighting a losing battle: eating related distress of men with advanced cancer and their female partners: a mixed methods study. *Palliat Med* 2007; 21(2): 129–137.
11. Hopkinson JB, Okamoto I, Addington-Hall JM. What to eat when off treatment and living with involuntary weight loss and cancer: a systematic search and narrative review. *Support Care Cancer* 2011; 19(1): 1-17.
12. Oberholzer R, Hopkinson JB, Baumann K, et al. Psychosocial effects of cancer cachexia: A systematic literature search and qualitative analysis. *J Pain Symptom Manage* 2013; 46(1): 77-95.
13. Hopkinson JB. Psychosocial impact of cancer cachexia. *J Cachexia Sarcopenia Muscle* 2014; 5: 89-94.
14. Reid J. Psychosocial, educational and communicative interventions for patients with cachexia and their family carers. *Curr Opin Support Palliat Care* 2014; 8(4): 334–338.
15. Cooper C, Burden ST, Cheng H, Molassiotis A. Understanding and managing cancer-related weight loss and anorexia: insights from a systematic review of qualitative research. *J Cachexia Sarcopenia Muscle* 2015; 6(1): 99-111.
16. Hopkinson, JB. Food connections: A qualitative exploratory study of weight- and eating-related distress in families affected by advanced cancer. *Eur J Oncol Nurs* 2016; 20: 87-96.

17. Wheelwright S, Darlington AS, Hopkinson JB, et al. A systematic review and thematic synthesis of quality of life in the informal carers of cancer patients with cachexia. *Palliat Med* 2016; 30(2): 149-160.
18. Amano K, Maeda I, Morita T, et al. Need for nutritional support, eating-related distress and experience of terminally ill cancer patients: A survey in an inpatient hospice. *BMJ Support Palliat Care* 2016; 6(3): 373-376.
19. Amano K, Maeda I, Morita T, et al. Eating-related distress and need for nutritional support of families of advanced cancer patients: a nationwide survey of bereaved family members. *J Cachexia Sarcopenia Muscle* 2016; 7(5): 527-534.
20. Amano K, Morita T, Miyamoto J, et al. Perception of need for nutritional support in advanced cancer patients with cachexia: a survey in palliative care settings. *Support Care Cancer* 2018; 26(8): 2793-2799.
21. Amano K, Morita T, Koshimoto S, et al. Eating-related distress in advanced cancer patients with cachexia and family members: a survey in palliative and supportive care settings. *Support Care Cancer* 2018; Dec 15. doi: 10.1007/s00520-018-4590-6.
22. Koshimoto S, Arimoto M, Saitou K, et al. Need and demand for nutritional counselling and their association with quality of life, nutritional status and eating-related distress among patients with cancer receiving outpatient chemotherapy: a cross-sectional study. *Support Care Cancer* 2019; Jan 14. doi: 10.1007/s00520-018-4628-9.
23. Kubrak C, Olson K, Jha N, et al. Nutrition impact symptoms: key determinants of reduced dietary intake, weight loss, and reduced functional capacity of patients with head and neck cancer before treatment. *Head Neck* 2010; 32(3): 290-300.
24. Anandavadivelan P, Martin L, Djärv T, et al. Nutrition Impact Symptoms Are Prognostic of Quality of Life and Mortality after Surgery for Oesophageal Cancer. *Cancers (Basel)*. 2018 Sep 7;10(9). pii: E318. doi:10.3390/cancers10090318.
25. Martin L, Kubrak C. How much does reduced food intake contribute to cancer-associated weight loss? *Curr Opin Support Palliat Care* 2018; 12(4): 410-419.
26. Bruera E, Kuehn N, Miller MJ, et al. The Edmonton symptom assessment system (ESAS): a simple method for the assessment of palliative care patients. *J Palliat Care* 1991; 7: 6-9.
27. Bauer J, Capra S, Ferguson M. Use of the scored patient generated subjective global assessment (PG-SGA) as a nutrition assessment tool in patients with cancer. *Eur J Clin Nutr* 2002; 56(8): 779-785.
28. Del Fabbro E, Orr TA, Stella SM. Practical approaches to managing cancer patients with weight loss. *Curr Opin Support Palliat Care* 2017; 11(4): 272-277.
29. Kubrak C, Olson K, Baracos VE. The head and neck symptom checklist©: an instrument to evaluate nutrition impact symptoms effect on energy intake and weight loss. *Support Care Cancer* 2013; 21(11): 3127-3136.



30. Del Fabbro E, Hui D, Dalal S, et al. Clinical outcomes and contributors to weight loss in a cancer cachexia clinic. *J Palliat Med* 2011; 14(9): 1004-1008.
31. Omlin A, Blum D, Wierecky J, et al. Nutrition impact symptoms in advanced cancer patients: frequency and specific interventions, a case-control study. *J Cachexia Sarcopenia Muscle* 2013; 4(1): 55-61.
32. Blum D, Stene GB, Solheim TS, et al. Validation of the consensus-definition for Cancer Cachexia and evaluation of a classification model—a study based on data from an international multicentre project (EPCRC-CSA). *Ann Oncol* 2014; 25(8): 1635-1642.
33. Calman KC. Quality of life in cancer patients – a hypothesis. *J Med Ethics* 1984; 10(3): 124-127.
34. McClemen SE, Degner LF, Harlos MS. Family beliefs regarding the nutritional care of a terminally ill relative: a qualitative study. *J Palliat Med* 2003; 6(5): 737-748.
35. McClement SE, Degner LF, Harlos M. Family responses to declining intake and weight loss in a terminally ill relative. Part 1: fighting back. *J Palliat Care* 2004; 20(2): 93-100.
36. McClement SE, Harlos M. When advanced cancer patients won't eat: family responses. *Int J Palliat Nurs* 2008; 14(4): 182-188.
37. Hopkinson JB. The Nourishing Role: Exploratory Qualitative Research Revealing Unmet Support Needs in Family Carers of Patients With Advanced Cancer and Eating Problems. *Cancer Nurs* 2018; 41(2): 131-138.
38. Hui D, De La Cruz M, Mori M, et al. Concepts and definitions for “supportive care,” “best supportive care,” “palliative care,” and “hospice care” in the published literature, dictionaries, and textbooks. *Support Care Cancer* 2013; 21(3): 659-685.
39. Gagnon B, Murphy J, Eades M, et al. A prospective evaluation of an interdisciplinary nutrition-rehabilitation program for patients with advanced cancer. *Curr Oncol* 2013; 20(6): 310-318.
40. Basch E. Missing Patients' Symptoms in Cancer Care Delivery--The Importance of Patient-Reported Outcomes. *JAMA Oncol* 2016; 2(4): 433-434.
41. Hébuterne X, Lemarié E, Michallet M, et al. Prevalence of malnutrition and current use of nutrition support in patients with cancer. *JPEN J Parenter Enteral Nutr* 2014; 38(2): 196-204.
42. Shen WQ, Yao L, Wang XQ, et al. Quality assessment of cancer cachexia clinical practice guidelines. *Cancer Treat Rev* 2018; 70: 9-15.
43. Kapoor N, Naufahu J, Tewfik S, et al. A prospective randomized controlled trial to study the impact of a nutrition-sensitive intervention on adult women with cancer cachexia undergoing palliative care in India. *Integr Cancer Ther* 2017; 16: 74-84.
44. Nasrah R, Kanbalian M, Van Der Borch C, et al. Defining the role of dietary intake in determining weight change in patients with cancer cachexia. *Clin Nutr* 2018; 37: 235-241.
45. de van der Schueren MAE, Laviano A, Blanchard H, et al. Systematic review and meta-analysis of the evidence for oral nutritional intervention on nutritional and clinical outcomes during

- chemo(radio)therapy: current evidence and guidance for design of future trials. *Ann Oncol* 2018; 29(5): 1141-1153.
46. Amano K, Morita T, Baba M et al. Effect of nutritional support on terminally ill patients with cancer in a palliative care unit. *Am J Hosp Palliat Care* 2013; 30(7): 728-731.
  47. Amano K, Maeda I, Ishiki H et al. A feasibility study to investigate the effect of nutritional support for advanced cancer patients in an inpatient hospice in Japan. *Palliat Med Hosp Care Open J* 2016; 2(2): 37-45.
  48. Torres-Vigil I, Cohen MZ, de la Rosa A, et al. Food or medicine: ethnic variations in perceptions of advanced cancer patients and their caregivers regarding artificial hydration during the last weeks of life. *BMJ Support Palliat Care* 2012; 2(3): 276-279.
  49. Morita T, Miyashita M, Shibagaki M, et al. Knowledge and beliefs about end-of-life care and the effects of specialized palliative care: a population-based survey in Japan. *J Pain Symptom Manage* 2006; 31(4): 306-316.
  50. Bükki J, Unterpaul T, Nübling G, et al. Decision making at the end of life—cancer patients' and their caregivers' views on artificial nutrition and hydration. *Support Care Cancer* 2014; 22(12): 3287-3299.
  51. Cohen MZ, Torres-Vigil I, Burbach BE, et al. The meaning of parenteral hydration to family caregivers and patients with advanced cancer receiving hospice care. *J Pain Symptom Manage* 2012; 43(5): 855-865.
  52. Druml C, Ballmer PE, Druml W, et al. ESPEN guideline on ethical aspects of artificial nutrition and hydration. *Clin Nutr* 2016; 35: 545-556.
  53. Aapro M, Arends J, Bozzetti F, et al., ESMO (European School of Medical Oncology). Early recognition of malnutrition and cachexia in the cancer patient: a position paper of a European School of Oncology Task Force. *Ann Oncol* 2014; 25(8): 1492-1499.
  54. Balstad TR, Kaasa S, Solheim TS. Multimodal nutrition/anabolic therapy for wasting conditions. *Curr Opin Clin Nutr Metab Care* 2014; 17(3): 226-235.
  55. Maddocks M, Hopkinson J, Conibear J, et al. Practical multimodal care for cancer cachexia. *Curr Opin Support Palliat Care* 2016; 10(4): 298-305.
  56. Hall CC, Cook J, Maddocks M, et al. Combined exercise and nutritional rehabilitation in outpatients with incurable cancer: a systematic review. *Support Care Cancer* 2019; 27(7): 2371-2384.
  57. Glare P, Jongs W, Zafiroopoulos B. Establishing a cancer nutrition rehabilitation program (CNRP) for ambulatory patients attending an Australian cancer center. *Support Care Cancer* 2011; 19(4): 445-454.
  58. Parmar MP, Swanson T, Jagoe RT. Weight changes correlate with alterations in subjective physical function in advanced cancer patients referred to a specialized nutrition and rehabilitation team. *Support Care Cancer* 2013; 21(7): 2049-2057.
  59. Parmar MP, Vanderbyl BL, Kanbalian M, et al. A multidisciplinary rehabilitation programme for cancer cachexia improves quality of life. *BMJ Support Palliat Care* 2017; 7(4): 441-449.

60. Uster A, Ruehlin M, Mey S, et al. Effects of nutrition and physical exercise intervention in palliative cancer patients: a randomized controlled trial. *Clin Nutr* 2018; 37(4): 1202-1209.
61. Xu YJ, Cheng JC, Lee JM, et al. A walk-and-eat intervention improves outcomes for patients with esophageal cancer undergoing neoadjuvant chemoradiotherapy. *Oncologist* 2015; 20(10): 1216-1222.
62. Wheelwright SJ, Hopkinson JB, Darlington AS, et al. Development of the EORTC QLQ-CAX24, A Questionnaire for Cancer Patients with Cachexia. *J Pain Symptom Manage* 2017; 53(2): 232-242.