

Orbifold equivalent potentials

Nils Carqueville* Ana Ros Camacho† Ingo Runkel†

nils.carqueville@scgp.stonybrook.edu

anaroscamacho@gmail.com

ingo.runkel@uni-hamburg.de

**Simons Center for Geometry and Physics,
Stony Brook, NY 11794, USA*

*†Fachbereich Mathematik, Universität Hamburg,
Bundesstraße 55, 20146 Hamburg, Germany*

To a graded finite-rank matrix factorisation of the difference of two homogeneous potentials one can assign two numerical invariants, the left and right quantum dimension. The existence of such a matrix factorisation with non-zero quantum dimensions defines an equivalence relation between potentials, giving rise to non-obvious equivalences of categories.

Restricted to ADE singularities, the resulting equivalence classes of potentials are those of type $\{A_{d-1}\}$ for d odd, $\{A_{d-1}, D_{d/2+1}\}$ for d even but not in $\{12, 18, 30\}$, and $\{A_{11}, D_7, E_6\}$, $\{A_{17}, D_{10}, E_7\}$ and $\{A_{29}, D_{16}, E_8\}$. This is the result expected from two-dimensional rational conformal field theory, and it directly leads to new descriptions of and relations between the associated (derived) categories of matrix factorisations and Dynkin quiver representations.

AMS 2010 Subject Classification: 18D05, 14B05

Keywords: orbifolds, ADE singularities, matrix factorisations, pivotal bicategories

Contents

1	Introduction and summary	2
1.1	Orbifold equivalence of potentials	2
1.2	Orbifold equivalence for simple singularities	5
2	Proofs	8
2.1	Pivotal bicategories	8
2.2	Matrix factorisations	11
2.3	Simple singularities	13
3	Comparison to conformal field theory	24

1 Introduction and summary

Let $k \subset \mathbb{C}$ be a field. By a *homogeneous potential in m variables* we mean a polynomial $V \in k[x_1, \dots, x_m]$ such that

$$V(\lambda^{|x_1|}x_1, \dots, \lambda^{|x_m|}x_m) = \lambda^2 V(x_1, \dots, x_m) \quad \text{for all } \lambda \in \mathbb{C}^\times \quad (1.1)$$

and

$$\dim_k(k[x_1, \dots, x_m]/(\partial_{x_1}V, \dots, \partial_{x_m}V)) < \infty \quad (1.2)$$

where $k[x_1, \dots, x_m]$ is viewed as a graded ring by assigning degrees $|x_i| \in \mathbb{Q}_+$ to the variables x_i . As we will only be concerned with homogeneous potentials, we shall refer to them simply as *potentials*.

The purpose of this paper is two-fold. Firstly, we will define an equivalence relation on the set \mathcal{P}_k of all potentials, where the number of variables in the polynomial ring is allowed to vary, but where the field k is kept fixed. Secondly, in the case $k = \mathbb{C}$ we will list all equivalence classes in the restricted set of simple singularities, leading to new equivalences of categories.

In the remainder of this section we describe these two points in more detail, leaving technical details and supplementary discussions for Sections 2 and 3.

1.1 Orbifold equivalence of potentials

The equivalence relation is defined by the existence of a matrix factorisation with certain properties.

We will write x for the sequence of variables x_1, \dots, x_m , and we pick a potential $V \in k[x]$. A *matrix factorisation of V* is a \mathbb{Z}_2 -graded free $k[x]$ -module M together with an odd $k[x]$ -linear endomorphism d_M , the *twisted differential*, such that $d_M \circ d_M = V \cdot 1_M$. Let us denote the two homogeneous subspaces of M with respect to the \mathbb{Z}_2 -grading by M^0 and M^1 , so that $M = M^0 \oplus M^1$. We call $M \equiv (M, d_M)$ a *graded matrix factorisation* if M^0 and M^1 are in addition

\mathbb{Q} -graded,¹ and if the following properties are satisfied: (i) Acting with x_i is an endomorphism of degree $|x_i|$ with respect to the \mathbb{Q} -grading on M . Consequently, V has degree 2. (ii) d_M has degree 1 with respect to the \mathbb{Q} -grading on M .

There are categories of (graded) matrix factorisations whose morphisms are (homogeneous) even linear maps up to homotopy with respect to the twisted differential. These are idempotent complete as they are triangulated and have arbitrary coproducts [Nee]. We denote by $\text{hmf}(k[x], V)$ the idempotent closure of the full subcategory of finite-rank matrix factorisations; this means that its objects are homotopy equivalent to direct summands of finite-rank matrix factorisations. Furthermore, $\text{hmf}^{\text{gr}}(k[x], V)$ is the full subcategory of graded matrix factorisations which are homotopy equivalent to finite-rank ones; $\text{hmf}^{\text{gr}}(k[x], V)$ is also idempotent complete, see [KST, Lem. 2.11].

Let now $V(x), W(y) \in \mathcal{P}_k$ be two potentials, where $x = x_1, \dots, x_m$ and $y = y_1, \dots, y_n$. Then $W(y) - V(x)$ is a potential in the ring $k[x, y]$. Let $X = (X, d_X)$ be a finite-rank graded matrix factorisation of $W(y) - V(x)$. The even and odd parts of $X = X^0 \oplus X^1$ necessarily have the same rank, say r , and we may as well assume that $X^0 = X^1 = k[x, y]^r$. The twisted differential is then given by two r -by- r matrices d^0, d^1 with entries in $k[x, y]$. We think of these as $k[x, y]$ -linear maps $d^0 : X^0 \rightarrow X^1$ and $d^1 : X^1 \rightarrow X^0$, and we call (X, d_X) *rank- r* .

We assign two numbers to X , the *left and right quantum dimensions* [CR2, CM2]

$$\begin{aligned} \dim_l(X) &= (-1)^{\binom{m+1}{2}} \text{Res} \left[\frac{\text{str}(\partial_{x_1} d_X \dots \partial_{x_m} d_X \partial_{y_1} d_X \dots \partial_{y_n} d_X) dy}{\partial_{y_1} W, \dots, \partial_{y_n} W} \right], \\ \dim_r(X) &= (-1)^{\binom{n+1}{2}} \text{Res} \left[\frac{\text{str}(\partial_{x_1} d_X \dots \partial_{x_m} d_X \partial_{y_1} d_X \dots \partial_{y_n} d_X) dx}{\partial_{x_1} V, \dots, \partial_{x_m} V} \right]. \end{aligned} \quad (1.3)$$

These formulas arose in the study of adjunctions in the bicategory of Landau-Ginzburg models, see [CR2, CM2] and Remark 1.8 below. Here we just point out the following properties:

- The definition of the quantum dimensions is independent of the \mathbb{Q} -grading on X . In fact, the same residue expressions also work for ungraded matrix factorisations; in this case however $\dim_l(X)$ and $\dim_r(X)$ are not necessarily numbers, but polynomials in $k[x]$ and $k[y]$, respectively, see Lemma 2.6.
- The quantum dimensions only depend on the isomorphism class of X in $\text{hmf}^{\text{gr}}(k[x, y], W - V)$ or $\text{hmf}(k[x, y], W - V)$, cf. Sections 2.1 and 2.2. Hence

¹Often one rescales the degrees $|x_i|$ by their least common denominator, so that one can work with a \mathbb{Z} -grading. For our purposes the present convention, where the degrees are normalised such that each potential has degree 2, is more convenient (the grading is then better behaved when taking differences of potentials). Note that a rescaled \mathbb{Z} -graded matrix factorisation gives a \mathbb{Q} -graded one, but not necessarily vice versa.

they are independent of the choice of isomorphism between X and $k[x, y]^r \oplus k[x, y]^r$.

- The quantum dimensions are multiplicative for a suitably defined tensor product. In addition, there is a duality $(-)^{\dagger}$ on matrix factorisations which exchanges left and right quantum dimensions (see again Sections 2.1 and 2.2).

Definition and Theorem 1.1. We say that two potentials $V(x), W(y) \in \mathcal{P}_k$ are *orbifold equivalent*, $V \sim W$, if there exists a finite-rank graded matrix factorisation of $W - V$ for which the left and right quantum dimensions are non-zero. This defines an equivalence relation on \mathcal{P}_k .

We shall prove the above statement, as well as Propositions 1.2 and 1.4, in Section 2.2. The name ‘orbifold equivalence’ has its roots in the study of orbifolds via defects in two-dimensional quantum field theories [FFRS2, DKR, CR3], see also Remark 1.8.

The hard part of establishing an orbifold equivalence $V \sim W$ is to construct an explicit matrix factorisation X of $W - V$ that exhibits the equivalence. On the other hand, checking a posteriori that a given X has non-zero quantum dimensions is easy thanks to the formulas (1.3). In particular, to verify our results on ADE singularities (or any other orbifold equivalence) all one has to do is to apply the residue formulas to the explicit matrices we provide (however laborious it may have been to construct them initially).

Two basic properties of orbifold equivalences are:

Proposition 1.2. Let $U(z), V(x), W(y) \in \mathcal{P}_k$.

- (i) (*Compatibility with external sums*) If $V \sim W$, then $V + U \sim W + U$.
- (ii) (*Knörrer periodicity*) $V \sim V + u^2 + v^2$.

To a potential $V \in \mathcal{P}_k$ one assigns a rational number, its *central charge* $c(V)$. Namely, for $V \in k[x_1, \dots, x_m]$ we have

$$c(V) = 3 \sum_{i=1}^m (1 - |x_i|). \quad (1.4)$$

It follows that in the case of ADE singularities V the central charge of V is related to the Coxeter number h_V of the root system associated to the potential V via $c(V) = 3 - \frac{6}{h_V}$. For example, if $V = x_1^d + x_1 x_2^2$, then $|x_1| = \frac{2}{d}$, $|x_2| = 1 - \frac{1}{d}$ and $c(V) = 3 - \frac{3}{d}$, which is the same as the central charge of $W = y_1^{2d} + y_2^2$. The following proposition gives some simple necessary conditions for any two potentials to be orbifold equivalent.

Proposition 1.3. Suppose $V, W \in \mathcal{P}_k$ are orbifold equivalent. Then

- (i) $m - n$ is even, where $V \in k[x_1, \dots, x_m]$ and $W \in k[y_1, \dots, y_n]$.
- (ii) $c(V) = c(W)$.

Part (i) is trivial since the supertrace in (1.3) is zero for an odd matrix. Part (ii) is proved in [CR3, Sect. 6.2].

We expect the converse of Proposition 1.3 to be false. For example, consider the family of potentials $x_1^3 + x_2^3 + x_3^3 + \kappa x_1 x_2 x_3$ with $\kappa \in \mathbb{C}$ which all have central charge 3 and whose zero locus is an elliptic curve in \mathbb{CP}^2 . In analogy with [FGRS, BBR] we expect the potentials for different values of κ to be orbifold equivalent iff the complex structure parameters of the corresponding curves are related by some $\mathrm{GL}(2, \mathbb{Q})$ transformation, resulting in infinitely many equivalence classes for these potentials.

If $V, W \in \mathcal{P}_k$ are orbifold equivalent, then the corresponding categories of matrix factorisations are closely related. Namely, let $X = (X, d_X)$ be a finite-rank graded matrix factorisation of $W(y) - V(x)$ with non-zero quantum dimensions. Let further $M = (M, d_M)$ be a graded matrix factorisation of $V(x)$. Then their tensor product $X \otimes M := (X \otimes_{k[x]} M, d_X \otimes 1_M + 1_X \otimes d_M)$ is a graded matrix factorisation of $W(y)$ (which is necessarily of infinite rank, but still equivalent to a finite-rank factorisation, for $M \neq 0$).

Proposition 1.4. Suppose that $V(x), W(y) \in \mathcal{P}_k$ are orbifold equivalent and let $X \in \mathrm{hmf}^{\mathrm{gr}}(k[x, y], W - V)$ have non-zero quantum dimensions. Then every matrix factorisation in $\mathrm{hmf}^{\mathrm{gr}}(k[y], W)$ occurs as a direct summand of $X \otimes M$ for some $M \in \mathrm{hmf}^{\mathrm{gr}}(k[x], V)$. This remains true if ‘ $\mathrm{hmf}^{\mathrm{gr}}$ ’ is replaced by ‘ hmf ’ everywhere.

1.2 Orbifold equivalence for simple singularities

Now we take $k = \mathbb{C}$ and consider the subset of potentials which define simple singularities.² These fall into an ADE classification and can be taken to be the following elements of $\mathbb{C}[x_1, x_2]$ (see e. g. [Yos, Prop. 8.5]):

$$\begin{aligned}
 V^{(\mathrm{A}_{d-1})} &= x_1^d + x_2^2 & c &= 3 - 3 \cdot \frac{2}{d} & (d \geq 2) \\
 V^{(\mathrm{D}_{d+1})} &= x_1^d + x_1 x_2^2 & c &= 3 - 3 \cdot \frac{2}{2d} & (d \geq 3) \\
 V^{(\mathrm{E}_6)} &= x_1^3 + x_2^4 & c &= 3 - 3 \cdot \frac{2}{12} \\
 V^{(\mathrm{E}_7)} &= x_1^3 + x_1 x_2^3 & c &= 3 - 3 \cdot \frac{2}{18} \\
 V^{(\mathrm{E}_8)} &= x_1^3 + x_2^5 & c &= 3 - 3 \cdot \frac{2}{30}
 \end{aligned} \tag{1.5}$$

²As we shall see in Section 2.3 we can also work over the cyclotomic field $k = \mathbb{Q}(\zeta)$ for an appropriate root of unity ζ . This will allow us to relate non-isomorphic matrix factorisations exhibiting a given orbifold equivalence via the Galois group, cf. Remarks 2.10, 2.12 and 2.13.

From Proposition 1.3 we know that for two potentials to be orbifold equivalent, their central charges have to agree. The preimages of the central charge function on the potentials in (1.5) are precisely the sets

$$\begin{aligned} & \{V^{(A_{d-1})}\} \quad \text{for } d \text{ odd,} \\ & \{V^{(A_{d-1})}, V^{(D_{d/2+1})}\} \quad \text{for } d \text{ even and } d \notin \{12, 18, 30\}, \\ & \{V^{(A_{11})}, V^{(D_7)}, V^{(E_6)}\}, \quad \{V^{(A_{17})}, V^{(D_{10})}, V^{(E_7)}\}, \quad \{V^{(A_{29})}, V^{(D_{16})}, V^{(E_8)}\}. \end{aligned} \quad (1.6)$$

In fact, this list exhausts the relevant orbifold equivalences:

Theorem 1.5. The orbifold equivalence classes of the potentials (1.5) are precisely those listed in (1.6).

This is our main result (together with Corollaries 1.6 and 1.7 below), which we prove in Section 2.3 by explicitly constructing graded matrix factorisations X with non-zero quantum dimensions for the equivalences $V^{(A_{2d-1})} \sim V^{(D_{d+1})}$ (already given in [CR3]), as well as $V^{(A_{11})} \sim V^{(E_6)}$, $V^{(A_{17})} \sim V^{(E_7)}$ and $V^{(A_{29})} \sim V^{(E_8)}$.

Higher dimensional ADE singularities are obtained by adding squares of further variables to the potentials in (1.5). Thus it immediately follows from Proposition 1.2(i) that Theorem 1.5 describes orbifold equivalences between general ADE singularities in any dimension.

Proposition 1.4 and Theorem 1.5 can be strengthened to equivalences of categories by invoking the general theory of equivariant completion of [CR3]:

Corollary 1.6. For V, W and X as in Proposition 1.4 we have

$$\text{hmf}^{\text{gr}}(k[y], W) \cong \text{mod}(X^\dagger \otimes X). \quad (1.7)$$

Here $\text{mod}(X^\dagger \otimes X)$ is the *category of modules over* $X^\dagger \otimes X$, which is made up of matrix factorisations of V together with a compatible action of the monoid $X^\dagger \otimes X \in \text{hmf}^{\text{gr}}(k[x, x'], V(x) - V(x'))$. For more details we refer to [CR3], or to [BCP, Car] for much shorter reviews.

In Section 2.3 we will explicitly compute $X^\dagger \otimes X$ for the matrix factorisations X giving rise to the orbifold equivalences of simple singularities. For those involving E-type singularities we find that $X^\dagger \otimes X$ decomposes into sums of the so-called permutation matrix factorisations P_S of $u'^d - u^d$ introduced in [BR1]. For a subset $S \subset \mathbb{Z}_d$ we denote by S^c its complement in \mathbb{Z}_d . Then one has $P_S = (\mathbb{C}[u, u'] \oplus \mathbb{C}[u, u'], d_{P_S})$ with

$$d_{P_S}^1 = \prod_{l \in S} (u' - \zeta_d^l u), \quad d_{P_S}^0 = \prod_{l \in S^c} (u' - \zeta_d^l u), \quad \text{where } \zeta_d = e^{2\pi i/d}. \quad (1.8)$$

Note that $P_{\{0\}}$ is nothing but the unit matrix factorisation. As shown in [BR1, DRCR] the tensor product in the case of consecutive index sets is

$$P_{\{m, m+1, \dots, m+a\}} \otimes P_{\{n, n+1, \dots, n+b\}} \cong \bigoplus_{i=|a-b|, \text{ step } 2}^{\min(a+b, 2d-4-a-b)} P_{\{m+n-\frac{1}{2}(a+b-i), \dots, m+n-\frac{1}{2}(a+b-3i)\}} \quad (1.9)$$

and the quantum dimensions were computed in [CR2, Sect. 3.3] to be

$$\dim_l(P_S) = \sum_{l \in S} \zeta_d^l, \quad \dim_r(P_S) = \sum_{l \in S} \zeta_d^{-l}. \quad (1.10)$$

Corollary 1.7. We have

$$\begin{aligned} \text{hmf}^{\text{gr}}(\mathbb{C}[x], V^{(E_6)}) &\cong \text{mod}(P_{\{0\}} \oplus P_{\{-3, -2, \dots, 3\}}), \\ \text{hmf}^{\text{gr}}(\mathbb{C}[x], V^{(E_7)}) &\cong \text{mod}(P_{\{0\}} \oplus P_{\{-4, -3, \dots, 4\}} \oplus P_{\{-8, -7, \dots, 8\}}), \\ \text{hmf}^{\text{gr}}(\mathbb{C}[x], V^{(E_8)}) &\cong \text{mod}(P_{\{0\}} \oplus P_{\{-5, -4, \dots, 5\}} \oplus P_{\{-9, -8, \dots, 9\}} \oplus P_{\{-14, -13, \dots, 14\}}). \end{aligned} \quad (1.11)$$

Remark 1.8. (i) The construction of the equivalence relation can be repeated in a number of slightly modified settings. For example, one can work with ungraded matrix factorisations, or one can work over any commutative ring k as in [CM2]. Or, instead of using a \mathbb{Q} -grading, one can consider matrix factorisations with R-charge in the sense of [CR2] (which is more general).

The setting used in this paper was chosen to be on the one hand as simple as possible – hence working over $k \subset \mathbb{C}$ – and on the other hand to be strong enough for us to be able to prove the decomposition of simple singularities into equivalence classes (1.6) – hence the homogeneous potentials and the \mathbb{Q} -grading. (In the ungraded setting we do not know how to exclude the existence of equivalences beyond those in (1.6).)

(ii) The decompositions (1.6) and (1.11) are expected from two-dimensional rational conformal field theory. According to the (conjectural) CFT/LG correspondence, the infrared fixed point of the Landau-Ginzburg model with potential V is a conformal field theory with central charge $c(V)$ [Mar, VW, HW]. Mathematically, this predicts a relation between the graded matrix factorisations of a given potential and the representation theory of a super vertex operator algebra. This relation is supported by explicit computations on both sides in examples but not understood (or even precisely formulated) in general.

For simple singularities, the fixed points are $\mathcal{N} = 2$ supersymmetric minimal models. The classification of the latter contains the above ADE series [Mar, VW, CV, Gan, Gra]. Matrix factorisations of the difference of two potentials are interpreted as line defects on the CFT side

[BR1, BR2, CR1, CR2]. It is known in rational CFT [FFRS1, FFRS2] that there are topological line defects with non-zero quantum dimensions, linking the minimal models in the equivalence classes corresponding to (1.6). We will compare this with our results in more detail in Section 3.

- (iii) It was shown in [KST] that for a simple singularity $V \in \mathbb{C}[x]$, $\text{hmf}^{\text{gr}}(\mathbb{C}[x], V)$ is equivalent to $\mathbb{D}^b(\text{Rep } \mathbb{C}Q)$, where Q is (any choice of) the associated Dynkin quiver. Thus by Theorem 1.5 and [CR3] the derived representation theory of ADE quivers enjoys orbifold equivalences analogous to (1.11).³ The monoids $X^\dagger \otimes X \cong P_{\{0\}} \oplus \dots$ translate into functors on $\mathbb{D}^b(\text{Rep } \mathbb{C}Q)$ whose actions on simple objects are easily computable.

Acknowledgements

We thank Hanno Becker, Ilka Brunner, Calin Lazaroiu, Wolfgang Lerche, Daniel Murfet, Daniel Plencner, Alexander Polishchuk, Daniel Roggenkamp, Eric Sharpe, Duco van Straten, Catharina Stroppel, and Atsushi Takahashi. In addition we are grateful to the organisers of the inspirational Oberwolfach workshop “Matrix Factorizations in Algebra, Geometry, and Physics”, without which numerous helpful discussions would not have taken place. We also acknowledge support from the German Science Foundation (DFG), N. C. and I. R. within the Collaborative Research Center 676 “Particles, Strings and the Early Universe”, A. R. C. as part of the Research Training Group 1670 “Mathematics Inspired by String Theory and QFT”.

2 Proofs

In this section we provide proofs of the results summarised above, with the claims of Sections 1.1 and 1.2 proven in Sections 2.2 and 2.3, respectively. The discussion for simple singularities can also be viewed as showcasing methods that may prove useful for constructing further orbifold equivalences between potentials.

The notion of orbifold equivalence actually makes sense between objects in any pivotal bicategory. For context and completeness this is explained in Section 2.1, after which we turn to concrete matrix factorisations.

2.1 Pivotal bicategories

In this section (only) we assume some familiarity with bicategories, referring to [Bor] for an introduction. We shall adopt the notation and conventions of [CR3]; more detailed explanations of the properties of duals and dimensions stated below can be found for example in [CR2, Sect. 2] and [CM2, Sect. 2.1].

³Such a relation between A- and D-type quivers was already proven in [RR] by different methods.

We denote the unit 1-morphism of an object a in a bicategory \mathcal{B} by I_a , its left action by λ , and the (horizontal) composition of 1-morphisms $X \in \mathcal{B}(a, b)$, $Y \in \mathcal{B}(b, c)$ is written $Y \otimes X \in \mathcal{B}(a, c)$.

We say that \mathcal{B} has *left adjoints* if for each $X \in \mathcal{B}(a, b)$ there is an $X^\dagger \in \mathcal{B}(b, a)$ together with 2-morphisms

$$\text{ev}_X : X^\dagger \otimes X \longrightarrow I_a, \quad \text{coev}_X : I_b \longrightarrow X \otimes X^\dagger \quad (2.1)$$

which are the counit and unit of an adjunction, see [GM]. Such a \mathcal{B} is *pivotal* if for all $a, b \in \mathcal{B}$ there are natural isomorphisms $\delta^{a,b}$ between the identity functor and $(-)^{\dagger\dagger}$ restricted to $\mathcal{B}(a, b)$, compatible with \otimes , see e.g. [Sch, Def. 5.1.4]. In this case X^\dagger is also right adjoint to X as exhibited by the maps

$$\tilde{\text{ev}}_X : X \otimes X^\dagger \longrightarrow I_b, \quad \widetilde{\text{coev}}_X : I_a \longrightarrow X^\dagger \otimes X \quad (2.2)$$

defined by $\text{ev}_{X^\dagger} \circ (\delta_X^{a,b} \otimes 1_{X^\dagger})$ and $(1_{X^\dagger} \otimes (\delta_X^{a,b})^{-1}) \circ \text{coev}_{X^\dagger}$, respectively.

From now on we assume that we are given a pivotal bicategory \mathcal{B} . For every $X \in \mathcal{B}(a, b)$ its *left* and *right quantum dimensions* are defined as

$$\dim_l(X) = \text{ev}_X \circ \widetilde{\text{coev}}_X \in \text{End}(I_a), \quad \dim_r(X) = \tilde{\text{ev}}_X \circ \text{coev}_X \in \text{End}(I_b). \quad (2.3)$$

One can show that the quantum dimensions only depend on the isomorphism class of X , and that

$$\dim_l(X^\dagger) = \dim_r(X). \quad (2.4)$$

The left quantum dimension of X is the image of 1_{I_b} under the map $\mathcal{D}_l(X) : \text{End}(I_b) \rightarrow \text{End}(I_a)$ given by

$$\mathcal{D}_l(X) = \text{ev}_X \circ [1_{X^\dagger} \otimes (\lambda_X \circ ((-) \otimes 1_X) \circ \lambda_X^{-1})] \circ \widetilde{\text{coev}}_X, \quad (2.5)$$

and similarly $\dim_r(X) = \mathcal{D}_r(X)(1_{I_a})$. One finds that for $Y \in \mathcal{B}(b, c)$ these operators satisfy

$$\mathcal{D}_l(X) \circ \mathcal{D}_l(Y) = \mathcal{D}_l(Y \otimes X), \quad \mathcal{D}_r(Y) \circ \mathcal{D}_r(X) = \mathcal{D}_r(Y \otimes X). \quad (2.6)$$

In our application to matrix factorisations, \mathcal{B} is a pivotal k -linear bicategory in which all quantum dimensions are multiples of the identity. For such a \mathcal{B} the definition of orbifold equivalence takes the form:

Definition 2.1. Two objects $a, b \in \mathcal{B}$ are *orbifold equivalent*, $a \sim b$, if there is $X \in \mathcal{B}(a, b)$ whose left and right quantum dimensions are non-zero multiples of the identity.

The following fact (whose proof is contained in Remark 2.3) shows that the name makes sense:

Theorem 2.2. Orbifold equivalence is an equivalence relation.

Remark 2.3. One may also consider the following slightly more general notion of orbifold equivalence in any pivotal bicategory: $a, b \in \mathcal{B}$ are orbifold equivalent if there is $X \in \mathcal{B}(a, b)$ such that

- (i) X has invertible left and right quantum dimensions, and
- (ii) $\mathcal{D}_l(X)$ and $\mathcal{D}_r(X)$ map invertible quantum dimensions to automorphisms.

Condition (ii) is needed for orbifold equivalence in this general setting to be an equivalence relation. This condition may be hard to check directly, as it requires knowing all invertible elements in $\text{End}(I_a)$ and $\text{End}(I_b)$ which arise as quantum dimension of some 1-morphism. A special case where condition (ii) is already implied by condition (i) is if \mathcal{B} is such that every Hom-category $\mathcal{B}(a, b)$ is R -linear for some fixed commutative ring R , and such that the left and right quantum dimensions of every 1-morphism $X \in \mathcal{B}(a, b)$ are in $R \cdot 1_{I_a}$ and $R \cdot 1_{I_b}$, respectively. In this case one may think of the left/right quantum dimension as an element of R , and one can check that $\dim_{l/r}(X \otimes Y) = \dim_{l/r}(X) \cdot \dim_{l/r}(Y)$, where the product on the right is in R . Graded finite-rank matrix factorisations are an example of this R -linear setting (see Lemma 2.6 below).

Theorem 2.2 holds in the more general case: to see reflexivity $a \sim a$ one can take $X = I_a$, for which $\mathcal{D}_l(I_a) = 1_{I_a} = \mathcal{D}_r(I_a)$. For symmetry and transitivity it is helpful to expand condition (ii) of the equivalence relation in more detail:

- (ii-r) for every $e \in \mathcal{B}$ and every $Z \in \mathcal{B}(e, a)$ such that $\dim_r(Z)$ is invertible in $\text{End}(I_a)$, we have that $\mathcal{D}_r(X)(\dim_r(Z))$ is invertible in $\text{End}(I_b)$;
- (ii-l) for every $f \in \mathcal{B}$ and every $Z' \in \mathcal{B}(b, f)$ such that $\dim_l(Z')$ is invertible in $\text{End}(I_b)$, we have that $\mathcal{D}_l(X)(\dim_l(Z'))$ is invertible in $\text{End}(I_a)$.

Of course, since (ii-r) holds for all Z , it also holds for all Z^\dagger , and hence $\mathcal{D}_r(X)$ also maps invertible left quantum dimensions to automorphisms. The same applies to (ii-l).

Symmetry $a \sim b \Leftrightarrow b \sim a$ follows by replacing X by X^\dagger . That X^\dagger satisfies (i) follows from (2.4). To see (ii-r) for X^\dagger , use $\mathcal{D}_r(X^\dagger) = \mathcal{D}_l(X)$ and that (ii-l) holds for X . Condition (ii-l) follows analogously.

To check transitivity, consider $X \in \mathcal{B}(a, b)$ and $Y \in \mathcal{B}(b, c)$ satisfying conditions (i) and (ii). Then $\dim_r(Y \otimes X) = \mathcal{D}_r(Y)(\dim_r(X))$ which is invertible by (i) for X and (ii-r) for Y . Dito for \dim_l . Next let $Z \in \mathcal{B}(e, a)$ be as in (ii-r) above. Then $\mathcal{D}_r(Y \otimes X)(\dim_r(Z)) = \mathcal{D}_r(Y)(\dim_r(X \otimes Z))$. Now $\dim_r(X \otimes Z)$ is invertible by (ii-r) for X and therefore $\mathcal{D}_r(Y)(\dim_r(X \otimes Z))$ is invertible by (ii-r) for Y . That $Y \otimes X$ satisfies (ii-l) is seen similarly.

We conclude our brief general discussion with an implication of the existence of orbifold equivalences in the setting of retracts and idempotent splittings. An object S in some category is called a *retract* of an object U if there are morphisms $e : S \rightarrow U$ and $r : U \rightarrow S$ such that $r \circ e = 1_S$. In particular, e is mono and so S is a subobject of U .

Proposition 2.4. Let $a, b, c, d \in \mathcal{B}$ and let $X \in \mathcal{B}(a, b)$, $Y \in \mathcal{B}(c, d)$ have invertible left quantum dimensions. Every $Z \in \mathcal{B}(a, c)$ is a retract of $Y^\dagger \otimes F \otimes X$ for some suitable (Z -dependent) $F \in \mathcal{B}(b, d)$.

Proof. We set $F = Y \otimes Z \otimes X^\dagger$ and define the maps $e : Z \rightarrow Y^\dagger \otimes F \otimes X$ and $r : Y^\dagger \otimes F \otimes X \rightarrow Z$ by

$$\begin{aligned} e &= \widetilde{\text{coev}}_Y \otimes 1_Z \otimes \widetilde{\text{coev}}_X, \\ r &= (\dim_1(Y)^{-1} \otimes 1_Z \otimes \dim_1(X)^{-1}) \circ (\text{ev}_Y \otimes 1_Z \otimes \text{ev}_X). \end{aligned} \quad (2.7)$$

Clearly, $r \circ e = 1_Z$. □

Remark 2.5. If the Hom-categories of \mathcal{B} are additive and if idempotent 2-morphisms split, we can improve on Proposition 2.4 slightly: instead of Z just being a retract, it now even occurs as a direct summand of $Y^\dagger \otimes F \otimes X$ for a suitable F . To see this, take the maps e, r from the proof, note that $p = e \circ r$ is an idempotent endomorphism of $Y^\dagger \otimes Y \otimes Z \otimes X^\dagger \otimes X$, and consider the decomposition $1 = p + (1 - p)$ into orthogonal idempotents.

2.2 Matrix factorisations

Potentials and (graded) matrix factorisations as defined in Section 1 form bicategories that are ‘as good as’ pivotal,⁴ as explained in detail in [CM2]. Accordingly we could present the results of this section as applications of a slight variant of the previous one. We will however take a different route and give more direct arguments.

Recall from [CR2, BFK, CM2] that every ungraded matrix factorisation X of $W(y_1, \dots, y_n) - V(x_1, \dots, x_m)$ has left and right adjoints. If we write $X^\vee = \text{Hom}_{k[x,y]}(X, k[x, y])$ and set $d_{X^\vee}(\nu) = (-1)^{|\nu|+1} \nu \circ d_X$ for \mathbb{Z}_2 -homogeneous $\nu \in X^\vee$, then they are given by

$${}^\dagger X = X^\vee[n], \quad X^\dagger = X^\vee[m], \quad (2.8)$$

respectively. These are (ungraded) matrix factorisations of $V - W$, and the associated adjunction maps as in (2.1), (2.2) are known explicitly, see [CM2] for details or [CM3] for a concise review. It follows that isolated singularities in an even number of variables and ungraded matrix factorisations form a pivotal bicategory $\mathcal{LG}^{\text{even}}$.

A straightforward computation of counting degrees shows that in the case of a graded matrix factorisation X its left and right adjunction maps of [CM2] are homogeneous of degree $\pm \sum_{j=1}^n (1 - |y_j|) = \pm \frac{1}{3}c(W)$ and $\pm \sum_{i=1}^m (1 - |x_i|) = \pm \frac{1}{3}c(V)$,

⁴One has pivotality ‘on the nose’ if one restricts to potentials in an even number of variables, as we do in our discussion of simple singularities in Section 2.3.

respectively, with respect to the adjoints in (2.8). Consequently the bicategory of potentials in an even number of variables, graded matrix factorisations, and their maps of arbitrary \mathbb{Q} -degree form a pivotal bicategory $\mathcal{LG}^{\text{gr,even}}$.

We remark that allowing maps of arbitrary \mathbb{Q} -degree is natural from the perspective of conformal field theory, where this degree corresponds to the R-charge. On the other hand, derived equivalences connecting graded matrix factorisations with representation theory [KST] or geometry [Buc, Orl] require restricting to the subcategory $\mathcal{LG}_0^{\text{gr,even}}$ with 2-morphisms only of degree zero. (Categories of graded matrix factorisations with maps of arbitrary degree are not even triangulated as mapping cones can fail to be graded matrix factorisations.) It follows that the full subcategories $\mathcal{LG}_0^{\text{gr,even}}(c) \subset \mathcal{LG}_0^{\text{gr,even}}$ of potentials with fixed central charge $c \in \mathbb{Q}$ are strictly pivotal if left and right adjoints in $\mathcal{LG}_0^{\text{gr}}$ are taken to be

$${}^\dagger X = X^\vee[n]\{\frac{1}{3}c(W)\}, \quad X^\dagger = X^\vee[m]\{\frac{1}{3}c(V)\}, \quad (2.9)$$

where $\{-\}$ denotes the \mathbb{Q} -grading shift.

Irrespective of the particular choice of bicategory, control over adjunctions leads to the explicit expressions for the quantum dimensions given in (1.3). For ungraded matrix factorisations these are polynomials, but in the graded case the quantum dimensions are just numbers:

Lemma 2.6. Quantum dimensions of graded matrix factorisations take values in k .

Proof. Let X be a graded matrix factorisation of $W - V$. Quantum dimensions are elements of the k -algebra of endomorphisms of the unit matrix factorisation of a potential. This algebra is isomorphic to the Milnor ring of the potential, and its only degree-zero component is k . Hence the statement follows for $\mathcal{LG}_0^{\text{gr}}$ by definition. For \mathcal{LG}^{gr} , $\dim_{1/r}(X)$ has degree $\pm \frac{1}{3}(c(W) - c(V))$, but by Proposition 1.3(ii) the quantum dimensions are non-zero only for $c(V) = c(W)$. \square

As a consequence of this lemma, in the graded case the statements “ X has non-zero quantum dimensions” and “ X has invertible quantum dimensions” are equivalent.

We may now proceed to prove the results advertised in Section 1.1.

Proof of Theorem 1.1. This is a corollary of [CM2, Prop.8.5]: reflexivity and symmetry follow analogously to Theorem 2.2, and transitivity follows from the fact that the quantum dimensions (1.3) are manifestly multiplicative up to a sign. \square

Proof of Proposition 1.2. From (1.3) it is clear that quantum dimensions are multiplicative (up to a sign) also for the external tensor product \otimes_k , showing part (i). Part (ii) is proven in [CR3, Sect. 7.2]. \square

Proof of Proposition 1.4. This is a direct consequence of the proof of Proposition 2.4 and Remark 2.5 as we are dealing with idempotent complete matrix factorisation categories. \square

2.3 Simple singularities

Let $W(y_1, y_2)$ be one of the potentials $V^{(D_{d/2+1})}, V^{(E_6)}, V^{(E_7)}, V^{(E_8)}$ in (1.5), and let $V(x_1, x_2) = V^{(A_{d-1})} = x_1^d + x_2^2$ be the corresponding A-type potential of the same central charge as W . To avoid too many indices, we rename

$$x_1 \rightsquigarrow u, \quad x_2 \rightsquigarrow v, \quad y_1 \rightsquigarrow x, \quad y_2 \rightsquigarrow y. \quad (2.10)$$

In this section we will give a finite-rank graded matrix factorisation of $W(x, y) - V(u, v)$ of non-zero left and right quantum dimension in each case, thus proving the results collected in Section 1.2. These matrix factorisations have all been constructed along the following lines:

- (i) Pick a matrix factorisation X_0 of $W(x, y) - v^2$ that is of low rank.
- (ii) Thinking of u as a deformation parameter, add to each entry of the matrix d_{X_0} the most general homogeneous polynomial of the form $u \cdot p(u, v, x, y)$ with the same total degree as the given entry. Let d_{X_u} be the resulting matrix.
- (iii) Reduce the number of free parameters in the polynomials p by absorbing some of them via a similarity transformation $d_{X_u} \mapsto \Phi \circ d_{X_u} \circ \Phi^{-1}$.
- (iv) Try to find a set of parameters (the remaining coefficients in the polynomials p) such that $d_{X_u} \circ d_{X_u} = (W - V) \cdot 1$.

Choosing a low-rank starting point in (i) and reducing the number of parameters via (iii) only serves to simplify the problem in (iv). We will work through these steps for $W = V^{(D_{d/2+1})}$ and $W = V^{(E_6)}$ in some detail, while our discussion will be briefer for $V^{(E_7)}$ and $V^{(E_8)}$.

$$\mathbf{V}^{(D_{d/2+1})} \sim \mathbf{V}^{(A_{d-1})}$$

We set $b = d/2 \geq 3$, so that $W = x^b + xy^2$ and $V = u^{2b} + v^2$, and we write $R = \mathbb{C}[u, v, x, y]$. As the starting point in step (i) we choose $X_0 = (X, d_{X_0})$ with \mathbb{Z}_2 -graded R -module $X = R^2 \oplus R^2$ and twisted differential

$$d_{X_0} = \begin{pmatrix} 0 & d_{X_0}^1 \\ d_{X_0}^0 & 0 \end{pmatrix} \quad \text{with} \quad d_{X_0}^1 = \begin{pmatrix} x & v \\ v & x^{b-1} + y^2 \end{pmatrix}. \quad (2.11)$$

Since $\det(d_{X_0}^1) = x^b + xy^2 - v^2$, the component $d_{X_0}^0$ is determined to be the adjunct matrix, $d_{X_0}^0 = (d_{X_0}^1)^\#$. Recall that the adjunct $M^\#$ of an invertible matrix M has entries which are polynomial in those of M and satisfies $M^{-1} = (\det M)^{-1} \cdot M^\#$.

The deformed matrix factorisation X_u has the same underlying R -module $R^2 \oplus R^2$. For step (ii) we need to pick the most general homogeneous deformation of $d_{X_0}^1$, which is

$$d_{X_u}^1 = \begin{pmatrix} x + u p_{11} & v + u p_{12} \\ v + u p_{21} & x^{b-1} + y^2 + u p_{22} \end{pmatrix} \quad \text{where } p_{ij} \in \mathbb{C}[u, v, x, y]. \quad (2.12)$$

The degrees of the variables are $|u| = 1/b$, $|v| = 1$, $|x| = 2/b$, and $|y| = 1 - 1/b$. Hence the total degrees of the p_{ij} have to be

$$|p_{11}| = \frac{1}{b}, \quad |p_{12}| = 1 - \frac{1}{b}, \quad |p_{21}| = 1 - \frac{1}{b}, \quad |p_{22}| = 2 - \frac{3}{b}. \quad (2.13)$$

From this it follows that $p_{11} = a_1 u$ for some $a_1 \in \mathbb{C}$, but the remaining p_{ij} will contain of the order b many free coefficients.

Moving to step (iii), we will now use degree-preserving row and column operations on $d_{X_u}^1$ to reduce the number of free coefficients. By applying the inverse operations to $d_{X_u}^0$ one produces in this way an isomorphic matrix factorisation. The most general such row and column manipulations turn out to be

$$\begin{pmatrix} 1 & 0 \\ f & 1 \end{pmatrix} d_{X_u}^1 \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} x + a_1 u^2 & v + u p_{12} + g(x + a_1 u^2) \\ v + u p_{21} + f(x + a_1 u^2) & * \end{pmatrix} \quad (2.14)$$

where $|f| = |g| = 1 - 2/b$. We see that f, g can be used to remove any x -dependence from p_{12} and p_{21} , and we arrive at the following reduced ansatz: $d_{X_u}^1$ of the form (2.12) with

$$\begin{aligned} p_{11} &= a_1 u, & p_{12} &= a_2 y + a_3 u^{b-1}, \\ p_{21} &= a_4 y + a_5 u^{b-1}, & p_{22} &= q_1 v + q_2 y + q_3, \end{aligned} \quad (2.15)$$

where $a_i \in \mathbb{C}$ and $q_i \in \mathbb{C}[u, x]$ with degrees $|q_1| = 1 - 3/b$, $|q_2| = 1 - 2/b$, and $|q_3| = 2 - 3/b$.

Step (iv) amounts to the tedious task of trying to find conditions such that $d_{X_u} \circ d_{X_u} = (W - V) \cdot 1$. The second component of the twisted differential is uniquely determined to be $d_{X_u}^0 = q / \det(d_{X_u}^1) \cdot (d_{X_u}^1)^\#$ with $q = x^b + x y^2 - u^{2b} - v^2$, and we need to find values of the deformation parameters so that this matrix has polynomial entries. Since $q \in \mathbb{C}[x, y, u, v]$ is irreducible, either q is a factor of $\det(d_{X_u}^1)$, or $\det(d_{X_u}^1)$ has to cancel against the entries of $(d_{X_u}^1)^\#$. Degree considerations show that the latter is not possible, and in fact $\det(d_{X_u}^1)$ equals q up to a multiplicative constant. By rescaling $d_{X_u}^1$ if necessary, without restriction of generality we can impose $\det(d_{X_u}^1) = q$. In solving this condition, one is lead to distinguish between two cases, $a_2 = 0$ and $a_2 \neq 0$. Setting $a_2 = 0$ produces a solution with zero left and right quantum dimension. On the other hand, keeping $s := a_2 \neq 0$ forces

$$d_{X_u}^1 = \begin{pmatrix} x - (su)^2 & v + y(su) \\ v - y(su) & \frac{x^b - (su)^{2b}}{x - (su)^2} + y^2 \end{pmatrix} \quad \text{where } s^{2b} = 1. \quad (2.16)$$

For $s = 1$ this is the solution given in [CR3, Sect. 7.3]. The quantum dimensions are

$$\dim_l(X_u) = -2s, \quad \dim_r(X_u) = -s^{-1}. \quad (2.17)$$

What remains to be done is to check that there exists a \mathbb{Q} -grading on $X_u = (X, d_{X_u})$ such that d_{X_u} has \mathbb{Q} -degree 1. It is in fact easy to write down all such gradings. The \mathbb{Q} -grading on the ring R is fixed by $|1| = 0$, with the variables u, v, x, y having degrees as stated below (2.12). Writing $R[\alpha]$, $\alpha \in \mathbb{Q}$, for R with $|1| = \alpha$, the possible gradings on X are

$$X^0 = R[\alpha] \oplus R[\alpha - 1 + \frac{2}{b}], \quad X^1 = R[\alpha - 1 + \frac{2}{b}] \oplus R[\alpha], \quad \alpha \in \mathbb{Q}. \quad (2.18)$$

Equivalently, the grading data on X may be presented as a ‘ U -matrix’, cf. e. g. [CR2],

$$U_X(\beta) = e^{i\alpha\beta} \cdot \text{diag} \left(1, e^{(2/b-1)i\beta}, e^{(2/b-1)i\beta}, 1 \right). \quad (2.19)$$

This completes the proof that the potentials $V^{(A_{d-1})}$ and $V^{(D_{d/2+1})}$ are orbifold equivalent. We can summarise the above discussion as follows.

Lemma 2.7. Let Y be a rank-two \mathbb{Q} -graded matrix factorisation of $x^b + xy^2 - u^{2b} - v^2$. Suppose that (i) Y has non-zero quantum dimensions and (ii) when setting u to zero, Y is equal to X_0 in (2.11). Then Y is isomorphic in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, v, x, y], x^b + xy^2 - u^{2b} - v^2)$ to X_u in (2.16) for some choice of s .

Let us analyse the matrix factorisation X_u a bit further. The automorphism σ of $\mathbb{C}[u]$ determined by $\sigma(u) = \zeta_d u$ leaves the potential u^d invariant. Given a matrix factorisation X of $W - u^d$ for some $W \in \mathbb{C}[x]$, twisting the $\mathbb{C}[u, x]$ -action on X by σ^l (with σ extended to $\mathbb{C}[u, x]$ via $\sigma(x_i) = x_i$) results again in a matrix factorisation of $W - u^d$ which we denote by X_{σ^l} . It follows from [CR1, Lem. 2.10] that

$$X_{\sigma^l} \cong X \otimes P_{\{-l\}} \quad (2.20)$$

in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, x], W - u^d)$, where $P_{\{-l\}}$ is a permutation matrix factorisation as in (1.10).

Remark 2.8. The observation (2.20) together with the classification given in Lemma 2.7 explains the $2b$ -th root of unity s appearing as a parameter in (2.16). To wit, given any 1-morphism $P : V \rightarrow V$ of non-zero quantum dimensions, the composition $X_u \otimes P : V \rightarrow W$ also has non-zero quantum dimensions. Consider the choice $P = P_{\{l\}} \otimes_{\mathbb{C}} I_{v^2}$, where I_{v^2} the unit 1-morphism for the potential v^2 with twisted differential $\begin{pmatrix} 0 & v'-v \\ v'+v & 0 \end{pmatrix}$. Then P is a 1-endomorphism of $u^{2b} + v^2$ and $X_u \otimes P$ is again a rank-two factorisation satisfying the conditions in Lemma 2.7. Hence $X_u \otimes P$ must be isomorphic to X_u for a possibly different choice of s . But different choices of s precisely amount to twisting the u -action by some power of the automorphism σ .

It was checked in [CR3, Sect. 7.3] that for $d \in \{2, 3, \dots, 10\}$, and up to a trivial factor of the unit I_{v^2} , the monoid $X_{u'}^\dagger \otimes X_u$ is isomorphic to

$$P_{\{0\}} \oplus P_{\{0,1,\dots,d-1\} \setminus \{\frac{d}{2}\}}. \quad (2.21)$$

Since $P_{\{0\}} = I_{u^{2d}}$ and $P_{\{0,1,\dots,d-1\} \setminus \{\frac{d}{2}\}} \cong P_{\{\frac{d}{2}\}}[1]$ acts as $u \mapsto -u$ via tensoring, the matrix factorisation (2.21) precisely encodes a \mathbb{Z}_2 -action, which also uniquely fixes its monoid structure. This \mathbb{Z}_2 -action was first discussed in the context of Dynkin quivers in [RR]. We conjecture that (2.21) tensored with I_{v^2} is isomorphic to $X_{u'}^\dagger \otimes X_u$ for all $d \in \mathbb{Z}_{\geq 2}$.

It is straightforward to compute an explicit basis for the endomorphisms of X_u in the homotopy category, e. g. by using the Singular code of [CDR]. We have done so for $d \in \{2, 3, \dots, 10\}$, with the result

$$\text{End}(X_u) \cong \left(\bigoplus_{i=0}^{d-2} \mathbb{C}[i \cdot \frac{2}{d}] \right) \oplus \mathbb{C}[\frac{d-2}{d}] \quad (2.22)$$

where as before $\mathbb{C}[j]$ denotes the one-dimensional subspace of maps of \mathbb{Q} -degree j . We expect (2.22) to be true for all d .

$$\mathbf{V}^{(\mathbf{E}_6)} \sim \mathbf{V}^{(\mathbf{A}_{11})}$$

Our starting point in step (i) now is the matrix factorisation $X_0 = (R^2 \oplus R^2, d_{X_0})$ of $x^3 + y^4 - v^2$ with $R = \mathbb{C}[u, v, x, y]$ and twisted differential

$$d_{X_0}^1 = \begin{pmatrix} y^2 - v & -x \\ x^2 & y^2 + v \end{pmatrix}, \quad d_{X_0}^0 = (d_{X_0}^1)^\# . \quad (2.23)$$

This is one of the six indecomposable objects of $\text{hmf}^{\text{gr}}(\mathbb{C}[v, x, y], V^{(\mathbf{E}_6)} - v^2)$ listed in [KST, Sect. 5]. The variable degrees are

$$|u| = \frac{1}{6}, \quad |v| = \frac{6}{6}, \quad |y| = \frac{3}{6}, \quad |x| = \frac{4}{6}. \quad (2.24)$$

Carrying out steps (ii) and (iii) leads to the possibility

$$d_{X_u}^1 = \begin{pmatrix} y^2 - v + a_1xu^2 + a_2u^6 & -x + a_3yu + a_4u^4 \\ x^2 + a_5yxu + a_6xu^4 + a_7vu^2 + q & y^2 + v + a_8xu^2 + a_9u^6 \end{pmatrix} \quad (2.25)$$

for the reduced ansatz, where $a_i \in \mathbb{C}$ and $q \in \mathbb{C}[u, y]$ with $|q| = 8/6$. Of course a different choice of similarity transformation may give a different (but isomorphic) reduced ansatz. Here, the row and column manipulations were used to absorb the terms yu^3 in the diagonal entries.

By the same argument as used in the D-case, we now need to solve the condition $\det(d_{X_u}^1) = x^3 + y^4 - u^{12} - v^2$ under the extra constraint that X_u has non-zero quantum dimensions. This leads to

$$d_{X_u}^1 = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{with} \quad a = y^2 - v + \frac{1}{2}x(su)^2 + \frac{2t+1}{8}(su)^6,$$

$$\begin{aligned}
b &= -x + y(su) + \frac{t+1}{4}(su)^4, \\
c &= x^2 + yx(su) + \frac{t}{4}x(su)^4 + \frac{2t+1}{4}y(su)^5 - \frac{9t+5}{48}(su)^8, \\
d &= y^2 + v + \frac{1}{2}x(su)^2 + \frac{2t+1}{8}(su)^6,
\end{aligned} \tag{2.26}$$

and $d_{X_0}^0 = (d_{X_0}^1)^\#$. Here, s and t can be any solution of

$$t^2 = \frac{1}{3}, \quad s^{12} = -576(26t - 15). \tag{2.27}$$

Note that s can be modified by a 12-th root of unity – the interpretation of this is as in Remark 2.8.

The quantum dimensions of X_u are

$$\dim_l(X_u) = s, \quad \dim_r(X_u) = 3(1-t)s^{-1}, \tag{2.28}$$

and all possible \mathbb{Q} -gradings on X_u are again easily found: the underlying \mathbb{Z}_2 -graded R -module $X = X_0 \oplus X_1$ has components with \mathbb{Q} -grading

$$X^0 = R[\alpha] \oplus R[\alpha - \frac{1}{3}], \quad X^1 = R[\alpha] \oplus R[\alpha - \frac{1}{3}], \quad \alpha \in \mathbb{Q}, \tag{2.29}$$

equivalently described by the U -matrix

$$U_{X_u}(\beta) = e^{i\alpha\beta} \cdot \text{diag}(1, e^{-i\beta/3}, 1, e^{-i\beta/3}). \tag{2.30}$$

As in the D-case, we can summarise the above as:

Lemma 2.9. Let Y be a rank-two \mathbb{Q} -graded matrix factorisation of $x^3 + y^4 - u^{12} - v^2$. Suppose that (i) Y has non-zero quantum dimensions and (ii) when setting u to zero, Y is equal to X_0 in (2.23). Then Y is isomorphic in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, v, x, y], x^3 + y^4 - u^{12} - v^2)$ to X_u in (2.26) for some solution s, t of (2.27).

Computing the endomorphism of X_u in the homotopy category one finds the 16-dimensional space

$$\text{End}(X_u) \cong \left(\bigoplus_{i=0}^{10} \mathbb{C}[i \cdot \frac{2}{12}] \right) \oplus \left(\bigoplus_{i=3}^7 \mathbb{C}[i \cdot \frac{2}{12}] \right). \tag{2.31}$$

As mentioned in Section 1.2 and explained in detail in [CR3], a matrix factorisation X of $W(y) - V(x)$ with invertible quantum dimensions allows us to describe *all* matrix factorisations of W in terms of modules over $A := X^\dagger \otimes X \in \text{hmf}^{\text{gr}}(k[x, x'], V(x) - V(x'))$. The matrix $d_{X^\dagger \otimes X}$ also depends on the y -variables, and hence the matrix factorisation A is of infinite rank. However, by the results of [DM] it is homotopy equivalent (and thus isomorphic in $\text{hmf}^{\text{gr}}(k[x, x'], V(x) - V(x'))$) to a finite-rank matrix factorisation.

The construction of this finite-rank factorisation and the explicit homotopy equivalence can be implemented on a computer; this was done in [CM1], where it

was used to compute Khovanov-Rozansky link invariants. In our present situation we can use this implementation to find that $X_u^\dagger \otimes X_u$ is equivalent to the matrix factorisation $A' \otimes_k I_{v^2}$, where the twisted differential of A' is represented by a 4-by-4 matrix a' with nonzero entries

$$\begin{aligned}
a'_{13} &= \left(\frac{76}{3} - 44t\right)u^8 + \left(\frac{136}{3} - 80t\right)u'u^7 + (44 - 76t)u'^2u^6 + \left(\frac{152}{3} - 88t\right)u'^3u^5 \\
&\quad + \left(\frac{208}{3} - 120t\right)u'^4u^4 + \left(\frac{152}{3} - 88t\right)u'^5u^3 + (20t - 12)u'^6u^2 \\
&\quad + (88t - \frac{152}{3})u'^7u + (52t - \frac{92}{3})u'^8, \\
a'_{14} &= (224 - 384t)u^5 + (832 - 1440t)u'u^4 + (1440 - 2496t)u'^2u^3 \\
&\quad + (1440 - 2496t)u'^3u^2 + (832 - 1440t)u'^4u + (224 - 384t)u'^5, \\
a'_{23} &= \left(\frac{5}{3} - \frac{19t}{6}\right)u^{11} + \left(\frac{49}{6} - \frac{43t}{3}\right)u'u^{10} + \left(\frac{287}{18} - 28t\right)u'^2u^9 + \left(\frac{361}{18} - 35t\right)u'^3u^8 \\
&\quad + \left(\frac{208}{9} - \frac{241t}{6}\right)u'^4u^7 + \left(\frac{475}{18} - 46t\right)u'^5u^6 + \left(\frac{64}{3} - \frac{223t}{6}\right)u'^6u^5 + \left(\frac{23}{6} - 7t\right)u'^7u^4 \\
&\quad + \left(\frac{65t}{3} - \frac{227}{18}\right)u'^8u^3 + \left(\frac{74t}{3} - \frac{259}{18}\right)u'^9u^2 + \left(\frac{65t}{6} - \frac{58}{9}\right)u'^{10}u + \left(\frac{5t}{3} - \frac{19}{18}\right)u'^{11}, \\
a'_{24} &= (16 - 28t)u^8 + (104 - 180t)u'u^7 + \left(\frac{908}{3} - 524t\right)u'^2u^6 + (524 - 908t)u'^3u^5 \\
&\quad + \left(\frac{1780}{3} - 1028t\right)u'^4u^4 + (448 - 776t)u'^5u^3 + \left(\frac{652}{3} - 376t\right)u'^6u^2 \\
&\quad + (60 - 104t)u'^7u + \left(\frac{20}{3} - 12t\right)u'^8, \\
a'_{31} &= \frac{3tu^4}{4} - \frac{3u'u^3}{4} + \frac{u'^2u^2}{4} + \frac{3u'^3u}{4} + \left(-\frac{3t}{4} - \frac{1}{4}\right)u'^4, \\
a'_{32} &= 6u - 6u', \\
a'_{41} &= \left(-\frac{13t}{32} - \frac{3}{16}\right)u^7 + \left(\frac{7t}{32} + \frac{1}{8}\right)u'u^6 + \left(-\frac{13t}{32} - \frac{31}{96}\right)u'^2u^5 + \left(\frac{3t}{32} + \frac{19}{96}\right)u'^3u^4 \\
&\quad + \left(\frac{t}{8} + \frac{1}{48}\right)u'^4u^3 + \left(\frac{t}{16} - \frac{1}{12}\right)u'^5u^2 + \left(\frac{t}{8} + \frac{11}{96}\right)u'^6u + \left(\frac{3t}{16} + \frac{13}{96}\right)u'^7, \\
a'_{42} &= \left(\frac{3t}{4} - \frac{1}{4}\right)u^4 + 2u'u^3 - 3u'^2u^2 + \frac{5u'^3u}{2} + \left(-\frac{3t}{4} - \frac{5}{4}\right)u'^4. \tag{2.32}
\end{aligned}$$

This is a graded matrix factorisation when accompanied by the U -matrix

$$U_{A'}(\beta) = \text{diag}\left(1, e^{-i\beta/2}, e^{i\beta/3}, e^{-i\beta/6}\right). \tag{2.33}$$

In anticipation of the comparison to conformal field theory in Section 3 we single out one of the roots of (2.27),

$$t_{\text{cft}} = -1/\sqrt{3}. \tag{2.34}$$

Choosing $t = t_{\text{cft}}$ and performing a series of row and column manipulations produces the isomorphism

$$\begin{pmatrix} \phi & 6 & 0 & 0 \\ -\frac{1}{32}(12t+7) & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \psi & 1 \end{pmatrix} : A' \longrightarrow P_{\{0\}} \oplus P_{\{-3,-2,\dots,3\}} \tag{2.35}$$

where $\phi = \frac{1}{4}(u+u')(3tu^2 - 3uu' + u'^2 + 3tu'^2)$ and $\psi = \frac{1}{24}(u^3 - 3tu^3 - 7u^2u' - 3t^2u' + 5uu'^2 - 3tuu'^2 - 5u'^3 - 3tu'^3)$. This proves the first third of Corollary 1.7.

Remark 2.10. One may wonder what the monoid $X_{u'}^\dagger \otimes X_u$ reduces to for the other solution $t = 1/\sqrt{3}$ of (2.27), i. e. what is the equivalent to the right-hand side of (2.35). As we will see momentarily, the other solution can be related to t_{cft} via the action of an appropriate Galois group.

Define $\zeta_d = e^{2\pi i/d}$ and consider the cyclotomic field $k = \mathbb{Q}(\zeta_d)$, i. e. the field obtained from \mathbb{Q} by adjoining the d -th primitive root of unity ζ_d . The Galois group is isomorphic to the group of units in \mathbb{Z}_d , $\text{Gal}(k/\mathbb{Q}) \cong \mathbb{Z}_d^\times$. Given $\nu \in \mathbb{Z}_d^\times$ the action of the corresponding Galois group element σ_ν is $\sigma_\nu(\zeta_d^a) = \zeta_d^{\nu a}$.

Let now $V \in \mathbb{Q}[x]$ be a potential with rational coefficients and let M be a finite-rank matrix factorisation of V over $k[x]$. We may take $M = (k[x]^{2r}, d_M)$, where d_M is a matrix with entries in $k[x]$. Let $\sigma \in \text{Gal}(k/\mathbb{Q})$ be an element of the Galois group and denote by $\sigma(d_M)$ the matrix obtained by applying σ to each entry. Since $\sigma(V) = V$, $\sigma(d_M)$ is still a factorisation of V , and we set $\sigma(M) = (k[x]^{2r}, \sigma(d_M))$. Analogously, if $f : M \rightarrow N$ is a morphism with entries in $k[x]$, then $\sigma(f)$ is a morphism from $\sigma(M)$ to $\sigma(N)$.

Let us apply this to the isomorphism in (2.35). Write $A'(t)$ for A' to highlight the t -dependence. We choose $k = \mathbb{Q}(\zeta_{12})$ so that all P_S are matrix factorisations over $k[u, u']$. Since

$$t_{\text{cft}} = -\frac{1}{3}(\zeta_{12} + \zeta_{12}^{-1}), \quad (2.36)$$

also $A'(t_{\text{cft}})$ has entries with coefficients in k . The same holds for the isomorphism (2.35), and so we get an isomorphism from $\sigma(A'(t_{\text{cft}}))$ to $\sigma(P_{\{0\}} \oplus P_{\{-3, -2, \dots, 3\}})$. Since the entries of $A'(t)$ are polynomials in u, u', t with rational coefficients, we have $\sigma(A'(t_{\text{cft}})) = A'(\sigma(t_{\text{cft}}))$, and $\sigma(P_S) = P_{\sigma_*(S)}$, where σ_* is the permutation of \mathbb{Z}_{12} induced by the action of σ on the 12-th roots of unity.

It turns out that the orbit of t_{cft} under $\text{Gal}(k/\mathbb{Q})$ covers all, namely both, roots of (2.27): we have $\sigma_5(t_{\text{cft}}) = -\frac{1}{3}(\zeta_{12}^5 + \zeta_{12}^{-5}) = 1/\sqrt{3}$ and $\sigma_{5*}(\{-3, -2, \dots, 3\}) = \{-5, -3, -2, 0, 2, 3, 5\}$ and so we obtain the isomorphism

$$X_{u'}^\dagger \otimes X_u \cong (P_{\{0\}} \oplus P_{\{-5, -3, -2, 0, 2, 3, 5\}}) \otimes_k I_{v^2} \quad \text{for } t = \sigma_5(t_{\text{cft}}). \quad (2.37)$$

Note that by construction $A'(t_{\text{cft}})$ and $A'(\sigma_5(t_{\text{cft}}))$ are Morita equivalent, i. e. the category $\text{mod}(X_{u'}^\dagger \otimes X_u)$ does not depend on the choice of solution t . The situation is analogous for the E_7 - and E_8 -singularities, cf. Remarks 2.12 and 2.13.

$\mathbf{V}(\mathbf{E}_7) \sim \mathbf{V}(\mathbf{A}_{17})$

In step (i) we pick the matrix factorisation $X_0 = (R^2 \oplus R^2, d_{X_0})$ of $x^3 + xy^3 - v^2$ with twisted differential

$$d_{X_0}^1 = \begin{pmatrix} v & -x \\ x^2 + y^3 & -v \end{pmatrix}, \quad d_{X_0}^0 = (d_{X_0}^1)^\# , \quad (2.38)$$

one of the seven indecomposable objects of $\text{hmf}^{\text{gr}}(\mathbb{C}[v, x, y], V(\mathbf{E}_7) - v^2)$ listed in [KST]. In step (iii) we again choose a similarity transformation that removes the

term xu^3 in the diagonal entries. The result of step (iv) reads

$$\begin{aligned} d_{X_u}^1 &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{with} \quad a = v - \frac{t^2-10t+19}{2} (su)^9 + (t-2)y(su)^5 + y^2(su), \\ & \quad b = -x + (2t-5)(su)^6 + y(su)^2, \\ & \quad c = x^2 + y^3 + (2t-5)^2 (su)^{12} + (2t-5)x(su)^6 \\ & \quad \quad \quad + 2(2t-5)y(su)^8 + xy(su)^2 + y^2(su)^4, \\ & \quad d = -v - \frac{t^2-10t+19}{2} (su)^9 + (t-2)y(su)^5 + y^2(su) \end{aligned} \quad (2.39)$$

and $d_{X_0}^0 = (d_{X_u}^1)^\#$. This time s and t can be any solution of

$$t^3 - 21t + 37 = 0, \quad s^{18} = 26220t^2 + 67488t - 376912. \quad (2.40)$$

We find that

$$\dim_l(X_u) = -2s, \quad \dim_r(X_u) = (-30 + 5t + 2t^2)s^{-1}, \quad (2.41)$$

and the the possible \mathbb{Q} -gradings on X_u are given by

$$U_{X_u}(\beta) = e^{i\alpha\beta} \cdot \text{diag}(1, e^{-i\beta/3}, 1, e^{-i\beta/3}) \quad \text{with} \quad \alpha \in \mathbb{Q}. \quad (2.42)$$

Lemma 2.11. Let Y be a rank-three \mathbb{Q} -graded matrix factorisation of $x^3 + xy^3 - u^{18} - v^2$. Suppose that (i) Y has non-zero quantum dimensions and (ii) when setting u to zero, Y is equal to X_0 in (2.38). Then Y is isomorphic in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, v, x, y], x^3 + xy^3 - u^{18} - v^2)$ to X_u in (2.39) for some solution s, t of (2.40).

As before we compute the endomorphisms of X_u in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, v, x, y], x^3 + xy^3 - u^{18} - v^2)$ to be the 27-dimensional space

$$\text{End}(X_u) \cong \left(\bigoplus_{i=0}^{16} \mathbb{C}[i \cdot \frac{2}{18}] \right) \oplus \left(\bigoplus_{i=4}^{12} \mathbb{C}[i \cdot \frac{2}{18}] \right) \oplus \mathbb{C}[\frac{8}{18}], \quad (2.43)$$

and using the code of [CM1] together with row and column manipulations as in the previous example, for the solution

$$t_{\text{cft}} = 3(\zeta_{18} + \zeta_{18}^{-1}) - 2(\zeta_{18}^2 + \zeta_{18}^{-2}) \quad \text{where} \quad \zeta_d = e^{2\pi i/d}, \quad (2.44)$$

to (2.40) we find that $X_{u'}^\dagger \otimes X_u$ is isomorphic, up to the factor I_{v^2} , to the rank-three graded matrix factorisation

$$P_{\{0\}} \oplus P_{\{-4, -3, \dots, 4\}} \oplus P_{\{-8, -7, \dots, 8\}}, \quad (2.45)$$

proving the second third of Corollary 1.7.

Remark 2.12. As in Remark 2.10 one can check that the other solutions t to (2.40) form a single orbit under the Galois group $\text{Gal}(\mathbb{Q}(\zeta_{18})/\mathbb{Q})$: the other two solutions are

$$\begin{aligned}\sigma_5(t_{\text{cft}}) &= 3(\zeta_{18}^5 + \zeta_{18}^{-5}) - 2(\zeta_{18}^{10} + \zeta_{18}^{-10}), \\ \sigma_7(t_{\text{cft}}) &= 3(\zeta_{18}^7 + \zeta_{18}^{-7}) - 2(\zeta_{18}^{14} + \zeta_{18}^{-14}).\end{aligned}\tag{2.46}$$

Computing the actions of σ_{5*} and σ_{7*} on $\{-4, -3, \dots, 4\}$ and $\{-8, -7, \dots, 8\}$, e.g. $\sigma_{5*}(\{-4, -3, \dots, 4\}) = \{0, \pm 2, \pm 3, \pm 5, \pm 8\}$, one thus finds that

$$\begin{aligned}X_{u'}^\dagger \otimes X_u &\cong (P_{\{0\}} \oplus P_{\{0, \pm 2, \pm 3, \pm 5, \pm 8\}} \oplus P_{\{-8, -7, \dots, 8\}}) \otimes_k I_{v^2} & \text{for } t = \sigma_5(t_{\text{cft}}), \\ X_{u'}^\dagger \otimes X_u &\cong (P_{\{0\}} \oplus P_{\{0, \pm 3, \pm 4, \pm 7, \pm 8\}} \oplus P_{\{-8, -7, \dots, 8\}}) \otimes_k I_{v^2} & \text{for } t = \sigma_7(t_{\text{cft}}).\end{aligned}\tag{2.47}$$

$$\mathbf{V}(\mathbf{E}_8) \sim \mathbf{V}(\mathbf{A}_{29})$$

The \mathbf{E}_8 -case is considerably more complicated than the cases already treated. This starts already in step (i) as the smallest factorisations of $V^{(\mathbf{E}_8)} - v^2$ are of rank four. Let us choose

$$d_{X_0}^1 = \begin{pmatrix} -v & 0 & x & y \\ 0 & -v & y^4 & -x^2 \\ x^2 & y & -v & 0 \\ y^4 & -x & 0 & -v \end{pmatrix}, \quad d_{X_0}^0 = \begin{pmatrix} v & 0 & x & y \\ 0 & v & y^4 & -x^2 \\ x^2 & y & v & 0 \\ y^4 & -x & 0 & v \end{pmatrix},\tag{2.48}$$

see [KST, Sect. 5]. This is a matrix factorisation of $x^3 + y^5 - v^2$.

In step (ii), the most generic homogeneous deformation of $d_{X_0}^1$ (deforming also the zero entries, of course) has 82 free parameters. Via the similarity transformation in step (iii) one can reduce this to 60 parameters. We refrain from giving this general deformation explicitly.

For step (iv) one has to use a different method than in the other cases, because now $\det(d_{X_0}^1) = (x^3 + y^5 - v^2)^2$ and imposing $\det(d_{X_u}^1) = (x^3 + y^5 - u^{30} - v^2)^2$ turns out to be impractical as it results in too many non-linear conditions. Instead, we make the ansatz $d_{X_u}^0 = (x^3 + y^5 - u^{30} - v^2)^{-1} \cdot (d_{X_u}^1)^\#$ and require that $d_{X_u}^0$ be a matrix with polynomial entries. As before, this leads to a (very long) matrix factorisation of $x^3 + y^5 - u^{30} - v^2$ in terms of two parameters s, t , where t satisfies an eighth order equation and s^{30} is equal to some polynomial in t . The eighth order equation, however, is a product of two fourth order ones, and we select one of these and use it to simplify the matrix factorisation. One is left with the matrix $m := d_{X_u}^1$, where with $\varsigma = su$ the matrix entries m_{ij} are as follows:

$$\begin{aligned}m_{11} &= -v - \frac{(1+t)(3+t)(5+7t)}{64} \varsigma^{15} - \frac{1+t}{4} \varsigma^5 x - \frac{19+47t+25t^2+5t^3}{192} \varsigma^9 y - \frac{1}{2} \varsigma^3 y^2, \\ m_{12} &= \varsigma,\end{aligned}$$

$$\begin{aligned}
m_{13} &= x + \frac{(-1+t)(23+36t+5t^2)}{96} \zeta^{10}, \\
m_{14} &= y, \\
m_{21} &= \frac{-138089-562209t-600371t^2-116355t^3}{11520} \zeta^{29} + \frac{-73-280t-285t^2-50t^3}{160} \zeta^{19} x \\
&\quad + \frac{-29-25t+25t^2+5t^3}{96} \zeta^9 x^2 + \frac{-2107-8545t-9085t^2-1735t^3}{960} \zeta^{23} y \\
&\quad + \frac{-33-57t-11t^2+5t^3}{64} \zeta^{13} xy + \frac{(5+7t)(13+36t+7t^2)}{384} \zeta^{17} y^2 - \frac{3+4t}{4} \zeta^7 xy^2 \\
&\quad + \frac{-35-49t+7t^2+5t^3}{96} \zeta^{11} y^3 - \zeta xy^3 - \frac{1}{2}(1+t)\zeta^5 y^4, \\
m_{22} &= -v + \frac{(1+t)(3+t)(5+7t)}{64} \zeta^{15} + \frac{1+t}{4} \zeta^5 x + \frac{19+47t+25t^2+5t^3}{192} \zeta^9 y + \frac{1}{2} \zeta^3 y^2, \\
m_{23} &= y^4 + \frac{3587+14687t+15785t^2+3125t^3}{1920} \zeta^{24} + \frac{(1-t)(23+36t+5t^2)}{96} \zeta^9 v \\
&\quad + \frac{43+102t+67t^2+12t^3}{96} \zeta^{14} x - \frac{(1+t)(81+126t+17t^2)}{384} \zeta^{18} y + \frac{2+3t}{4} \zeta^8 xy \\
&\quad + \frac{(2+t)(7+6t-5t^2)}{96} \zeta^{12} y^2 + \zeta^2 xy^2 + \frac{1+2t}{4} \zeta^6 y^3, \\
m_{24} &= -x^2 + \frac{(-1+t)(23+36t+5t^2)}{96} \zeta^{10} x + \frac{2+21t+32t^2+9t^3}{48} \zeta^{14} y, \\
m_{31} &= x^2 + \frac{(1-t)(23+36t+5t^2)}{96} \zeta^{10} x - \frac{2+21t+32t^2+9t^3}{48} \zeta^{14} y, \\
m_{32} &= y, \\
m_{33} &= -v + \frac{-37-39t+29t^2+15t^3}{192} \zeta^{15} + \frac{1+t}{4} \zeta^5 x + \frac{-65-73t+37t^2+5t^3}{192} \zeta^9 y - \frac{1}{2} \zeta^3 y^2, \\
m_{34} &= \frac{(1-t)(23+36t+5t^2)}{96} \zeta^{11} + \zeta x + \frac{1+t}{2} \zeta^5 y, \\
m_{41} &= y^4 + \frac{3587+14687t+15785t^2+3125t^3}{1920} \zeta^{24} + \frac{(-1+t)(23+36t+5t^2)}{96} \zeta^9 v \\
&\quad + \frac{43+102t+67t^2+12t^3}{96} \zeta^{14} x - \frac{(1+t)(81+126t+17t^2)}{384} \zeta^{18} y + \frac{2+3t}{4} \zeta^8 xy \\
&\quad + \frac{(2+t)(7+6t-5t^2)}{96} \zeta^{12} y^2 + \zeta^2 xy^2 + \frac{1+2t}{4} \zeta^6 y^3, \\
m_{42} &= -x + \frac{(1-t)(23+36t+5t^2)}{96} \zeta^{10}, \\
m_{43} &= -\frac{569+2615t+2855t^2+425t^3}{1920} \zeta^{19} + \frac{17+t-37t^2-5t^3}{96} \zeta^9 x + \frac{-17-17t+13t^2+5t^3}{64} \zeta^{13} y \\
&\quad - \frac{1+2t}{4} \zeta^7 y^2 - \zeta y^3, \\
m_{44} &= -v + \frac{37+39t-29t^2-15t^3}{192} \zeta^{15} - \frac{1+t}{4} \zeta^5 x + \frac{65+73t-37t^2-5t^3}{192} \zeta^9 y + \frac{1}{2} \zeta^3 y^2.
\end{aligned}$$

The parameters s, t can be any solution of the equations

$$\begin{aligned}
s^{30} &= \frac{1}{4}(45308593275 t^3 - 32199587625 t^2 - 973905678975 t - 395277903075), \\
5 t^4 - 110 t^2 - 120 t - 31 &= 0.
\end{aligned} \tag{2.49}$$

As already noted above, the matrix $d_{X_u}^0$ is given by $(x^3+y^5-u^30-v^2)^{-1} \cdot (d_{X_u}^1)^\#$, which has polynomial entries provided s, t solve (2.49). It is now straightforward to determine the \mathbb{Q} -gradings on X_u to be given by

$$U_{X_u}(\beta) = e^{i\alpha\beta} \cdot \text{diag} \left(1, e^{-14i\beta/15}, e^{-i\beta/3}, e^{-3i\beta/5}, 1, e^{-14i\beta/15}, e^{-i\beta/3}, e^{-3i\beta/5} \right) \tag{2.50}$$

with $\alpha \in \mathbb{Q}$, and compute the quantum dimensions to be

$$\dim_1(X_u) = 2s, \quad \dim_r(X_u) = \frac{5}{16}(-27 - 86t - 3t^2 + 4t^3) s^{-1}, \tag{2.51}$$

which are indeed non-zero for all choices of s, t satisfying (2.49). This concludes the proof of Theorem 1.5.

Because we have made a number of restricting assumptions before arriving at (X, d_{X_u}) , we cannot claim a statement analogous to Lemmas 2.7, 2.9 and 2.11.

Finally we compute the endomorphisms of X_u in $\text{hmf}^{\text{gr}}(\mathbb{C}[u, v, x, y], x^5 + y^3 - u^{30} - v^2)$ to be the 60-dimensional space

$$\text{End}(X_u) \cong \left(\bigoplus_{i=0}^{28} \mathbb{C}[i \cdot \frac{2}{30}] \right) \oplus \left(\bigoplus_{i=5}^{23} \mathbb{C}[i \cdot \frac{2}{30}] \right) \oplus \left(\bigoplus_{i=9}^{19} \mathbb{C}[i \cdot \frac{2}{30}] \right) \oplus \mathbb{C}[\frac{28}{30}]. \quad (2.52)$$

For the solution

$$t_{\text{cft}} = -\frac{1}{5} \left(7 + 4(\zeta_{30} + \zeta_{30}^{-1}) + 8(\zeta_{30}^2 + \zeta_{30}^{-2}) - 16(\zeta_{30}^3 + \zeta_{30}^{-3}) \right) \quad (2.53)$$

of (2.49), $X_{u'}^\dagger \otimes X_u$ is isomorphic, up to the factor I_{v^2} , to the rank-four graded matrix factorisation

$$P_{\{0\}} \oplus P_{\{-5, -4, \dots, 5\}} \oplus P_{\{-9, -8, \dots, 9\}} \oplus P_{\{-14, -13, \dots, 14\}}. \quad (2.54)$$

This concludes the proof of Corollary 1.7.

Remark 2.13. The other solutions of (2.49) are found via the Galois group $\text{Gal}(\mathbb{Q}(\zeta_{30})/\mathbb{Q})$ to be

$$\begin{aligned} \sigma_7(t_{\text{cft}}) &= -\frac{1}{5} \left(7 + 4(\zeta_{30}^7 + \zeta_{30}^{-7}) + 8(\zeta_{30}^{14} + \zeta_{30}^{-14}) - 16(\zeta_{30}^{21} + \zeta_{30}^{-21}) \right), \\ \sigma_{11}(t_{\text{cft}}) &= -\frac{1}{5} \left(7 + 4(\zeta_{30}^{11} + \zeta_{30}^{-11}) + 8(\zeta_{30}^{22} + \zeta_{30}^{-22}) - 16(\zeta_{30}^3 + \zeta_{30}^{-3}) \right), \\ \sigma_{13}(t_{\text{cft}}) &= -\frac{1}{5} \left(7 + 4(\zeta_{30}^{13} + \zeta_{30}^{-13}) + 8(\zeta_{30}^{26} + \zeta_{30}^{-26}) - 16(\zeta_{30}^9 + \zeta_{30}^{-9}) \right). \end{aligned} \quad (2.55)$$

The corresponding decompositions of $X_{u'}^\dagger \otimes X_u$ are, with $Z = P_{\{0\}} \oplus P_{\{-14, -13, \dots, 14\}}$ and up to the factor I_{v^2} ,

$$\begin{aligned} Z \oplus P_{\{0, \pm 2, \pm 5, \pm 7, \pm 9, \pm 14\}} \oplus P_{\{0, \pm 2, \pm 3, \pm 4, \pm 5, \pm 7, \pm 9, \pm 11, \pm 12, \pm 14\}} & \quad \text{for } t = \sigma_7(t_{\text{cft}}), \\ Z \oplus P_{\{0, \pm 3, \pm 5, \pm 8, \pm 11, \pm 14\}} \oplus P_{\{0, \pm 2, \pm 3, \pm 5, \pm 6, \pm 8, \pm 9, \pm 11, \pm 13, \pm 14\}} & \quad \text{for } t = \sigma_{11}(t_{\text{cft}}), \\ Z \oplus P_{\{0, \pm 3, \pm 4, \pm 5, \pm 8, \pm 9, \pm 13\}} \oplus P_{\{0, \pm 1, \pm 4, \pm 5, \pm 8, \pm 9, \pm 12, \pm 13, \pm 14\}} & \quad \text{for } t = \sigma_{13}(t_{\text{cft}}). \end{aligned} \quad (2.56)$$

Another method to construct orbifold equivalences

There is another way, slightly different from the one described at the beginning of Section 2.3, of obtaining the orbifold equivalences $V^{(\text{D}_{d/2+1})} \sim V^{(\text{A}_{d-1})}$, $V^{(\text{E}_6)} \sim V^{(\text{A}_{11})}$, and $V^{(\text{E}_7)} \sim V^{(\text{A}_{17})}$. Roughly, this method starts with a matrix factorisation M of a potential W and computes the deformations [Lau, Sij] M_u

of M . Then instead of setting the deformation parameters to be solutions of the obstruction equations (which would give factorisations of W), one tries to re-interpret some of the parameters u as variables of another potential V while choosing the remaining parameters such that M_u becomes a factorisation of $W - V$.

As this method may prove useful to construct further orbifold equivalences, we illustrate it in more detail in the example of $V^{(E_6)} \sim V^{(A_{11})}$:

- (i) Start with the matrix factorisation (2.23) of $W = V^{(E_6)} - v^2$ and compute its deformations, using e. g. the implementation of [CDR].⁵
- (ii) One obtains a 4-by-4 matrix D with polynomial entries in the variables v, x, y and two deformation parameters u_1, u_2 . By construction D squares to $W \cdot 1 + R$, where the matrix R vanishes if u_1, u_2 satisfy the obstruction equations.
- (iii) Interpret the parameter u_1 as the (rescaled) variable u and choose $u_2 \in \mathbb{C}[u]$ such that D becomes a factorisation of $V^{(E_6)} - u^{12} - v^2$. This turns out to be isomorphic to X_u of (2.26) with $t = -1/\sqrt{3}$.

In the cases $W + v^2 \in \{V^{(D_{d/2+1})}, V^{(E_6)}, V^{(E_7)}\}$ this method is especially straightforward as the matrix D already squares to $(W + f) \cdot 1$, where f is a polynomial with leading term $u_1^{d/2}, u_1^{12}, u_1^{18}$, respectively. On the other hand, for $W = V^{(E_8)} - v^2$ the square of D is a more generic matrix, rendering the problem $D^2 = (W - u^{30}) \cdot 1$ much more computationally involved.

3 Comparison to conformal field theory

In this section we describe how the orbifold equivalences established in Theorem 1.5 compare to – and are in fact predicted by – the correspondence between $\mathcal{N} = 2$ supersymmetric Landau-Ginzburg models (LG models) and $\mathcal{N} = 2$ superconformal field theories (CFTs) in two dimensions. We do not aim for a self-contained discussion, but we provide pointers to the literature where more details can be found.

One can argue that for a given LG model there is an associated CFT, namely the infrared fixed point theory [Mar, VW, HW]. The Virasoro central charge of the infrared CFT is computed from the potential of the LG model via (1.4). The renormalisation group flow also provides a map from certain boundary conditions and defects lines in the LG model to certain conformal boundary conditions and defects lines in the CFT, see e. g. [KL, BHLS, BR1]. In general, charges and correlators of fields in the LG model vary along the flow. However, by $\mathcal{N} = 2$ supersymmetry the charges and correlators in a subsector of the LG model

⁵Of course the indecomposable object (2.23) has no nontrivial deformations, but that does not matter here.

consisting of chiral primary fields are preserved and can be directly compared to their CFT equivalents. Below, we will carry out this comparison for the defects X_u of Section 2.3 and the charge spectrum of the chiral primary fields supported on these defects.

Conformal field theories form a bicategory, where objects are CFTs, 1-morphisms are topological defect lines and 2-morphisms are topological junction fields, see [DKR] for details. This bicategory has adjoints and is pivotal, hence the general concept of orbifold equivalence is applicable. There cannot be a topological defect joining two CFTs of different Virasoro central charge. As a consequence, equality of central charges provides a *necessary* condition for an orbifold equivalence to exist, cf. Proposition 1.3. Contrary to the matrix factorisation framework, for CFTs a useful *sufficient* condition is known for the existence of an orbifold equivalence [FFRS2]:

Let V be a rational vertex operator algebra. Suppose two CFTs have unique bulk vacua and their algebra of bulk fields contains $V \otimes_{\mathbb{C}} \bar{V}$ as a subalgebra. Then these two CFTs are orbifold equivalent.

Here the bar over V indicates that the second factor is embedded in the anti-holomorphic fields.

Those LG models whose potentials define simple singularities are believed to renormalise to CFTs that contain the $\mathcal{N} = 2$ minimal super Virasoro vertex operator algebra with the same central charge. The latter are rational, so by the above criterion all these CFTs of the same central charge are orbifold equivalent.

One may expect topological defects between two infrared CFTs to have analogues already in the corresponding LG model. Furthermore, one may expect that composition of topological defects commutes with renormalisation group flow. If so, the orbifold equivalences of infrared CFTs should exist also for LG models. This is the reason why Theorem 1.5 is expected from the above CFT criterion.

After these qualitative considerations, we now turn to quantitative, more technical comparisons. The correspondence between topological defect lines of LG models of A-type singularities and those of the associated diagonal $\mathcal{N} = 2$ CFTs motivates the following conjecture, whose ingredients we explain directly after stating it.

Conjecture 3.1. For any integer $d \geq 3$, consider the monoidal subcategory of $\text{hmf}^{\text{gr}}(\mathbb{C}[x, x'], x^d - x'^d)$ whose morphisms only have \mathbb{Q} -degree zero, and which is generated by $\{P_{\{a, a+1, \dots, a+b\}} \mid a, b \in \mathbb{Z}_d\}$ with respect to tensor products and direct sums. This subcategory is monoidally equivalent to $(\mathcal{C}_{d-2}^{\text{su}(2)} \boxtimes \mathcal{C}_{2d}^{\text{u}(1)})_{\text{NS}}$.

The matrix factorisations P_S are those introduced in (1.8). The relevant \mathbb{Q} -grading on P_S follows from the R-charge discussion in [CR1, Sect. 3.2]. By $\mathcal{C}_k^{\text{su}(2)}$ we denote the modular tensor category of integrable highest weight representations of the affine Lie algebra $\widehat{\mathfrak{su}}(2)$ at level k . It has simple objects labelled

by $\{0, 1, \dots, k\}$. $\mathcal{C}_{2d}^{u(1)}$ is the modular tensor category of representations of the rational lattice vertex operator algebra built out of single free boson which has simple objects labelled by $\{0, 1, \dots, 2d - 1\}$, and \boxtimes denotes the Deligne product, see e. g. [BK]. Details and references to the original literature can be found in [CR1, App. A.2].

We set $\mathcal{D} = \mathcal{C}_{d-2}^{su(2)} \boxtimes \mathcal{C}_{2d}^{u(1)}$ and denote the simple objects of \mathcal{D} by $U_{l,m}$, where $l \in \{0, 1, \dots, d-2\}$ and $m \in \{0, 1, \dots, 2d-1\}$. The notation $(-)_\text{NS}$ refers to passing to the full subcategory generated by the $U_{l,m}$ with $l+m$ even. Via the coset construction, this describes the NS sector of the representations of the $\mathcal{N} = 2$ minimal super Virasoro vertex operator algebra at central charge $c = 3(1 - 2/d)$, see again [CR1, App. A.2] for details and references.⁶

The status of Conjecture 3.1 is currently as follows.

- The functor which conjecturally provides the tensor equivalence acts on simple objects as $P_{\{a, a+1, \dots, a+b\}} \mapsto U_{b, b+2a}$. It is verified in [BR1] that this is compatible with the tensor product on the level of isomorphism classes.
- Some, but far from all, associativity isomorphisms were proved to be compatible in [CR1].
- For odd d , the conjecture is proved in [DRCR].

Comparison of algebra objects

According to [FRS, FFRS], the full conformal field theories (with unique bulk vacuum) that can be constructed starting from a rational vertex operator algebra V are parametrised by Morita classes of Δ -separable symmetric Frobenius algebras in the modular tensor category of representations of V . The algebras relevant for the CFT describing the infrared fixed point of an LG model with ADE-type potential are predicted from [Gan, Gra] to be non-trivial only in the $\mathfrak{su}(2)$ factor of \mathcal{D} . More specifically, they are representatives of the ADE classification of Morita classes of such algebras in $\mathcal{C}_{d-2}^{su(2)}$ given in [Ost]. As objects in \mathcal{D} these algebras are

$$\begin{aligned}
F^{(A_{d-1})} &= U_{0,0} && \text{for } d \geq 2, \\
F^{(D_{d/2+1})} &= U_{0,0} \oplus U_{d-2,0} && \text{for } d \in 2\mathbb{Z}_+, \\
F^{(E_6)} &= U_{0,0} \oplus U_{6,0} && \text{for } d = 12, \\
F^{(E_7)} &= U_{0,0} \oplus U_{8,0} \oplus U_{16,0} && \text{for } d = 18, \\
F^{(E_8)} &= U_{0,0} \oplus U_{10,0} \oplus U_{18,0} \oplus U_{28,0} && \text{for } d = 30. \tag{3.1}
\end{aligned}$$

⁶In [CR1] the $(-)_\text{NS}$ is missing in (A.38) and (A.45). This is a typo or an error, depending on one's disposition towards the authors.

Topological defects between a diagonal, i. e. A-type, CFT and another CFT from the above list with the same value of d given by some algebra F are described by F -modules in \mathcal{D} [FFRS1]. This is consistent with the point of view of orbifold equivalences. There, the algebra describing the theory on one side of a topological defect X in terms of the other is $X^\dagger \otimes X$, see [CR3] for details. In the present setting one has $X = {}_F F$, $X^\dagger = F_F$, and $F_F \otimes_F {}_F F \cong F$ as algebras.

On the matrix factorisation side, the objects underlying the algebras describing the D- and E-type singularities as orbifolds of A-type singularities are those in (2.21), (2.35), (2.45) and (2.54) above. Under the tensor equivalence of Conjecture 3.1, they are indeed mapped to the corresponding objects in the list (3.1). (It would of course be enough to land in the same Morita class, but for our choices of matrix factorisations we get the actual representatives chosen in (3.1).)

Comparison of defect spectra

Given the above observations on the objects underlying the algebras establishing the orbifold equivalences, it is natural to expect that the matrix factorisations of the potential differences described in Section 2.3 get mapped to the topological defects described by the modules ${}_F F$ for F the corresponding algebra in (3.1). This expectation can be tested by comparing the spectra of chiral primaries.

The chiral primaries in \mathcal{D} are the ground states in the representations labelled $U_{l,l}$ with $l \in \{0, 1, \dots, d-2\}$; their charge is l/d . The space of chiral primaries of holomorphic and antiholomorphic labels l and m , respectively, on the defect ${}_F F$ is isomorphic to the vector space

$$\mathrm{Hom}_F(F \otimes U_{l,l} \otimes U_{m,-m}, F) \cong \mathrm{Hom}(U_{l,l} \otimes U_{m,-m}, F), \quad (3.2)$$

where ‘ Hom_F ’ comprises only F -module maps in \mathcal{D} , and ‘ Hom ’ all maps in \mathcal{D} . We refer to [FFRS1] for an explanation of this formula. The total charge (as seen on the LG side) of these chiral primaries is $(l+m)/d$.

Let us consider the example $F = F^{(E_6)}$ in some detail. The first thing to note is that $\mathrm{Hom}(U_{l,l} \otimes U_{m,-m}, U_{n,0})$ has dimension zero unless $l = m$. (Actually the fusion rules in $\mathcal{C}_{2d}^{u(1)}$ tell us that $l - m \cong 0 \pmod{2d}$, but for the given range on l and m this just amounts to $l = m$.) For $l = m$, the space is one-dimensional if n is even and $n/2 \leq l \leq d - 2 - n/2$, as follows from the fusion rules of $\mathcal{C}_{d-2}^{su(2)}$. From the summand $U_{0,0}$ in $F^{(E_6)}$ we therefore get one state at each charge $l/6$, $l \in \{0, 1, \dots, 10\}$, and from the summand $U_{6,0}$ another state of charge $l/6$ for each $l \in \{3, 4, \dots, 7\}$. Hence the total dimension of the space of chiral primaries is 16, and we have perfect agreement with (2.31).

It is straightforward to carry out the analogous computations for $F = F^{(D_{d/2+1})}$, $F = F^{(E_7)}$ and $F = F^{(E_8)}$, again consistent with the charges and multiplicities of the matrix factorisation results listed in (2.21), (2.43) and (2.52), respectively.

References

- [BBR] C. Bachas, I. Brunner, and D. Roggenkamp, *A worldsheet extension of $O(d, d|\mathbb{Z})$* , *JHEP* **1210** (2012), 1210, [[arXiv:1205.4647](#)].
- [BCP] I. Brunner, N. Carqueville, and D. Plencner, *A quick guide to defect orbifolds*, *Proceedings of Symposia in Pure Mathematics* **88** (2014), 231–241, [[arXiv:1310.0062](#)].
- [BFK] M. Ballard, D. Favero and L. Katzarkov, *A category of kernels for equivariant factorizations and its implications for Hodge theory*, [[arXiv:1105.3177](#)].
- [BHLS] I. Brunner, M. Herbst, W. Lerche, and B. Scheuner, *Landau-Ginzburg Realization of Open String TFT*, *JHEP* **0611** (2003), 043, [[hep-th/0305133](#)].
- [BK] B. Bakalov and A. A. Kirillov Jr., *Lectures on Tensor Categories and Modular Functors*, American Mathematical Society, Providence, 2001.
- [Bor] F. Borceux, *Handbook of categorical algebra 1*, volume 50 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, 1994.
- [BR1] I. Brunner and D. Roggenkamp, *B-type defects in Landau-Ginzburg models*, *JHEP* **0708** (2007), 093, [[arXiv:0707.0922](#)].
- [BR2] I. Brunner and D. Roggenkamp, *Defects and Bulk Perturbations of Boundary Landau-Ginzburg Orbifolds*, *JHEP* **0804** (2008), 001, [[arXiv:0712.0188](#)].
- [Buc] R.-O. Buchweitz, *Maximal Cohen-Macaulay Modules and Tate-Cohomology over Gorenstein Rings*, (1986), [University of Toronto Research Repository](#).
- [Car] N. Carqueville, *Equivariant completion*, *Oberwolfach Report No. 44/2013*, also available at [nils.carqueville.net/OWR-Beq.pdf](#).
- [CDR] N. Carqueville, L. Dowdy, and A. Recknagel, *Algorithmic deformation of matrix factorisations*, *JHEP* **04** (2012), 014, [[arXiv:1112.3352](#)].
- [CM1] N. Carqueville and D. Murfet, *Computing Khovanov-Rozansky homology and defect fusion*, *Algebraic & Geometric Topology* **14** (2014), 489–537, [[arXiv:1108.1081](#)].
- [CM2] N. Carqueville and D. Murfet, *Adjunctions and defects in Landau-Ginzburg models*, [[arXiv:1208.1481](#)].

- [CM3] N. Carqueville and D. Murfet, *A toolkit for defect computations in Landau-Ginzburg models*, [[arXiv:1303.1389](#)].
- [CR1] N. Carqueville and I. Runkel, *On the monoidal structure of matrix bi-factorisations*, *J. Phys. A: Math. Theor.* **43** (2010), 275401, [[arXiv:0909.4381](#)].
- [CR2] N. Carqueville and I. Runkel, *Rigidity and defect actions in Landau-Ginzburg models*, *Comm. Math. Phys.* **310** (2012), 135–179, [[arXiv:1006.5609](#)].
- [CR3] N. Carqueville and I. Runkel, *Orbifold completion of defect bicategories*, to appear in *Quantum Topology*, [[arXiv:1210.6363](#)].
- [CV] C. Cecotti and C. Vafa, *On Classification of $N = 2$ Supersymmetric Theories*, *Comm. Math. Phys.* **158** (1993), 569–644, [[hep-th/9211097](#)].
- [DKR] A. Davydov, L. Kong, and I. Runkel, *Field theories with defects and the centre functor*, *Mathematical Foundations of Quantum Field Theory and Perturbative String Theory*, Proceedings of Symposia in Pure Mathematics, AMS, 2011, [[arXiv:1107.0495](#)].
- [DM] T. Dyckerhoff and D. Murfet, *Pushing forward matrix factorisations*, *Duke Math. J.* **162**, Number 7 (2013), 1249–1311, [[arXiv:1102.2957](#)].
- [DRCR] A. Davydov, A. Ros Camacho, and I. Runkel, *$N = 2$ minimal conformal field theories and matrix bifactorisations of x^d* , [[arXiv:1409.2144](#)].
- [FFRS] J. Fjelstad, J. Fuchs, I. Runkel, and C. Schweigert, *Uniqueness of open/closed rational CFT with given algebra of open states*, *Adv. Theor. Math. Phys.* **12** (2008), 1283–1375, [[hep-th/0612306](#)].
- [FFRS1] J. Fröhlich, J. Fuchs, I. Runkel and C. Schweigert, *Duality and defects in rational conformal field theory*, *Nucl. Phys. B* **763** (2007), 354–430, [[hep-th/0607247](#)].
- [FFRS2] J. Fröhlich, J. Fuchs, I. Runkel and C. Schweigert, *Defect lines, dualities, and generalised orbifolds*, *Proceedings of the XVI International Congress on Mathematical Physics*, Prague, August 3–8, 2009, [[arXiv:0909.5013](#)].
- [FGRS] J. Fuchs, M. R. Gaberdiel, I. Runkel and C. Schweigert, *Topological defects for the free boson CFT*, *J. Phys. A* **40** (2007), 11403, [[arXiv:0705.3129](#)].

- [FRS] J. Fuchs, I. Runkel, and C. Schweigert, *TFT construction of RCFT correlators. I: Partition functions*, Nucl. Phys. B **646** (2002), 353–497, [[hep-th/0204148](#)].
- [Gan] T. Gannon, *$U(1)^m$ modular invariants, $N = 2$ minimal models, and the quantum Hall effect*, Nucl. Phys. B **491** (1997), 659–688, [[hep-th/9608063](#)].
- [GM] S. I. Gelfand and Y. I. Manin, *Methods of Homological Algebra*, Springer Monographs in Mathematics, Springer, 2010.
- [Gra] O. Gray, *On the complete classification of the unitary $N = 2$ minimal superconformal field theories*, Comm. Math. Phys. **312** (2012), 611–654, [[arXiv:0812.1318](#)].
- [HW] P. Howe and P. West, *$N = 2$ superconformal models, Landau-Ginzburg hamiltonians and the ϵ expansion*, Phys. Lett. B **223** (1989), 377–385; *Chiral Correlators In Landau-Ginzburg Theories And $N = 2$ Superconformal Models*, Phys. Lett. B **227** (1989), 397–405; *Fixed points in multifield Landau-Ginzburg models*, Phys. Lett. B **244** (1990), 270–274.
- [KL] A. Kapustin and Y. Li, *D-branes in Landau-Ginzburg Models and Algebraic Geometry*, JHEP **0312** (2003), 005, [[hep-th/0210296](#)].
- [KST] H. Kajiura, K. Saito, and A. Takahashi, *Matrix Factorizations and Representations of Quivers II: type ADE case*, Adv. Math. **211** (2007), 327–362, [[math.AG/0511155](#)].
- [Lau] O. A. Laudal, *Matric Massey products and formal moduli I*, Lecture Notes in Mathematics **1183** (1983), 218–240.
- [Mar] E. J. Martinec, *Algebraic Geometry and Effective Lagrangians*, Phys. Lett. B **217** (1989), 431–437.
- [Nee] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- [Orl] D. Orlov, *Derived Categories of Coherent Sheaves and Triangulated Categories of Singularities*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., **270**, 503–531, Birkhäuser Boston, Inc., Boston, MA, 2009, [[math.AG/0503632](#)].
- [Ost] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups **8** (2003), 177–206, [[math.QA/0111139](#)].
- [RR] I. Reiten and C. Riedtmann, *Skew group algebras in the representation theory of Artin algebras*, J. Algebra **92** (1985), 224–282.

- [Sch] G. Schaumann, *Duals in tricategories and in the tricategory of bimodule categories*, Ph.D. thesis, available at <https://opus4.kobv.de/opus4-fau/frontdoor/index/index/docId/3732>.
- [Siq] A. Siqueland, *The method of computing formal moduli*, *J. of Algebra* **241** (2001), 291–327.
- [VW] C. Vafa and N. Warner, *Catastrophes and the classification of conformal theories*, *Phys. Lett. B* **218** (1989), 51–58.
- [Yos] Y. Yoshino, *Maximal Cohen-Macaulay Modules Over Cohen-Macaulay Rings*, London Mathematical Society Lecture Note Series **146**, Cambridge University Press, 1990.