

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/133532/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Arbyn, Marc, Bruni, Laia, Kelly, Daniel, Basu, Partha, Poljak, Mario, Gultekin, Murat, Bergeron, Christine, Ritchie, David and Weiderpass, Elisabete 2020. Tackling cervical cancer in Europe amidst the COVID-19 pandemic. *Lancet Public Health* 5 (8), E425. 10.1016/S2468-2667(20)30122-5

Publishers page: <https://www.thelancet.com/journals/lanpub/article/...>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Tackling cervical cancer in Europe amidst the COVID-19 pandemic

According to estimates for 2018,¹ approximately 33 000 cases of cervical cancer occurred and 15 000 people died from the disease in Europe (see map in appendix). Human papillomavirus (HPV) vaccine coverage is relatively low in countries with the highest incidence and screening performance is heterogeneous among European countries. Cytological screening followed by treatment of screen-detected cervical lesions has resulted in substantial decreases in the burden of cervical cancer in western and northern Europe; but in eastern Europe, cervical cancer incidence and mortality remain comparatively high.²

Today, new powerful tools are available for primary and secondary prevention of cervical cancer, among which prophylactic HPV vaccines, and screening using validated HPV tests for women—including some tests that can be applied on self-collected samples, a strategy that might be used to reach underscreened populations.³

In May, 2018, the WHO Director General launched a call to eliminate cervical cancer by vaccinating at least 90% of girls by age 15 years, by offering screening at least twice in a lifetime to 70% or more of the target age populations and treat more than 90% of women with screen-detected lesions.

Recently, a large group of experts from diverse professional societies and cancer organisations supporting WHO's call to eliminate cervical cancer, proposed a series of concerted actions to implement organised integrated HPV vaccination and HPV-based screening, and requested European health authorities to endorse the principles of the WHO elimination initiative, mobilise resources to update evidence-based guidelines, and translate them into quantified and monitored preventive activities.⁴

However, in the first half of 2020, due to the dramatic COVID-19 pandemic, cervical cancer prevention activities have been disrupted in many European countries. We are concerned and urge the public health community to maintain sufficient resources to sustain HPV vaccination and cancer screening in the future.

Importantly, the COVID-19 pandemic might also generate opportunities for more efficient prevention, by promoting more cost-effective, evidence-based protocols, by focusing on women who are at high-risk, extending HPV testing on self-collected samples, and discouraging inefficient policies, such as screening with two tests. We welcome the unprecedented collaborations between the cancer and infectious disease communities, who have been working jointly to tackle the spread of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) by applying the experience of HPV test evaluation to protocols for comparing and validating SARS-CoV-2 assays and by bringing modellers together in the COVID-19 and Cancer Taskforce Global Modelling Consortium.

MA was supported by Directorate-General for Research and Innovation of the European Commission (Regional Invasive Species and Climate Change Network, grant number 847845); Belgian Foundation Against Cancer (IHUVAC project) and from the VALCOR project. LB's research unit has received unrestricted research grants from Merck, Sharpe & Dohme and GlaxoSmithKline. MG received travel support and honoraria from Merck, Sharpe & Dohme to be a speaker. DK, PB, MP, CB, DR, and EW declare no competing interests. Where authors are identified as personnel of the International Agency for Research on Cancer or WHO, the authors alone are responsible for the views expressed in this Article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer or WHO.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

**Marc Arbyn, Laia Bruni, Daniel Kelly, Partha Basu, Mario Poljak, Murat Gultekin, Christine Bergeron, David Ritchie, Elisabete Weiderpass*
marc.arbyn@sciensano.be

Unit of Cancer Epidemiology, Belgian Cancer Centre, Sciensano, Brussels B1050, Belgium (MA);

Consortium for Biomedical Research in Cancer Epidemiology Research Program, Catalan Institute of Oncology - IDIBELL, Barcelona, Spain (LB); Royal College of Nursing, School of Healthcare Sciences, University of Cardiff, Cardiff, UK (DK); International Agency for Research on Cancer, Lyon, France (PB, EW); Institute of Microbiology and Immunology, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia (MP); Hacettepe University Faculty of Medicine, Department of Obstetrics and Gynecology, Division of Gynaecological Oncology, Ankara, Turkey (MG); Laboratoire Cerba, Cergy Pontoise France, Paris, France (CB); and Association of European Cancer Leagues, Brussels, Belgium (DR)

- 1 Arbyn M, Weiderpass E, Bruni L, et al. Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *Lancet Glob Health* 2020; **8**: e191–203.
- 2 Arbyn M, Antoine J, Mägi M, et al. Trends in cervical cancer incidence and mortality in the Baltic countries, Bulgaria and Romania. *Int J Cancer* 2011; **128**: 1899–907.
- 3 Arbyn M, Smith SB, Temin S, et al. Detecting cervical precancer and reaching underscreened women by using HPV testing on self samples: updated meta-analyses. *BMJ* 2018; **363**: k4823.
- 4 Kyrgiou, M, Arbyn M, Bergeron C. et al. Cervical screening: ESGO-EFC position paper of the European Society of Gynaecologic Oncology (ESGO) and the European Federation of Colposcopy (EFC). *Br J Cancer* 2020; published online June 8. <https://doi.org/10.1038/s41416-020-0920-9>.
- 5 Gultekin M, Ramirez PT, Broutet N, Hutubessy R. World Health Organization call for action to eliminate cervical cancer globally. *Int J Gynecol Cancer* 2020; **30**: 426–27.



Lancet Public Health 2020

Published Online
 July 13, 2020
[https://doi.org/10.1016/2468-2667\(20\)30122-5](https://doi.org/10.1016/2468-2667(20)30122-5)

See Online for appendix

For WHO DG call see <https://www.who.int/dg/speeches/detail/cervical-cancer-an-ncd-we-can-overcome>.