

Effects of a functional COMT polymorphism on brain anatomy and cognitive function in adults with velo-cardio-facial syndrome

T. van Amelsvoort^{1*}, J. Zinkstok¹, M. Figuee¹, E. Daly², R. Morris², M. J. Owen³, K. C. Murphy²,
L. De Haan¹, D. H. Linszen¹, B. Glaser³ and D. G. M. Murphy²

¹ Department of Psychiatry, Academic Medical Centre, Amsterdam, The Netherlands

² Department of Psychological Medicine, Institute of Psychiatry, King's College London, London, UK

³ Department of Psychological Medicine, Cardiff University, Cardiff, UK

Background. Velo-cardio-facial syndrome (VCFS) is associated with deletions at chromosome 22q11, abnormalities in brain anatomy and function, and schizophrenia-like psychosis. Thus it is assumed that one or more genes within the deleted region are crucial to brain development. However, relatively little is known about how genetic variation at 22q11 affects brain structure and function. One gene on 22q11 is catechol-O-methyltransferase (COMT): an enzyme that degrades dopamine and contains a functional polymorphism (*Val*¹⁵⁸*Met*) affecting enzyme activity. Here, we investigated the effect of COMT *Val*¹⁵⁸*Met* polymorphism on brain anatomy and cognition in adults with VCFS.

Method. The COMT *Val*¹⁵⁸*Met* polymorphism was genotyped for 26 adults with VCFS on whom DNA was available. We explored its effects on regional brain volumes using hand tracing approaches; on regional grey- and white-matter density using computerized voxel-based analyses; and measures of attention, IQ, memory, executive and visuospatial function using a comprehensive neuropsychological test battery.

Results. After corrections for multiple comparisons *Val*-hemizygous subjects, compared with *Met*-hemizygotes, had a significantly larger volume of frontal lobes. Also, *Val*-hemizygotes had significantly increased grey matter density in cerebellum, brainstem, and parahippocampal gyrus, and decreased white matter density in the cerebellum. No significant effects of COMT genotype on neurocognitive performance were found.

Conclusions. COMT genotype effects on brain anatomy in VCFS are not limited to frontal regions but also involve other structures previously implicated in VCFS. This suggests variation in COMT activity is implicated in brain development in VCFS.

Received 19 October 2006; Revised 21 March 2007; Accepted 30 March 2007; First published online 10 May 2007

Key words: Cognition, COMT, MRI, VCFS, 22q11.

Introduction

Velo-cardio-facial syndrome (VCFS) is a genetic disorder associated with microdeletions at chromosome 22q11 (Shprintzen *et al.* 1978), typical facial morphology, cardiovascular anomalies, velopharyngeal insufficiencies, mild to borderline learning disabilities, cognitive deficits and a high prevalence of psychiatric disorders including ADHD (Papolos *et al.* 1996) and schizophrenia-like psychosis (Murphy *et al.* 1999). This high prevalence of psychiatric disorders is likely to be caused by haploinsufficiency of one or more genes on 22q11.

One of the candidate genes located at 22q11 is the gene for catechol-O-methyltransferase (COMT), an enzyme that degrades dopamine. Particularly in the prefrontal cortex (PFC) degradation of dopamine is largely dependent on activity of COMT, whereas in other brain regions degradation of dopamine is mainly by monoamino-oxidase (MAO) and the dopamine transporter (DAT) (Chen *et al.* 2004). A common mutation in the COMT gene, leading to an amino acid substitution [Valine (*Val*) to Methionine (*Met*)] results in decreased enzymatic activity with the *Met*/*Met* variant of COMT showing 40% less enzymatic activity than the *Val*/*Val* variant (Lachman *et al.* 1996; Chen *et al.* 2004).

The COMT *Val*¹⁵⁸*Met* polymorphism has been reported to: (1) affect midbrain tyrosine hydroxylase levels and dopamine synthesis (Akil *et al.* 2003);

* Address for correspondence: T. van Amelsvoort, M.D.,
Department of Psychiatry, Academic Medical Centre, Tafelbergweg
25, 1105 BC, Amsterdam, The Netherlands.
(Email: t.a.vanamelsvoort@amc.uva.nl)

(2) modulate dopaminergic interactions between the PFC and the midbrain (Meyer-Lindenberg *et al.* 2005; Smolka *et al.* 2005); and (3) affect cognitive functions dependent on PFC, including working memory and attention (Egan *et al.* 2001; Malhotra *et al.* 2002; Meyer-Lindenberg *et al.* 2005). People with VCFS carry only one copy of the COMT allele on their intact chromosome. It has been suggested that dopaminergic neurotransmission in VCFS is compromised, thereby increasing susceptibility for psychosis (Dunham *et al.* 1992). Therefore studying the COMT *Val¹⁵⁸Met* polymorphism in VCFS may increase our understanding of brain development and predisposition for psychosis in this population.

An increasing number of studies have investigated the effects of the COMT *Val¹⁵⁸Met* polymorphism on cognitive function in people with VCFS. Some have reported that in VCFS children the low-activity *Met* allele is associated with better performance on prefrontal-dependent cognitive tasks (Bearden *et al.* 2004; Shashi *et al.* 2006). However Glaser *et al.* (2006) reported no difference on tasks of executive function between *Met*-hemizygous and *Val*-hemizygous groups in a combined children/adult VCFS population. In addition the COMT *Val¹⁵⁸Met* polymorphism may affect overall cognitive ability. For example, some studies have reported that the *Met* allele is associated with a decline in verbal IQ (Gothelf *et al.* 2005), and poorer performance on a task of language expression and spatial working memory (Baker *et al.* 2005); whereas others found the *Met* allele to be associated with higher verbal IQ (Shashi *et al.* 2006). Thus the effect of COMT *Val¹⁵⁸Met* polymorphism and cognitive function in people with VCFS is not easily understood, and the discrepancy in results of the reported studies may have arisen because of differences in dopaminergic neurotransmission due to developmental changes during adolescence, since both children, adolescents, and young adults were included in the study populations (Wahlstrom *et al.* 2007). The only study to date on COMT *Val¹⁵⁸Met* polymorphism in adults with VCFS showed that those with the *Met* allele displayed more severe excitement symptoms, and worse performance on theory of mind, Trails B, olfactory identification, communication and social functioning compared with those with the *Val* allele (Bassett *et al.* 2007). However, thus far there is no evidence that the COMT *Val¹⁵⁸Met* polymorphism affects the prevalence of schizophrenia in adults with VCFS (Bassett *et al.* 2007; Murphy *et al.* 1999).

The effects of the COMT *Val¹⁵⁸Met* polymorphism on brain anatomy in people with VCFS have been less frequently reported, but Gothelf *et al.* (2005) found the *Met* allele was associated with a smaller frontal lobe volume. Others, however (Kates *et al.* 2006), did not

find an effect of the COMT *Val¹⁵⁸Met* polymorphism alone, but did observe a gender-moderated effect of COMT genotype on both frontal lobe anatomy but not on frontal-lobe-dependent cognitive tasks.

Thus to date there have been relatively few studies on the effect of COMT *Val¹⁵⁸Met* polymorphism in people with VCFS, and those which are available focused on children and adolescents. Hence, it is unclear how the COMT *Val¹⁵⁸Met* polymorphism affects brain anatomy and cognitive function in adults with VCFS – and especially in adults. It has been suggested that PFC dopamine levels peak during adolescence and decline thereafter (Andersen *et al.* 1997), and that COMT activity reaches its optimal levels in early adulthood in the PFC, which might explain the discrepancies in studies on COMT *Val¹⁵⁸Met* polymorphism in children, adolescents and young adults with VCFS (Tunbridge *et al.* 2006). We previously reported on brain anatomy and cognitive function in an adult VCFS population (van Amelsvoort *et al.* 2004a,b). We have retrospectively obtained COMT genotype data from this sample. Here, we explore the relationship between COMT *Val¹⁵⁸Met* polymorphism and (1) brain morphometry and (2) measures of intelligence, attention, working memory, visuospatial function and memory in adults with VCFS. Because COMT is also expressed in abundance in non-frontal brain regions (Tunbridge *et al.* 2006), we tested the hypothesis that in adults with VCFS, genetic variation in COMT activity is associated with anatomical differences in frontal brain regions but also in other regions including cerebellum, regions which have been implicated in VCFS in previous studies. We further hypothesized that, in agreement with Bassett *et al.* (2007), in adults with VCFS those who were *Val*-hemizygous would perform better on frontal cognitive tests.

Method

Subjects

The study was approved by the local Ethics Committee and after complete description of the study to the subjects, written informed consent was obtained from themselves and/or their carers. All subjects were screened for medical conditions affecting brain functioning using a semi-structured clinical interview and routine clinical blood tests. To establish a DSM-IV (APA, 1994) diagnosis a semi-structured psychiatric interview was performed [Schedule for Clinical Assessment in Neuropsychiatry (SCAN); Wing *et al.* 1990] as described elsewhere (Murphy *et al.* 1999). We included 26 subjects with clinical features of VCFS and an established 22q11 deletion who have participated in a previously reported study on brain anatomy

and/or cognitive function in VCFS (Van Amelsvoort *et al.* 2004a,b). Twelve subjects had psychosis, five *val*-hemizygotes and seven *met*-hemizygotes, were clinically stable and taking antipsychotic medication. No other psychiatric disorders were present in the 26 subjects with VCFS.

Genetics

Blood samples were collected from 26 participants, and DNA was extracted. Fluorescence *in situ* hybridization (FISH) was used to detect a 22q11 deletion (Oncor Inc., Gaithersburg, MD). The COMT Val¹⁵⁸Met polymorphism was genotyped using the SNaPshot technique of single base extension (Applied Biosystems, Foster City, CA, USA) according to manufacturer's instructions. The initial PCR reaction was performed using a Touchdown-PCR-protocol, with the following primers: forward: 5'-ACTGTGGCT-ACTCAGCTGTG-3' and reverse: 5'-CCTTTTCCA-GGCTGACAA-3'. The allele at the SNP position was determined by use of a 30 bp extension primer (5'-ATCACCCAGCGGATGGTGGATTTTCGCTGGC-3'). All alleles were resolved on an ABI 3100 sequencer (Applied Biosystems, Foster City, CA, USA).

Magnetic resonance imaging acquisition and measurements

Magnetic resonance imaging (MRI) of the brain was performed on a 1.5-T MRI system (Signa; General Electric Co, Milwaukee, WI, USA) at the Maudsley Hospital, London, UK. A coronal volumetric spoiled gradient acquisition in the steady state dataset covering the whole head was acquired (repetition time, 13.8 ms; echo time, 2.8 ms; 124 sections; 1.5-mm section thickness). This dataset was used to perform manual tracing of brain volumes, using Measure software (Barta *et al.* 1997). Volumetric analysis by manual tracing was performed for total intracranial volume, caudate, putamen, hippocampus, amygdala, frontal, occipito-parietal and temporal lobes, cerebellum and ventricular and peripheral cerebrospinal fluid (CSF) by means of region of interest boundaries as previously described (Murphy *et al.* 1992, 1993a,b; Van Amelsvoort *et al.* 2001, 2004a). The volume of each region was calculated by multiplying the summed pixel cross-sectional areas by section thickness. Intra-rater and inter-rater reliabilities (range, 0.90–0.99) were determined by intra-class correlation computation for all brain regions traced by two operators and were highly significant ($F > 4.0$ and $p < 0.002$).

In addition, for determination of grey and white matter densities we carried out voxel based morphometry on a second dataset that we acquired in the

same individuals: a whole brain near axial dual-echo fast spin-echo (FSE) dataset aligned with the anterior commissure (AC)–posterior commissure (PC) plane [relaxation time (TR)=4000 ms, effective echo time (TE)=20 and 85 ms, 3 mm slice thickness]. This dataset was used to examine between-group differences in grey and white matter volume using a previously published methodology (Sigmundsson *et al.* 2001; McAlonan *et al.* 2002; Van Amelsvoort *et al.* 2004a). Voxels representing extracerebral tissue were automatically set to zero (Suckling *et al.* 1999a) and the probability of each intracerebral voxel belonging to grey matter, white matter, CSF, or dura/vasculature tissue classes was then estimated by a modified fuzzy clustering algorithm (Suckling *et al.* 1999b). On the basis of prior results, we equated these probabilities to the proportional volumes of each tissue class in the often heterogeneous volume of tissue represented by each voxel (Bullmore *et al.* 1995). Thus, for example, if the probability of grey matter class membership was 0.8 for a given voxel, it was assumed that 80% of the tissue represented by that voxel was grey matter. Because the voxel size was predetermined (2.2 mm³), we then estimated the volume in millilitres of grey matter, white matter and CSF in each voxel. Summing these voxel tissue class volumes over all intracerebral voxels yielded global tissue class volumes. To allow estimation of between-group differences at each intracerebral voxel (spatial extent statistics), the short echo (proton-density-weighted) FSE images were co-registered using an affine transformation (Press *et al.* 1992; Brammer *et al.* 1997) with a template image in the coordinate system of standard space as defined by Talairach & Tournoux (1988). This individually estimated transformation was then applied to each of that subject's grey and white tissue probability maps.

Neuropsychological testing

Overall intellectual functioning, memory, visuospatial and perceptual ability, executive functioning, and attention were measured. Details of the tests employed have been described elsewhere (Van Amelsvoort *et al.* 2004b) but the battery was chosen on suitability for a learning disabled population and addressing measures of overall IQ, attention, memory, executive function and visuospatial function. The test battery included: a short version of the Wechsler Adult Intelligence Scale – Revised (WAIS-R; Wechsler, 1981) consisting of five subtests (Vocabulary, Comprehension, Similarities, Block Design and Object Assembly); the Doors and People Test of Visual and Verbal Recall and Recognition (Baddeley *et al.* 1994); two subtests from the Wechsler Memory Scale – Revised (WMS-R; Wechsler, 1987): the Logical Memory (immediate and

delayed recall of two short stories) and Paired Associates (immediate and delayed recall of matched pairs) subtests; the Visual Space and Object Perception Battery (VOSP; Warrington & James, 1991); Computerised Tower of London Task (3-D CTL Test) (Shallice, 1982; Morris *et al.* 1988, 1990); Computerised Executive Golf Task (Morris *et al.* 1988; Baker *et al.* 1996); Controlled Oral Word Association Test (Benton & Hamsher, 1976); Weigl test (Goldstein & Scheerer, 1941); and the Continuous Performance Test (Conner, 1995).

Statistics

The study population was divided into two groups according to COMT genotype: *val*-hemizygous and *met*-hemizygous VCFS subjects. Differences in age, and IQ between the two groups were compared using a *t* test; differences in gender and psychiatric diagnosis using a χ^2 test.

Manually traced brain volumes, computerized tissue class volumes, and cognitive variables

Data were analysed with SPSS 12.0 for Windows (SPSS Inc., Chicago, IL). COMT effects on total and regional brain volumes, total tissue class volumes and cognitive variables were examined using a univariate analysis of covariance (ANCOVA) using COMT genotype, and gender as fixed factors and age and total intracranial volume (in case of MRI data) as covariates. We subsequently employed Bonferroni corrections for multiple testing using $p < 0.0045$ for the manually traced volumes, $p < 0.017$ for the tissue class volumes, and $p < 0.0011$ for the cognitive variables as levels of statistical significance (two-tailed).

Analysis of MRI data using computerized voxel-wise analysis

Between-group differences in grey and white matter were localized by fitting an appropriate GLM at each intracerebral voxel. Inference was via a permutation distribution of spatial extent statistics with significance levels set to control for multiple comparisons by having less than one estimated false positive region (cluster) across the image ($p < 0.001$). In brief, the processing proceeded as follows. Maps of the standardized GLM model coefficient of interest (group) at each voxel were thresholded such that only voxels with probability < 0.05 were retained. The sum of voxelwise statistics for each three-dimensional suprathreshold cluster was the test statistic, the sign indicating a relative excess or deficit in local tissue density. Significance testing of the clusters was performed using a null distribution of this test statistic similarly obtained after repeatedly randomly

Table 1. Characteristics of 26 adults with velo-cardio-facial syndrome

	<i>Val</i> (n=14)	<i>Met</i> (n=12)	<i>p</i>
Age (yr \pm S.D.)	30.3 (10.6)	37.33 (10.6)	0.10
Gender, M/F	6/8	3/9	0.43
Psychosis	5	7	0.43
FSIQ (\pm S.D.)	71.7 (10.6)	72.32 (13.2)	0.90




Fig. 1. Mean volumes (ml) of frontal lobes in *Met* hemizygous and *Val* hemizygous velo-cardio-facial syndrome people.

permuting the relevant factor in the GLM and refitting of the model (Bullmore *et al.* 1999).

Results

We studied 26 adults with VCFS. Data on age, gender, psychiatric diagnosis and intellectual function are presented in Table 1. Age, psychiatric diagnosis, full scale IQ (FSIQ) and gender did not differ between *val*- and *met*-hemizygous groups. However, the women in our sample were significantly older than the men; therefore age was introduced as a covariate in our analyses. Psychiatric diagnosis and FSIQ did not vary as a function of gender. We were unable to acquire MRI scans on seven subjects and neuropsychological data on three subjects. There were no differences in age, psychiatric diagnosis, FSIQ or gender between *val*- and *met*-hemizygous subgroups with the MRI or the cognitive data.

COMT effects on manually traced brain volumes

After accounting for total intracranial volume and age there was a significant COMT effect on volumes of frontal ($F = 30.09$, $p = 0.000$), and temporal lobes ($F = 10.17$, $p = 0.007$), cerebellum ($F = 8.30$, $p = 0.013$); and peripheral CSF (CSF excluding ventricles) ($F = 10.84$, $p = 0.006$) (Fig. 1). There were no gender effects or COMT \times gender interactions. No significant

Table 2. Manually traced brain and computerized tissue class volumes in *Val*-hemizygous subjects and *Met*-hemizygous subjects (corrected for intracranial volume)

Brain structure	Group mean (s.d.), ml		<i>p</i> value	<i>F</i>	df
	<i>Val</i> (<i>n</i> =10)	<i>Met</i> (<i>n</i> =9)			
Total intracranial volume	1270.639 (50.074)	1314.991 (55.782)	0.571	0.337	1, 14
Frontal lobes	499.154 (5.479)	454.957 (6.216)	0.000**	30.09	1, 13
Occipitoparietal lobes	365.178 (11.253)	363.977 (12.766)	0.946	0.005	1, 13
Temporal lobes	136.855 (5.177)	111.586 (5.873)	0.007*	10.17	1, 13
Putamen	6.632 (0.468)	7.140 (0.497)	0.473	0.548	1, 12
Caudate	7.967 (0.422)	7.588 (0.449)	0.552	0.375	1, 12
Peripheral CSF	126.961 (11.793)	184.760 (13.377)	0.006*	10.84	1, 13
Hippocampus	5.339 (0.307)	4.693 (0.344)	0.197	1.888	1, 11
Amygdala	4.467 (0.329)	4.032 (0.368)	0.406	0.746	1, 11
Cerebellum	109.859 (3.621)	95.740 (4.108)	0.013*	8.30	1, 13
Ventricles	0.995 (0.543)	2.623 (0.616)	0.074	3.768	1, 13
Tissue class volumes					
Total grey matter	599.90 (81.90)	560 (101.49)	0.31	1.15	1, 11
Total white matter	577.12 (77.65)	576.26 (70.28)	0.38	0.82	1, 11
Total CSF	158.26 (42.42)	216.20 (105.37)	0.028*	6.38	1, 11

CSF, Cerebrospinal fluid.

* $p < 0.05$, ** $p < 0.01$.

COMT, gender or COMT \times gender interactions were found in the volume of any other brain region or ventricles (Table 2). After correction for multiple comparisons ($p < 0.0045$) only frontal lobe volumes remained significantly different, with larger volumes in the *val*-hemizygous group compared with the *met*-hemizygous group.

COMT effects on total tissue class volumes

Having accounted for total intracranial volume and age, there was no gender effect on total CSF, grey, and white matter volume. There was a significant COMT effect on total CSF volume ($p = 0.028$, $F = 6.38$). In addition there was a significant COMT \times gender interaction on total white matter volume ($p = 0.015$, $F = 8.31$) and total CSF volume ($p = 0.003$, $F = 14.86$) (Table 2). After correction for multiple comparisons ($p < 0.017$) only a COMT \times gender interaction on CSF remained significant.

COMT effects on regional grey matter distribution

There was a significant difference between the VCFS *Val*-hemizygous and *Met*-hemizygous group in grey matter volume at four spatially extensive 3D voxel clusters. The *Met*-hemizygous group had a significant reduced grey matter volume in left parahippocampal gyrus and brainstem, and bilaterally in cerebellum. There were no significant clusters of

Table 3. Regional differences in grey and white matter volume

Cerebral region	<i>n</i>	Talairach coordinates (mm)			Side
		x	y	z	
Grey matter					
<i>Met</i> < <i>Val</i> Deficit					
Cerebellum	2948	9.3	-65.1	-19.4	R
Cerebellum	116	-34.3	59.3	-16.6	L
Brainstem	10	-11.0	-36.8	-26.0	L
Parahippocampal gyrus	208	-21.9	46.5	-4.8	L
White matter					
<i>Met</i> > <i>Val</i> Excess					
Cerebellum	4776	7.2	-57.8	21.4	R

n, Number of voxels in each cluster.The cluster-wise probability is $p = 0.001$.

excess grey matter in the *Met*-hemizygous group (Table 3, Fig. 2).

COMT effects on regional white matter distribution

The *Met*-hemizygous group had a significantly increased white matter volume in cerebellum. There were no clusters of reduced white matter in the *Met*-hemizygous group (Table 3, Fig. 3).




Fig. 2. Relative deficits (blue) and excesses (red) in grey matter volume in *Met* hemizygous compared with *Val* hemizygous VCFS people. The maps are oriented with the right side of the brain shown on the left side of each panel. The z coordinate for each row of axial slices in the standard space of Talairach & Tournoux (1988) is given in millimetres.

COMT effects on cognitive variables

Accounting for age, there were significant COMT effects on two subtests of the Continuous Performance Test: Variability of Standard Error ($p=0.031$, $F=5.64$), and Standard Error Block Change ($p=0.023$, $F=6.42$). Also there was a significant COMT effect on total CPT index score ($p=0.019$, $F=6.85$) and on the minimal amount of moves needed (MAM) in the Tower of London task ($p=0.019$, $F=6.96$) (Table 4).

Significant COMT \times gender interactions were found for Standard Error Block Change of the CPT ($p=0.011$, $F=8.42$), Strategy Formation ($p=0.009$, $F=8.76$) and Between Search Errors ($p=0.017$, $F=6.95$) on the Golf Spatial Working Memory task, MAM of the Tower of London task ($p=0.006$, $F=10.16$), Delayed Logical Memory of Wechsler memory Scale revised ($p=0.021$, $F=6.78$), Object Discrimination subtest of VOSP ($p=0.018$, $F=6.79$). After Bonferroni correction for multiple comparisons none of the findings remained significant.

Discussion

We found that in adults with VCFS, the COMT *Val*¹⁵⁸*Met* polymorphism is associated with significant differences in brain anatomy affecting frontal lobe volume and grey matter density in cerebellum, brainstem and parahippocampal gyrus, and white matter density in cerebellum.

However, our study has limitations. These include the relatively small sample size with insufficient power to look for interactions, and the multiple testing we carried out. Nevertheless, corrections for multiple comparisons were applied for all anatomical and cognitive variables. Also it is unlikely that differences in age, psychiatric comorbidity, and gender can fully account for our findings (as these factors did not differ between *Val*- and *Met*-hemizygous groups). In addition, we controlled for gender and age in the statistical analysis. Also, as noted by others (Chow *et al.* 1999), the effect size for structural brain abnormalities in people with VCFS is relatively large. Nevertheless




Fig. 3. Relative deficits (blue) and excesses (red) in white matter volume in *Met* hemizygous compared with *Val* hemizygous velo-cardio-facial syndrome people. The maps are oriented with the right side of the brain shown on the left side of each panel. The z coordinate for each row of axial slices in the standard space of Talairach & Tournoux (1988) is given in millimetres.

our findings should be interpreted with caution and further studies of a longitudinal nature using larger samples of people with and without VCFS are required.

Our study is the first to demonstrate that the effect of the COMT *Val*¹⁵⁸*Met* polymorphism is not limited to frontal brain regions in people with VCFS. Because of the paucity of dopamine transporters in the prefrontal cortex, COMT is responsible for dopamine degradation in this region (Chen *et al.* 2004). Therefore, effects of the COMT *Val*¹⁵⁸*Met* polymorphism on brain volume have previously been hypothesized to be most pronounced in frontal cortex, and this has been the focus of most studies on brain anatomy and COMT *Val*¹⁵⁸*Met* polymorphism (Ho *et al.* 2005). Our results, using a whole brain anatomy approach, suggest that COMT *Val*¹⁵⁸*Met* polymorphism may affect other brain regions as well, including cerebellum, a region particularly compromised in VCFS (Van Amelsvoort *et al.*

2004a; Simon *et al.* 2005; Campbell *et al.* 2006). COMT mRNA is expressed in several human brain regions including the frontal, temporal, and parietal lobes, and cerebellum, amygdala, putamen, thalamus and spinal cord (Hong *et al.* 1998). Therefore it is not surprising that the *Val*¹⁵⁸*Met* polymorphism affects other brain regions than the PFC, which supports our findings. Also, COMT may be crucial for dopaminergic degradation in PFC, but more relevant for regulating norepinephrine (another substrate of COMT) in other brain regions. Both these neurochemical systems affect brain development and function. Thus COMT *Val*¹⁵⁸*Met* effects on brain anatomy may not only reflect differences in dopamine degradation but also differences in (nor)epinephrine metabolism (Parini *et al.* 1988).

The underlying biological mechanism for our finding of larger bulk frontal lobe volumes and increased grey matter densities in *Val*-hemizygotes is unclear.

Table 4. Neuropsychological test scores for velo-cardio-facial syndrome Met and Val subgroups

	Val	Met	p value	F	df
Intellectual functioning					
WAIS-R	(n = 12)	(n = 11)			1, 18
Vocabulary	4.58 ± 2.19	4.18 ± 2.23	0.27	1.27	
Comprehension	4.42 ± 1.44	4.45 ± 1.51	0.33	1.00	
Similarities	5.58 ± 0.79	4.91 ± 1.97	0.14	2.36	
Block Design	5.75 ± 3.38	6.00 ± 2.90	0.58	0.32	
Object Assembly	4.83 ± 2.66	4.64 ± 3.04	0.41	0.71	
VIQ	73.25 ± 7.59	71.36 ± 9.80	0.22	1.61	
PIQ	73.58 ± 15.97	77.36 ± 18.42	0.24	0.63	
FSIQ	71.67 ± 10.55	72.32 ± 13.21	0.38	0.82	
V-P Discrepancy	-0.33 ± 12.57	-6.0 ± 12.18	0.80	0.06	
Memory					
Doors and People	(n = 12)	(n = 9)			1, 16
People	5.58 ± 3.32	5.22 ± 2.73	0.25	1.44	
Doors	3.92 ± 2.02	5.78 ± 3.77	0.93	0.01	
Shapes	5.08 ± 4.23	6.11 ± 3.92	0.68	0.18	
Names	6.67 ± 5.23	6.44 ± 3.84	0.39	0.80	
Overall	3.33 ± 3.55	4.88 ± 4.02	0.67	0.20	
Visual	4.27 ± 2.49	6.38 ± 3.54	0.96	0.00	
Verbal	5.67 ± 4.56	6.25 ± 3.01	0.50	0.47	
Recall	4.67 ± 3.28	5.62 ± 2.77	0.37	0.86	
Recognition	4.64 ± 3.38	6.13 ± 4.12	0.81	0.06	
Overall Forgetting	10.50 ± 4.54	11.22 ± 2.82	0.78	0.08	
WMS-R	(n = 11)	(n = 9)			1, 15
Logical Memory Immediate	23.09 ± 11.26	20.89 ± 13.0	0.19	1.90	
Logical Memory Delayed	6.18 ± 5.83	5.50 ± 4.84	0.14	2.49	
Paired Associates Immediate	12.36 ± 5.45	13.89 ± 6.86	0.76	0.09	
Paired Associates Delayed	10.0 ± 5.51	12.00 ± 5.45	0.97	0.00	
Verbal Memory Index	67.82 ± 16.6	71.13 ± 12.92	0.48	0.08	
Visuospatial/perceptual functioning					
VOSP	(n = 12)	(n = 11)			1, 18
Incomplete Letters	19.33 ± 0.65	19.00 ± 2.10	0.26	1.33	
Silhouettes	13.83 ± 2.79	17.09 ± 4.91	0.13	2.49	
Object Decision	13.67 ± 3.17	14.82 ± 4.45	0.79	0.07	
Progressive Silhouettes	12.75 ± 2.60	11.09 ± 2.59	0.59	0.30	
Dot Counting	9.75 ± 0.45	9.82 ± 0.41	0.95	0.00	
Position Discrimination	16.00 ± 3.79	17.55 ± 2.77	0.58	0.34	
Number Location	6.58 ± 3.50	6.09 ± 3.18	0.53	0.40	
Cube Analysis	6.83 ± 2.86	6.18 ± 2.52	0.45	0.59	
Number of Passes	4.75 ± 1.29	5.55 ± 1.75	0.62	0.25	
Spatial Working Memory					
Executive Golf Task	(n = 12)	(n = 11)			1, 18
Within Search Errors	0.60 ± 0.85	0.91 ± 0.91	0.38	0.82	
Between Search Errors	4.63 ± 2.16	5.69 ± 2.64	0.10	3.07	
Strategy Formation	12.35 ± 1.16	12.55 ± 1.12	0.32	1.06	
Planning and Problem-Solving					
Tower of London	(n = 12)	(n = 9)			1, 16
Moves Above Minimum	2.74 ± 1.04	2.02 ± 1.35	0.03*	5.89	
Planning (s)	4.09 ± 1.94	4.56 ± 2.41	0.24	1.51	
Verbal fluency					
Total	(n = 12)	(n = 7)			1, 14
	22.17 ± 9.52	23.43 ± 15.70	0.25	1.47	
Weigl					
(n = 12)	(n = 10)			χ^2	
Sorting	100%	90%	0.45	1.59	
Shifting	33.3%	60%	0.53	1.28	

Table 4 (cont.)

	<i>Val</i>	<i>Met</i>	<i>p</i> value	<i>F</i>	df
Attention					
CPT (<i>T</i> scores)	(<i>n</i> = 12)	(<i>n</i> = 9)			1, 16
Omission Errors (percentiles)	96.85 ± 4.44	95.74 ± 6.36	0.71	0.14	
Hit Reaction Times (<i>T</i> scores)	40.70 ± 7.32	43.97 ± 8.39	0.19	1.91	
Variability of s.e.'s (<i>T</i> scores)	69.11 ± 14.78	75.02 ± 11.99	0.02*	6.20	
Hit s.e. Block Change (<i>T</i> scores)	59.76 ± 14.66	45.41 ± 22.95	0.02*	7.11	
Commission Errors (<i>T</i> scores)	62.05 ± 13.21	61.23 ± 14.14	0.67	0.19	
Perceptual Sensitivity (<i>T</i> scores)	71.28 ± 11.32	69.58 ± 11.01	0.59	0.31	
Risk Taking (<i>T</i> scores)	83.28 ± 15.30	82.22 ± 18.41	0.46	0.58	
CPT Index Score	10.50 ± 4.19	12.50 ± 3.99	0.04*	5.28	

Values are group means ± s.d.

* $p < 0.05$.

One possible explanation could be differences in brain maturation between the two genotype groups. Qualitative imaging studies have reported brain anomalies in children and adults with VCFS, for example agenesis of the corpus callosum, white matter hyperintensities and septum pellucidum abnormalities (Mitnick *et al.* 1994; Chow *et al.* 1999; Van Amelsvoort *et al.* 2001). Quantitative imaging studies have also reported evidence for disturbed brain maturation in people with VCFS. For example, several studies of brain anatomy have reported that people with VCFS compared with controls often have abnormalities in either frontal lobe anatomy (with subtle differences in grey/white matter composition) and/or posterior brain regions including cerebellum (Van Amelsvoort *et al.* 2004a; Simon *et al.* 2005; Campbell *et al.* 2006). Normal brain maturation starts from posterior/inferior brain regions and takes place last in frontal regions (Sowell *et al.* 1999) and is associated with an increase in frontal white matter and a decrease in grey matter during adolescence (Nagy *et al.* 2004). Abnormalities in brain maturation (e.g. programmed cell death) could therefore lead to differences in the relative proportions of grey and white matter and hence frontal lobe volume. Since normal brain maturation is accompanied by increasingly efficient cognitive processing (Levy & Goldman-Rakic, 2000), the finding of decreased efficiency of prefrontal cognitive processing in those possessing the *Val* allele (Egan *et al.* 2001) perhaps suggests that possessing the *Val* allele may slow down the maturation process – and hence the relative proportion of grey and white matter and/or bulk volume of frontal regions. A so-called anterior-posterior dichotomy in anatomical differences between people with VCFS and controls has been suggested (Kates *et al.* 2001; Van Amelsvoort *et al.* 2004a; Simon *et al.* 2005), and could also explain differences in brain

anatomy within the VCFS population between those hemizygous for *Val* or *Met*. How COMT haploinsufficiency in VCFS affects the dopaminergic system is still unknown, although a preliminary open study by Graf *et al.* (2001) suggested increased baseline brain dopamine levels in three of four patients with VCFS and the *Met*-allele, as measured by the level of a catecholamine metabolite, homovanillic acid, in cerebrospinal fluid compared with calibrated standards and laboratory controls. These preliminary findings suggest that haploinsufficiency in COMT is associated with dysregulation of dopaminergic systems. Results from animal and human studies suggest that dopamine has a trophic action during early brain maturation and later influences prefrontal cortical specification (Nieoullon, 2002). Therefore, differences in dopaminergic clearance as a result of *Val*- or *Met*-hemizygosity could result in differences in (frontal) brain maturation because of differences in dopamine signalling during early brain development. Our finding of relative cerebellar grey matter increase and white matter decrease in the *Val*-hemizygotes suggests that dynamic interplay between grey and white matter tissue composition might take place during brain maturation with grey matter volume decrease due to pruning and white matter volume increase during myelination. These findings further support the hypothesis that brain maturation may be slower in *val*-hemizygotes than in *met*-hemizygotes.

Our findings on cognitive performance did not survive Bonferroni corrections for multiple comparisons. This is probably partially due to our small sample size. The current literature suggests that cognitive tasks classically thought to be measures of 'frontal' regions are modulated by the COMT polymorphism. As mentioned before the results on COMT polymorphism and cognitive function in VCFS are

inconsistent also because various study groups have used different cognitive measures and study populations included people of different ages. Future studies with larger sample sizes and of a longitudinal nature should clarify the role of the COMT polymorphism on specific cognitive tasks during lifespan in VCFS.

In conclusion, our results suggest that genetic variation in COMT activity affects brain anatomy in adults with VCFS, and this extends outside frontal brain regions. This suggests variation in COMT activity is implicated in brain development in VCFS. Future studies are required to investigate changes across the lifespan.

Declaration of Interest

None.

References

- Akil M, Kolachana BS, Rothmond DA, Hyde TM, Weinberger DR, Kleinman JE (2003). Catechol-O-methyltransferase genotype and dopamine regulation in the human brain. *Journal of Neuroscience* **23**, 2008–2013.
- Andersen SL, Dumont NL, Teicher MH (1997). Developmental differences in dopamine synthesis inhibition by (+/-)-7-OH-DPAT. *Naunyn-Schmiedeberg's Arch Pharmacol* **356**, 173–181.
- APA (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn (DSM-IV). American Psychiatric Association: Washington, DC.
- Baddeley AD, Emslie H, Nimmo-Smith I (1994). *The Doors and People Test: a Test of Visual and Verbal Recall and Recognition*. Thames Valley Test Company: Bury St. Edmunds.
- Baker K, Baldeweg T, Sivagnanasundaram S, Scambler P, Skuse D (2005). COMT Val108/158 Met modifies mismatch negativity and cognitive function in 22q11 deletion syndrome. *Biological Psychiatry* **58**, 23–31.
- Baker SC, Rogers RD, Owen AM, Frith CD, Dolan RJ, Frackowiak RS, Robbins TW (1996). Neural systems engaged by planning: a PET study of the Tower of London task. *Neuropsychologia* **34**, 515–526.
- Barta PE, Dhingra L, Royall R, Schwartz E (1997). Improving stereological estimates for the volume of structures identified in three-dimensional arrays of spatial data. *Journal of Neuroscience Methods* **75**, 111–118.
- Bassett AS, Caluseriu O, Weksberg R, Young DA, Chow EW (2007). Catechol-O-methyl transferase and expression of schizophrenia in 73 adults with 22q11 deletion syndrome. *Biological Psychiatry*. doi:10.1016/j.biopsych.2006.07.038.
- Bearden CE, Jawad AF, Lynch DR, Sokol S, Kanes SJ, McDonald-McGinn DM, Saitta SC, Harris SE, Moss E, Wang PP, Zackai E, Emanuel BS, Simon TJ (2004). Effects of a functional COMT polymorphism on prefrontal cognitive function in patients with 22q11.2 deletion syndrome. *American Journal of Psychiatry* **161**, 1700–1702.
- Benton AL, Hamsher KS (1976). *Multilingual Aphasia Examination*. University of Iowa: Iowa City, IA.
- Brammer MJ, Bullmore ET, Simmons A, Williams SC, Grasby PM, Howard RJ, Woodruff PW, Rabe-Hesketh S (1997). Generic brain activation mapping in functional magnetic resonance imaging: a nonparametric approach. *Magnetic Resonance Imaging* **15**, 763–770.
- Bullmore E, Brammer M, Rouleau G, Everitt B, Simmons A, Sharma T, Frangou S, Murray R, Dunn G (1995). Computerized brain tissue classification of magnetic resonance images: a new approach to the problem of partial volume artifact. *Neuroimage* **2**, 133–147.
- Bullmore ET, Suckling J, Overmeyer S, Rabe-Hesketh S, Taylor E, Brammer MJ (1999). Global, voxel, and cluster tests, by theory and permutation, for a difference between two groups of structural MR images of the brain. *IEEE Transactions on Medical Imaging* **18**, 32–42.
- Campbell LE, Daly E, Toal F, Stevens A, Azuma R, Catani M, Ng V, Van Amelsvoort T, Chitnis X, Cutter W, Murphy DG, Murphy KC (2006). Brain and behaviour in children with 22q11.2 deletion syndrome: a volumetric and voxel-based morphometry MRI study. *Brain* **129**, 1218–1228.
- Chen J, Lipska BK, Halim N, Ma QD, Matsumoto M, Melhem S, Kolachana BS, Hyde TM, Herman MM, Apud J, Egan MF, Kleinman JE, Weinberger DR (2004). Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain. *American Journal of Human Genetics* **75**, 807–821.
- Chow EW, Mikulis DJ, Zipursky RB, Scutt LE, Weksberg R, Bassett AS (1999). Qualitative MRI findings in adults with 22q11 deletion syndrome and schizophrenia. *Biological Psychiatry* **46**, 1436–1442.
- Conner CK (1995). *Continuous Performance Test*. Multi-Health Systems Inc.: Toronto, Canada.
- Dunham I, Collins J, Wadey R, Scambler P (1992). Possible role for COMT in psychosis associated with velo-cardio-facial syndrome. *Lancet* **340** (8831), 1361–1362.
- Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE, Goldman D, Weinberger DR (2001). Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proceedings of the National Academy of Sciences USA* **98**, 6917–6922.
- Glaser B, Debbane M, Hinard C, Morris MA, Dahoun SP, Antonarakis SE, Eliez S (2006). No evidence for an effect of COMT Val158Met genotype on executive function in patients with 22q11 deletion syndrome. *American Journal of Psychiatry* **163**, 537–539.
- Goldstein KH, Scheerer M (1941). Abstract and concrete behaviour: an experimental study with special tests. *Psychological Monographs* **53**, no. 2.
- Gothelf D, Eliez S, Thompson T, Hinard C, Penniman L, Feinstein C, Kwon H, Jin S, Jo B, Antonarakis SE, Morris MA, Reiss AL (2005). COMT genotype predicts longitudinal cognitive decline and psychosis in 22q11.2 deletion syndrome. *Nature Neuroscience* **8**, 1500–1502.
- Graf WD, Unis AS, Yates CM, Sulzbacher S, Dinulos MB, Jack RM, Dugaw KA, Paddock MN, Parson WW (2001).

- Catecholamines in patients with 22q11.2 deletion syndrome and the low-activity COMT polymorphism. *Neurology* 57, 410–416.
- Ho BC, Wassink TH, O'Leary DS, Sheffield VC, Andreasen NC** (2005). Catechol-O-methyl transferase Val158Met gene polymorphism in schizophrenia: working memory, frontal lobe MRI morphology and frontal cerebral blood flow. *Molecular Psychiatry* 10, 287–298.
- Hong J, Shu-Leong H, Tao X, Lap-Ping Y** (1998). Distribution of catechol-O-methyltransferase expression in human central nervous system. *Neuroreport* 9, 2861–2864.
- Kates WR, Antshel KM, AbdulSabur N, Colgan D, Funke B, Fremont W, Higgins AM, Kucherlapati R, Shprintzen RJ** (2006). A gender-moderated effect of a functional COMT polymorphism on prefrontal brain morphology and function in velo-cardio-facial syndrome (22q11.2 deletion syndrome). *American Journal of Medical Genetics, B: Neuropsychiatric Genetics* 141, 274–280.
- Kates WR, Burnette CP, Jabs EW, Rutberg J, Murphy AM, Grados M, Geraghty M, Kaufmann WE, Pearlson GD** (2001). Regional cortical white matter reductions in velocardiofacial syndrome: a volumetric MRI analysis. *Biological Psychiatry* 49, 677–684.
- Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM** (1996). Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* 6, 243–250.
- Levy R, Goldman-Rakic PS** (2000). Segregation of working memory functions within the dorsolateral prefrontal cortex. *Experimental Brain Research* 133, 23–32.
- Malhotra AK, Kestler LJ, Mazzanti C, Bates JA, Goldberg T, Goldman D** (2002). A functional polymorphism in the COMT gene and performance on a test of prefrontal cognition. *American Journal of Psychiatry* 159, 652–654.
- McAlonan GM, Daly E, Kumari V, Critchley HD, Van Amelsvoort T, Suckling J, Simmons A, Sigmundsson T, Greenwood K, Russell A, Schmitz N, Happe F, Howlin P, Murphy DG** (2002). Brain anatomy and sensorimotor gating in Asperger's syndrome. *Brain* 125, 1594–1606.
- Meyer-Lindenberg A, Kohn PD, Kolachana B, Kippenhan S, McInerney-Leo A, Nussbaum R, Weinberger DR, Berman KF** (2005). Midbrain dopamine and prefrontal function in humans: interaction and modulation by COMT genotype. *Nature Neuroscience* 8, 594–596.
- Mitnick RJ, Bello JA, Shprintzen RJ** (1994). Brain anomalies in velo-cardio-facial syndrome. *American Journal of Medical Genetics* 54, 100–106.
- Morris RG, Downes JJ, Robbins TW** (1990). The nature of the dysexecutive syndrome in Parkinson's disease. In *Lines of Thinking* (ed. K.J. Gilhooly, M.T.G. Keane, R.H. Logie and G. Erdos), pp. 247–258. John Wiley: New York.
- Morris RG, Downes JJ, Sahakian BJ, Evenden JL, Heald A, Robbins TW** (1988). Planning and spatial working memory in Parkinson's disease. *Journal of Neurology, Neurosurgery & Psychiatry* 51, 757–766.
- Murphy DG, DeCarli CD, Daly E, Gillette JA, McIntosh AR, Haxby JV, Teichberg D, Schapiro MB, Rapoport SI, Horwitz B** (1993b). Volumetric magnetic resonance imaging in men with dementia of the Alzheimer type: correlations with disease severity. *Biological Psychiatry* 34, 612–621.
- Murphy DG, DeCarli C, Daly E, Haxby JV, Allen G, White BJ, McIntosh AR, Powell CM, Horwitz B, Rapoport SI** (1993a). X-chromosome effects on female brain: a magnetic resonance imaging study of Turner's syndrome. *Lancet* 342, 1197–1200.
- Murphy DG, DeCarli C, Schapiro MB, Rapoport SI, Horwitz B** (1992). Age-related differences in volumes of subcortical nuclei, brain matter, and cerebrospinal fluid in healthy men as measured with magnetic resonance imaging. *Archives of Neurology* 49, 839–845.
- Murphy KC, Jones LA, Owen MJ** (1999). High rates of schizophrenia in adults with velo-cardio-facial syndrome. *Archives of General Psychiatry* 56, 940–945.
- Nagy Z, Westerberg H, Klingberg T** (2004). Maturation of white matter is associated with the development of cognitive functions during childhood. *Journal of Cognitive Neurosciences* 16, 1227–1233.
- Nieoullon A** (2002). Dopamine and the regulation of cognition and attention. *Progress in Neurobiology* 67, 53–83.
- Papolos DF, Faedda GL, Veit S, Goldberg R, Morrow B, Kucherlapati R, Shprintzen RJ** (1996). Bipolar spectrum disorders in patients diagnosed with velo-cardio-facial syndrome: does a hemizygous deletion of chromosome 22q11 result in bipolar affective disorder? *American Journal of Psychiatry* 153, 1541–1547.
- Parini A, Coupry I, Laude D, Diop L, Vincent M, Sassard J, Dausse JP** (1988). Noradrenaline content and adrenergic receptors in kidney and heart of the prehypertensive and hypertensive Lyon rat strain. *American Journal of Hypertension* 1, 140–145.
- Press WH, Taukolsy SA, Vetterling WT, Flannery BP** (1992). *Numerical Recipes in C: The Art of Scientific Computing*, 2nd edn. Cambridge University Press: Cambridge.
- Shallice T** (1982). Specific impairments of planning. *Philosophical Transactions of the Royal Society (London)* 298, 199–209.
- Shashi V, Keshavan MS, Howard TD, Berry MN, Basehore MJ, Lewandowski E, Kwapil TR** (2006). Cognitive correlates of a functional COMT polymorphism in children with 22q11.2 deletion syndrome. *Clinical Genetics* 69, 234–238.
- Shprintzen RJ, Goldberg RB, Lewin ML, Sidoti EJ, Berkman MD, Argamaso RV, Young D** (1978). A new syndrome involving cleft palate, cardiac anomalies, typical facies, and learning disabilities: velo-cardio-facial syndrome. *Cleft Palate Journal* 15, 56–62.
- Sigmundsson T, Suckling J, Maier M, Williams S, Bullmore E, Greenwood K, Fukuda R, Ron M, Toone B** (2001). Structural abnormalities in frontal, temporal, and limbic regions and interconnecting white matter tracts in schizophrenic patients with prominent negative symptoms. *American Journal of Psychiatry* 158, 234–243.
- Simon TJ, Ding L, Bish JP, McDonald-McGinn DM, Zackai EH, Gee J** (2005). Volumetric, connective, and morphologic changes in the brains of children with chromosome 22q11.2 deletion syndrome: an integrative study. *Neuroimage* 25, 169–180.

- Smolka MN, Schumann G, Wrase J, Grusser SM, Flor H, Mann K, Braus DF, Goldman D, Buchel C, Heinz A** (2005). Catechol-O-methyltransferase val158met genotype affects processing of emotional stimuli in the amygdala and prefrontal cortex. *Journal of Neuroscience* **25**, 836–842.
- Sowell ER, Thompson PM, Holmes CJ, Jernigan TL, Toga AW** (1999). In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. *Nature Neuroscience* **2**, 859–861.
- Suckling J, Brammer MJ, Lingford-Hughes A, Bullmore ET** (1999a). Removal of extracerebral tissues in dual-echo magnetic resonance images via linear scale-space features. *Magnetic Resonance Imaging* **17**, 247–256.
- Suckling J, Sigmundsson T, Greenwood K, Bullmore ET** (1999b). A modified fuzzy clustering algorithm for operator independent brain tissue classification of dual echo MR images. *Magnetic Resonance Imaging* **17**, 1065–1076.
- Talairach J, Tournoux P** (1988). *Co-Planar Stereotaxic Atlas of the Human Brain*. Thieme Medical Publishers: New York.
- Tunbridge EM, Harrison PJ, Weinberger DR** (2006). Catechol-o-methyltransferase, cognition, and psychosis: Val(158)Met and beyond. *Biological Psychiatry* **60**, 141–151.
- Van Amelsvoort T, Daly E, Henry J, Robertson D, Ng V, Owen M, Murphy KC, Murphy DG** (2004a). Brain anatomy in adults with velocardiofacial syndrome with and without schizophrenia: preliminary results of a structural magnetic resonance imaging study. *Archives of General Psychiatry* **61**, 1085–1096.
- Van Amelsvoort T, Daly E, Robertson D, Suckling J, Ng V, Critchley H, Owen MJ, Henry J, Murphy KC, Murphy DG** (2001). Structural brain abnormalities associated with deletion at chromosome 22q11: quantitative neuroimaging study of adults with velo-cardio-facial syndrome. *British Journal of Psychiatry* **178**, 412–419.
- Van Amelsvoort T, Henry J, Morris R, Owen M, Linszen D, Murphy K, Murphy D** (2004b). Cognitive deficits associated with schizophrenia in velo-cardio-facial syndrome. *Schizophrenia Research* **70**, 223–232.
- Wahlstrom D, White T, Hooper CJ, Vrshek-Schallhorn S, Oetting WS, Brott MJ, Luciana M** (2007). Variations in the catechol O-methyltransferase polymorphism and prefrontally guided behaviors in adolescents. *Biological Psychiatry* **61**, 626.
- Warrington EK, James M** (1991). *The Visual Object and Space Perception Battery*. Thames Valley Test Company: Titchfield, UK.
- Wechsler D** (1981). *The Wechsler Adult Intelligence Scale – Revised*. Psychological Corporation: New York.
- Wechsler D** (1987). *Wechsler Memory Scale – Revised: Manual*. The Psychological Corporation: San Antonio, TX.
- Wing JK, Babor T, Brugha T, Burke J, Cooper JE, Giel R, Jablenski A, Regier D, Sartorius N** (1990). SCAN. Schedules for Clinical Assessment in Neuropsychiatry. *Archives of General Psychiatry* **47**, 589–593.