

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/81603/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Maier, Wolfgang D. , Howard, H. M., Smithies, R. H., Yang, S. H., Barnes, S.-J., O'Brien, H., Huhma, H. and Gardoll, S. 2015. Magmatic ore deposits in mafic-ultramafic intrusions of the Giles Event, Western Australia. *Ore Geology Reviews* 71 , pp. 405-436. 10.1016/j.oregeorev.2015.06.010

Publishers page: <http://dx.doi.org/10.1016/j.oregeorev.2015.06.010>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Magmatic ore deposits in mafic-ultramafic intrusions of the Giles Event, Western**
2 **Australia**

3 W.D. Maier¹, H.M. Howard², R.H. Smithies², S.H. Yang³, S-J. Barnes⁴, H. O'Brien⁵, H. Huhma⁵, and S.
4 Gardoll⁶

5 ¹School of Earth and Ocean Sciences, Cardiff University, Cardiff, Wales,UK

6 ²Geological Survey of Western Australia, Perth, Australia

7 ³Oulu Mining School, University of Oulu, Oulu 90014, Finland

8 ⁴Sciences de la Terre, Université du Québec á Chicoutimi, Chicoutimi G7H 2B1, Canada

9 ⁵Geological Survey of Finland (GTK), FI-02151 Espoo, Finland

10 ⁶Department of Applied Geology, Curtin University, GPO Box U1987, Perth WA 6845.

11

12 **Abstract**

13 More than 20 layered intrusions were emplaced at c. 1075 Ma across >100 000 km² in the
14 Mesoproterozoic Musgrave Province of central Australia as part of the c. 1090–1040 Ma Giles
15 Event of the the Warakurna Large Igneous Province (LIP). Some of the intrusions, including
16 Wingellina Hills, Pirntirri Mulari, The Wart, Ewarara, Kalka, Claude Hills, and Gosse Pile contain
17 thick ultramafic segments comprising wehrlite, harzburgite, and websterite. Other intrusions,
18 notably Hinckley Range, Michael Hills, and Murray Range, are essentially of olivine-gabbroitic
19 composition. Intrusions with substantial troctolitic portions comprise Morgan Range and
20 Cavenagh Range, as well as the Bell Rock, Blackstone, and Jameson–Finlayson ranges which are
21 tectonically dismembered blocks of an originally contiguous intrusion, here named Mantamaru,
22 with a strike length of >170 km and a width of > 20 km, constituting one of the world's largest
23 layered intrusions.

24 Over a time span of > 200 my, the Musgrave Province was affected by near continuous high-
25 temperature reworking under a primarily extensional regime. This began with the 1220–1150 Ma
26 intracratonic Musgrave Orogeny, characterized by ponding of basalt at the base of the lithosphere,
27 melting of lower crust, voluminous granite magmatism, and widespread and near-continuous,
28 mid-crustal ultra-high-temperature (UHT) metamorphism. Direct ascent of basic magmas into the
29 upper crust was inhibited by the ductile nature of the lower crust and the development of
30 substantial crystal-rich magma storage chambers. In the period between c. 1150 and 1090 Ma
31 magmatism ceased, possibly because the lower crust had become too refractory, but mid-crustal
32 reworking was continuously recorded in the crystallisation of zircon in anatectic melts. Renewed
33 magmatism in the form of the Giles Event of the Warakurna LIP began at around 1090 Ma and was
34 characterized by voluminous basic and felsic volcanic and intrusive rocks grouped into the
35 Warakurna Supersuite. Of particular interest in the context of the present study are the Giles
36 layered intrusions which were emplaced into localized extensional zones. Rifting, emplacement of
37 the layered intrusions, and significant uplift all occurred between 1078 and 1075 Ma, but mantle-
38 derived magmatism lasted for >50 m.y., with no time progressive geographical trend, suggesting
39 that magmatism was unrelated to a deep mantle plume, but instead controlled by plate
40 architecture.

41 The Giles layered intrusions and their immediate host rocks are considered to be prospective
42 for (i) platinum group element (PGE) reefs in the ultramafic–mafic transition zones of the
43 intrusions, and in magnetite layers of their upper portions, (ii) Cu–Ni sulfide deposits hosted within
44 magma feeder conduits of late basaltic pulses, (iii) vanadium in the lowermost magnetite layers of
45 the most fractionated intrusions, (iv) apatite in unexposed magnetite layers towards the evolved
46 top of some layered intrusions, (v) ilmenite as granular disseminated grains within the upper
47 portions of the intrusions, (vi) iron in tectonically thickened magnetite layers or magnetite pipes of

48 the upper portions of intrusions, (vii) gold and copper in the roof rocks and contact aureoles of the
49 large intrusions, and (viii) lateritic nickel in weathered portions of olivine-rich ultramafic intrusions.

50

51 **Keywords:** Musgrave Province, Giles event, layered intrusions, PGE deposits, magnetite layers

52

53 **1. Introduction**

54 The Musgrave Province of central Australia hosts one of the most important clusters of mafic–
55 ultramafic layered intrusions globally (Fig. 1), referred to as the Giles Complex (Daniels, 1974) or
56 the Giles intrusions (Smithies et al., 2009). Together with broadly contemporaneous bimodal
57 volcanism of the Bentley Supergroup and basic magmatism of the Warakurna Large Igneous
58 Province (LIP)(Wingate et al., 2004), the Giles intrusions constitute the Warakurna Supersuite,
59 formed during the c. 1090 to 1040 Ma Giles Event. The prospectivity of the Giles intrusions for
60 magmatic ore deposits remains poorly understood. This is partly due to sparse exposure and
61 because much of the study area belongs to the Ngaanyatjarra – Anangu Pitjantjatjara –
62 Yankunytjatjara Central Reserve into which access is strictly regulated. However, the enormous
63 volume of mafic igneous rocks and the remarkable size of some of the intrusions (up to several
64 1000 km²) reflect a high flux of mantle derived magma and heat into the crust. This is considered
65 to be favorable for the formation of magmatic and hydrothermal ore deposits. Two world-class
66 deposits have been discovered so far, namely the Nebo–Babel magmatic Ni–Cu deposit (Seat et
67 al., 2007, 2009) and the Wingellina Ni laterite deposit (Metals X Ltd, 2013). In the present paper
68 we review the ore potential of the Giles intrusions and related mafic intrusive rocks of the
69 Warakurna Supersuite.

70

71 **2. Past work**

72 Systematic geologic research on the Musgrave Province began with a mapping program (at 1:250
73 000 scale) in the 1960s (Geological Survey of Western Australia - reported in Daniels, 1974), during
74 which the Blackstone, Murray, and Morgan ranges, and parts of the Cavenagh and Jameson
75 ranges, were mapped (see Fig 1 for localities). Nesbitt and Talbot (1966) subsequently proposed
76 that some of the layered intrusions are tectonised remnants of an originally much larger body.
77 Other important early contributions on the layered intrusions include the papers by Goode and
78 Krieg (1967), Goode (1970, 1976a,b, 1977a,b,c, 1978), and Goode and Moore (1975). A large
79 multidisciplinary study of the Musgrave Province beginning in the late 1980s (Australian Geological
80 Survey Organisation, now Geoscience Australia: Glikson, 1995; Glikson et al., 1996) also focused
81 mainly on the Giles intrusions (Ballhaus and Glikson, 1989, 1995; Ballhaus and Berry, 1991; Clarke
82 et al., 1995a,b; Glikson, 1995; Sheraton and Sun, 1995; Stewart, 1995). Recent work includes
83 studies by Seat et al. (2007, 2009) and Godel et al. (2011), on the Nebo–Babel Ni–Cu ore deposit,
84 and by Evins et al. (2010a,b) and Aitken et al (2013), on the structural evolution during the Giles
85 Event.

86

87 **3. Regional geology**

88 In the present paper, the term ‘Musgrave Province’ is used to refer to high-grade metamorphic
89 rocks affected by the 1220-1150 Ma Musgrave Orogeny, covering an area up to 800 km long and
90 350 km wide in central Australia (Fig. 1). The Western Australian segment of the Musgrave
91 Province is referred to as the ‘west Musgrave Province’. On geophysical images the Province is
92 delineated by a series of east-trending anomalies. It is tectonically bound by the Neoproterozoic to
93 Paleozoic sedimentary rocks of the Amadeus Basin in the north and the Officer Basin in the south
94 (Edgoose et al., 2004).

95 The Musgrave Province lies at the junction between the North, South and West Australian
96 Cratons. While some models suggest these cratonic elements of the Australian Craton
97 amalgamated as early as c. 1700 Ma (e.g. Li, 2000; Wingate and Evans, 2003), most models agree
98 that final amalgamation pre-dates the Musgrave Orogeny at c. 1220 Ma (Giles et al., 2004; Betts
99 and Giles, 2006; Caood and Korsch, 2008; Smithies et al., 2010, 2011; Kirkland et al., 2013).

100 With the exception of c. 1575 Ma rocks in the Wannarn area, the basement to the west
101 Musgrave Province is not exposed. However, isotopic data on the detrital components in
102 paragneisses, and on zircon xenocrysts, indicate that the basement is dominated by two major
103 juvenile crust formation events, at 1600–1550 Ma and 1950–1900 Ma (Kirkland et al., 2013).

104 The oldest magmatic rocks in the study area comprise felsic calc-alkaline rocks of the
105 Papulankutja Supersuite at c. 1400 Ma (Howard et al., 2011b; Kirkland et al., 2013). However, the
106 oldest event recognizable throughout the west Musgrave Province is the Mount West Orogeny,
107 characterized by emplacement of calc-alkaline granites of the Wankanki Supersuite mainly within
108 the central and southeastern part of the west Musgrave Province (Evins et al., 2009; Smithies et
109 al., 2009). Crystallization ages cluster between c. 1326 and 1312 Ma (Gray, 1971; Sun et al., 1996;
110 White et al., 1999; Bodorkos et al., 2008a–e; Kirkland et al., 2008a–f; Smithies et al., 2009). The
111 rocks are typically metaluminous, calcic to calc-alkaline granodiorites and monzogranites,
112 compositionally resembling Phanerozoic granites of the Andean continental arc (Smithies et al.,
113 2010). The Mount West Orogeny may have been triggered by the amalgamation of the North,
114 West, and South Australian Cratons and associated subduction and accretion (Giles et al., 2004;
115 Betts and Giles, 2006; Smithies et al., 2010, 2011; Kirkland et al., 2013).

116 The 1220–1150 Ma Musgrave Orogeny formed in an intracratonic (Wade et al., 2008;
117 Smithies et al., 2009, 2010) or back-arc setting (Smithies et al., 2013). Deformation and
118 metamorphism was of high grade resulting in abundant mylonites. The main magmatic

119 components are charnockitic and rapakivi granites of the Pitjantjatjara Supersuite. The earliest
120 Pitjantjatjara granites are strongly Yb-depleted interpreted to have formed through deep-crustal
121 melting in the presence of garnet. A transition from these to Yb-undepleted granites derived from
122 shallower depth is diachronous, attributed to removal of the lower crust and mantle lithosphere,
123 previously thickened during the Mount West Orogeny (Smithies et al., 2010, 2011). Intrusion of
124 the high-T Pitjantjatjara granites (Smithies et al., 2010, 2011) coincided with a 70–100 m.y. period
125 of regional ultra-high-temperature (UHT) metamorphism (King, 2008; Kelsey et al., 2009, 2010;
126 Smithies et al., 2010, 2011), characterized by lower to mid-crustal temperatures of >1000°C, along
127 a geothermal gradient of $\geq 35\text{--}40^\circ\text{C}/\text{km}$ (King, 2008; Kelsey et al., 2009, 2010). Such conditions are
128 consistent with removal of the lithospheric mantle. Thermal modeling and zircon geochronology
129 indicates that mid-crustal temperatures remained elevated (>800°C) in the period between the
130 Musgrave Orogeny and the Giles Event (Smithies et al., 2015), mainly due to the accumulation of
131 the highly radiogenic Pitjantjatjara granites.

132 The c. 1090 to 1040 Ma Giles Event was characterized by voluminous mafic to felsic
133 magmatism (Fig. 2), including the layered mafic-ultramafic ‘Giles intrusions’ (G1), massive gabbro
134 (G2) locally mixed and mingled with granite, various dyke suites including the Alcurra Dolerite
135 suite, granite plutons, as well as mafic and felsic lavas, volcanoclastic and sedimentary rocks
136 forming the Bentley Supergroup. All these components are grouped into the Warakurna
137 Supersuite interpreted to have accumulated in the long-lived intracontinental Ngaanyatjarra Rift
138 (Evins et al., 2010b; Aitken et al., 2013). Based in part on the extensive outcrop of the Warakurna
139 Supersuite across ~ 1.5 million km^2 the Giles magmatism has been interpreted as the result of a
140 mantle plume (Wingate et al., 2004; Morris and Pirajno, 2005). However, the most conservative
141 estimates for the duration of mantle magmatism are >30 m.y. with a likelihood it continued for
142 significantly longer (Smithies et al., 2015, and in press), with no time-progressive geographical

143 trend or track, inconsistent with a simple plume model (Smithies et al., 2013, 2015, in press; Evins
144 et al., 2010a,b).

145 Younger events include the 580–530 Ma intracratonic Petermann Orogeny, during which
146 many of the Giles intrusions were fragmented. Additional younger events are reflected by several
147 regional dolerite dyke suites (at c. 1000, c. 825, and c. 750 Ma) and low-volume felsic magmatism
148 (at c. 995 Ma and c. 625 Ma)(Howard et al., 2015).

149

150 **4. Tectonic subdivision**

151 Past workers divided the Musgrave Province into a number of sub-zones that show distinct
152 structural and metamorphic characteristics. The sub-zones are separated by major west- and
153 west-northwesterly trending faults, including the south-dipping Woodroffe Thrust (Fig. 1). In the
154 eastern portion of the west Musgrave Province, there is a marked north-to-south change in the
155 pressure of granulite-facies metamorphism. In the northern segment, high-pressure (10–14 kbar)
156 metamorphism during the Petermann Orogeny has masked the effects of older metamorphism
157 (Scrimgeour and Close, 1999). To the south, the metamorphic overprint of the Petermann
158 Orogeny is not as marked and evidence for relatively low-pressure, but high-temperature
159 Mesoproterozoic metamorphism is preserved (Clarke et al., 1995b). The boundary between these
160 two regimes lies close to the west-trending, near vertical Mann Fault (Fig. 1).

161 The western part of the study area is subdivided into three distinct zones, namely the Walpa
162 Pulka, Tjuni Purlka, and Mamutjarra Zones (from northeast to southwest; Howard et al., 2014; Fig.
163 2). The Tjuni Purlka Zone represents a northwest-trending belt of multi-generational (c. 1220,
164 1075, and 550 Ma) shearing and mafic magmatism of the Warakurna Supersuite (Fig. 2) that
165 remained active throughout much of the Giles Event. The Walpa Pulka Zone (Fig. 2) is a deep-
166 crustal domain hosting abundant c. 1220–1150 Ma granites of the Pitjantjatjara Supersuite that

167 were emplaced during the Musgrave Orogeny. Mafic intrusions are rare and restricted to small
168 bodies north of the Hinckley Range. The Mamutjarra Zone in the south (Fig. 2) contains several
169 Giles intrusions and the c. 1345–1293 Ma calc-alkaline granites of the Wankanki Supersuite,
170 emplaced during the Mount West Orogeny.

171

172 **5. Analytical methods**

173 Samples were prepared for analysis at GSWA using a jaw crusher followed by milling in a tungsten
174 carbide mill. The mill was tested for possible contaminants, with only cobalt being significant
175 (≤ 157 ppm in grinding tests). Major elements were determined by wavelength-dispersive X-ray
176 fluorescence spectrometry (XRF) on fused disks using methods similar to those of Norrish and
177 Hutton (1969). Precision is better than 1% of the reported values. Concentrations of Ba, Cr, Cu, Ni,
178 Sc, V, Zn, and Zr were determined by wavelength-dispersive XRF on pressed pellets using methods
179 similar to those of Norrish and Chappell (1977), whereas Cs, Ga, Nb, Pb, Rb, Sr, Ta, Th, U, Y, and
180 REE were analysed by inductively coupled plasma–mass spectrometry (ICP-MS) using methods
181 similar to those of Eggins et al. (1997), on solutions obtained by dissolution of fused glass disks.
182 Precision for trace elements is better than 10% of the reported values. All whole-rock major and
183 trace element data for the silicate rocks are listed in Table 1 of the supplementary data. All
184 analyses and analytical details can be obtained from the WACHEM database (at
185 <http://geochem.dmp.wa.gov.au/geochem/>). Selected major and trace elements for the massive
186 magnetite seams were determined by instrumental neutron activation analysis (INAA) at The
187 University of Québec at Chicoutimi, Canada (Table 2, supplementary data).

188 Platinum, Pd, and Au were analysed by lead collection fire assay of 40 g of sample, followed
189 by ICP quantitation. The detection limit was 1 ppb for each element. For selected samples,
190 complete PGE spectra were obtained by ICP-MS at The University of Quebec at Chicoutimi.

191 Analytical details are given in Barnes et al. (2010). Platinum group element data are shown with
192 the bulk of the lithophile element data in Table 1, supplementary data.

193 Sulfur isotopes were analysed by Rafter GNS Sciences at the New Zealand National Isotope
194 Centre at Lower Hutt, New Zealand. Samples were measured in duplicate in tin capsules with
195 equal amounts of V₂O₅ on a EuroVector elemental analyzer connected to a GVI IsoPrime mass
196 spectrometer. All results are averages and standard deviations of duplicates are reported with
197 respect to Vienna Canyon Diablo Troilite (VCDT) standard, normalized to internal standards
198 R18742, R2268, and R2298 with accepted $\delta^{34}\text{S}$ values of -32‰ , $+3.3\text{‰}$, and $+8.6\text{‰}$, respectively.
199 The external precision for this instrument is better than 0.3 for $\delta^{34}\text{S}$. All data are listed in Table 3,
200 supplementary data.

201 In situ Sr isotope data were determined by laser ablation ICP-MS at the Geological Survey of
202 Finland (GTK). Analytical procedures are provided in Yang et al. (2013b), and the data are provided
203 in Table 4, supplementary data.

204 For most samples, Sm–Nd isotopic analyses were determined by isotope dilution at the
205 VIEPS Radiogenic Isotope Laboratory, Department of Earth Sciences, La Trobe University, Victoria.
206 Analytical techniques follow those of Waight et al. (2000). All quoted ϵNd values are initial values
207 calculated at the time of igneous crystallization. For some samples, Nd isotopes were determined
208 at GTK Espoo. Analytical details are given in Maier et al. (2013a). All Sm–Nd isotope data are listed
209 in Table 5, supplementary data.

210 The composition of olivine was determined at The University of Oulu in Oulu, Finland, using
211 a JEOL JXA-8200 electron microprobe at an accelerating voltage of 15 kV and a beam current of 30
212 nA, which allowed an approximately 150 ppm detection limit for nickel. The accuracy of analyses
213 was monitored using reference material of similar compositions. Compositional data are listed in
214 Table 1.

215

216 **6. Geology and petrology of the Giles intrusions**

217

218 **6.1. Introduction**

219 The Giles intrusions in western Australia show considerable variation in terms of size (from a few
220 km² to >3000 km²), depth of emplacement (from < 1kbar to possibly as much as 12 kbar), whole
221 rock and mineral composition, as well as style of associated mineral deposits. In the following
222 section, we group the intrusions by their predominant lithologies. Intrusions with important
223 ultramafic segments of wehrlite, harzburgite, websterite and (olivine) orthopyroxenite include
224 Wingellina Hills, Pirntirri Mulari, The Wart (Fig. 1). Intrusions that are predominantly
225 leucogabbronoritic are more common, comprising Hinckley Range, Michael Hills, Latitude Hill,
226 Murray Range, Morgan Range, Cavenagh Range and Saturn. The Blackstone, Jameson-Finlayson,
227 and Bell Rock ranges also belong to this group, but they are now believed to be tectonically
228 segmented portions of an originally single body, hereafter called the Mantamaru intrusion. In
229 addition, a number of smaller intrusions or fragments of intrusions occur to the north of the Tjuni
230 Purlka Zone including Mt Muir and Lehmann Hills.

231 Several of the mafic intrusions contain thick troctolitic successions, namely the Cavenagh
232 and Morgan ranges and Mantamaru. Anorthosites may form relatively thin layers (centimetres to
233 several tens of centimetres) in the mafic intrusions, but attaining thicknesses of several tens of
234 metres at Kalka in South Australia.

235

236 **6.2. Mafic-ultramafic layered intrusions**

237 **6.2.1. Pirntirri Mulari**

238 The intrusion is located ~30 km north of Blackstone Community (Fig. 2). It was grouped with the
239 gabbronoritic Murray Range by Ballhaus and Glickson (1995), but in view of the distinct lithologies
240 and composition of the intrusions (ultramafic-mafic for Pirntirri Mulari, mafic for Murray) this
241 interpretation is rejected here. The exposed portion of the intrusion has the shape of a wedge,
242 with the upper mafic portion being markedly wider than the lower ultramafic portion (Fig. 3). The
243 layers strike about 150° and mostly dip steeply to the southwest (60–90°). The body is about 5 km
244 wide and has a stratigraphic thickness of about 3 km.

245 Most of the rocks are medium-grained (olivine) websterites, peridotites, (olivine)
246 orthopyroxenites, and (olivine) gabbronorites. At the southwestern and northeastern margins of
247 the intrusion occur interlayered websteritic and gabbronoritic orthocumulates, whereas
248 adcumulate peridotites and orthopyroxenites are concentrated in the centre. Based on
249 compositional and field evidence, we propose that the succession youngs to the northeast,
250 implying that the intrusion is slightly overturned. Many of the rocks show textural equilibration,
251 expressed by 120° grain boundaries, abundant bronzite and spinel exsolutions in clinopyroxene,
252 small granoblastic plagioclase grains containing spinel inclusions, orthoclase exsolution blebs and
253 two-pyroxene spinel symplectite at contacts between plagioclase and olivine. These features are
254 common to many other Giles intrusions and were previously interpreted to reflect relatively high
255 crystallization pressures and temperatures (Ballhaus and Glikson, 1989).

256 The basal contact of the intrusion is not exposed, but in view of the increased proportion of
257 orthocumulates towards the southwestern exposed edge, we argue that the basal contact is
258 proximal to this. The top contact is likely of a tectonic nature, as suggested by the presence of a
259 mylonite zone and by the relatively unevolved chemical composition of the uppermost gabbros
260 (Mg# 0.7, Cr/V > 1) ([Table 1 data repository](#)).

261 Most exposed contacts between layers are sharp (Fig. 4a). In the centre of the intrusion
262 (GSWA189374), coarse-grained websterite underlies gabbronorite with an undulose contact, and
263 has locally injected the gabbronorite (Fig. 4b). The gabbronorite is more deformed than the
264 pyroxenite, and is altered near the contact, consistent with intrusion of pyroxenite below older
265 gabbronorite. Elsewhere (e.g., GSWA189368), fine-grained lherzolite overlies coarse-grained
266 peridotite with an undulose, sharp contact. The lherzolite contains inclusions of the peridotite and
267 is interpreted to have crystallized from a new magma influx that was quenched and partially
268 eroded its floor. Other ultramafic layers are underlain by pegmatoidal gabbronoritic
269 orthocumulate layers (Fig. 4c), consisting of 40% plagioclase, 30% orthopyroxene, 20%
270 clinopyroxene, and accessory phlogopite and chromite. Relatively high concentrations of
271 incompatible trace elements (e.g. P, LREE, Nb, Rb) in the pegmatoid suggest that it may have
272 formed in response to upward percolation of evolved melt or fluid.

273 The lower portion of the intrusion, particularly below and above pyroxenite layers, contains
274 several horizons showing textural evidence for considerable syn-magmatic cumulate mobility,
275 such as abundant ultramafic and anorthositic schlieren within gabbronorite (Fig. 4d).

276 The concentrations of the platinum group elements (PGE) in the intrusion are mostly <10
277 ppb (Fig. 5), and Cu/Pd is at the level of the primitive mantle (4000–7000), suggesting that the
278 magma was fertile and sulfur undersaturated during crystallization. However, at a stratigraphic
279 height of ~ 2600 m above the base of the intrusion, Cu/Pd increases sharply, suggesting that a
280 sulfide melt had fractionated from the magma at this stage. This sample has low PGE
281 concentrations, although it contains nearly 500 ppm Cu. It is located in the mafic–ultramafic
282 transition interval, analogous to the stratigraphic position of PGE reefs in the Bushveld Complex
283 and many other PGE-mineralized intrusions. A sulfide-bearing pyroxenite from an equivalent
284 stratigraphic level collected along the southeastern edge of the intrusion (Fig. 3) has similar

285 chromium, copper, and nickel concentrations to the equivalent internal horizon, but higher Pt+Pd
286 (172 ppb) and gold (17 ppb) concentrations. Assay results using a NITON portable XRF showed
287 elevated Ni and Cu over a stratigraphic interval of 5–10 m, and peak concentrations of 0.43% Cu
288 and 0.7% Ni in weathered rock (Redstone Resources Ltd, 2008a). No detailed sampling of this
289 interval has yet been conducted to establish the width and grade of a putative reef horizon.

290

291 **6.2.2. Wingellina Hills**

292 The intrusion is approximately 12 km long and up to 3 km wide (Fig. 6). Drilling by Acclaim
293 Minerals and Metals X Ltd, and mapping by GSWA, established dips of about 65–75° to the
294 southwest, and younging in the same direction indicating that the intrusion is not overturned. The
295 strike of the layering is 110–120°. The exposed stratigraphic thickness amounts to 2.5 km,
296 assuming that the stratigraphy is not duplicated by tectonism. The gabbros and pyroxenites tend
297 to be unaltered at the surface, whereas the peridotites are commonly deeply weathered to a
298 depth of about 60 m to >200 m, particularly in shear zones.

299 The central portion of the intrusion contains numerous cyclic units consisting of basal
300 pegmatoidal (ortho) pyroxenite, overlain by clinopyroxenite and then peridotite (olivine–spinel
301 cumulate), wehrlite, and gabbronorite (Ballhaus and Glickson, 1989). The gabbronorite may
302 contain fragments and schlieren of ultramafic material, and it may display convoluted and folded
303 layering on a scale of 1–2-centimetres, analogous to Pirntirri Mulari (Fig. 4).

304 The composition of the basal to central portion of the intrusion has been studied in several
305 stratigraphic boreholes. The basal contact has been intersected by a reverse circulation (RC)
306 borehole (WPRC23, see Maier et al., 2014 for details). It is interpreted as intrusive because there is
307 a well-defined, > 100 m wide basal compositional reversal that is characteristic of layered
308 intrusions (Latypov et al., 2011). Overlying the contact is a 1–2m thick zone consisting of hybrid,

309 possibly contaminated gabbroic rocks (~10% MgO). A rock chip collected ~2 m above the contact
310 consists of medium-grained, moderately deformed olivine gabbronorite, implying relatively
311 insignificant tectonism of the contact zone (R Coles, 2013, written comm., 27 May).

312 In the next 300 m of the drill core, basal olivine gabbronorite is first overlain by pyroxenite
313 and then by progressively more magnesian harzburgite. This, in turn, is followed by peridotite and
314 wehrlite and then by about 20 m of websterite and >40 m of olivine gabbronorite. The contacts
315 between rock types can be either sharp or gradational. Within the websterite occurs a PGE reef
316 (described below) that has been identified along a strike length of 2–3 km (Fig. 6). The remainder
317 of the intrusion consists of layers of peridotite, wehrlite, pyroxenite, and olivine gabbronorite,
318 described in more detail in Ballhaus and Glickson (1989).

319 The compositional variation in the interval hosting the PGE reef is shown for borehole
320 WPRCO-064 (Fig. 7). The contact between websterite and wehrlite is sharp, reflected by a marked
321 decrease in MgO and an increase in Cr concentration. Platinum-group element concentrations
322 increase through the websterite layer to a maximum of 2ppm Pt+Pd+Au, at a level about 5–7 m
323 beneath the top of the layer. The peak grades occur over a 1-m interval, and grades in excess of 1
324 ppm PGE occur over 3–5 m. Above the reef, the concentrations of the PGE decrease relatively
325 rapidly over a height of a few meters. Gold and Cu concentrations remain relatively low (<5 ppb
326 Au, <10 ppm Cu) throughout the PGE-enriched zone, but peak just above the PGE reef, with up to
327 330 ppb Au in a 1-m interval, and up to 400 ppm Cu in rocks located a further 2 m above the Au
328 peak. Gold and Cu concentrations then decrease with further stratigraphic height, although at a
329 slower rate than the PGE levels. Similar metal distribution patterns have been observed in several
330 other layered intrusions (e.g., the c. 2.58 Ga Great Dyke, Wilson et al., 1989; the c. 2.925 Ga Munni
331 Munni intrusion, Barnes, 1993; and the c. 3.03 Ga Stella intrusion, Maier et al., 2003b), where they
332 were referred to as 'offset patterns' (Barnes, 1993). In comparison to the Bushveld PGE reefs (~1–

333 2% S), the PGE reefs at Wingellina Hills are relatively sulfur-poor (mostly <500 ppm S), possibly due
334 to metamorphic devolatilization. This renders the reefs nearly invisible in hand specimen.

335 The thickness and grade of the PGE reef shows considerable variation along-strike. In two of
336 the three analysed boreholes (WPRC0-064 and WPRCD0-083) concentrations of >500 ppb PGE
337 occur over a thickness of 8 m. The bulk concentrations of PGE+Au normalized to a width of 1 m
338 are similar in the two holes (10.6 ppm in WPRC0-064, 9.6 ppm in WPRCD0-083). Borehole WPRC0-
339 043 contains >100 ppb PGE+Au over 12 m, and total PGE+Au contents of 5.5 ppm normalized to 1
340 m, lower by about 40% than in the two other holes. This is unlikely to be the result of alteration,
341 but implies significant variation in PGE grade of the reef along strike. Notably, the bulk PGE
342 contents normalized to a width of 1m of the reef interval (at least in holes WPRCD0-064 and
343 WPRCD0-083) are broadly similar to the PGE contents of the combined Merensky Reef and UG2
344 chromitite of the Bushveld Complex. However, the Bushveld reefs are much narrower and thus
345 more economic to mine.

346 The Wingellina Hills laterite consists of yellow-brown to dark brown ochre material
347 composed of goethite, manganese oxides, gibbsite, and kaolinite produced by weathering of
348 peridotite. This constitutes the Wingellina nickel laterite deposit, discovered by INCO in 1956 (187
349 Mt at about 1% Ni and 0.08% Co; >167 Mt is classified as probable mining reserve; Metals X Ltd,
350 2013). Limonitic ochre is also present at Claude Hills (4.5 Mt at 1.5% Ni; Goode, 2002), in the
351 Pirntirri Mulari intrusion, and in the southeastern part of the Bell Rock intrusion where
352 excavations revealed laterite, ochre, and chalcedonic veins above a zone of saprolite (Howard et
353 al., 2011b). The laterites formed by selective leaching of SiO₂ and MgO, resulting in residual
354 concentration of alumina, iron oxides, and nickel, developed particularly prominently along shear
355 zones. The ore is exposed at surface and has an average thickness of 80 m (maximum 200 m). The
356 deposit has a high aspect- and very low strip ratio. It is locally cut by semiprecious, pale green

357 chrysoprase mined artisinally since the 1960s, particularly in the Kalka area (Goode, 2002). The
358 lateritic profiles may have some potential for scandium deposits, particularly where the parent
359 rock consisted of clinopyroxenite, but to our knowledge, no relevant investigations have so far
360 been conducted.

361

362 **6.2.3. The Wart**

363 The Wart is a relatively small mafic-ultramafic body, located about 20 km south of the community
364 of Wingellina. The main block of the intrusion measures about 5 km x 2 km striking about 130° and
365 dipping steeply at 80–90° to the southwest. Several smaller mafic slivers occur to the northwest of
366 the main body. The stratigraphic thickness of the body is ~1–2 km. Ballhaus and Glikson (1995),
367 who studied the Wart in some detail suggested that it represents the lower portion of the Bell
368 Rock intrusion.

369 The Wart shares certain lithological and compositional characteristics with Pirntirri Mulari.
370 Both contain layers of medium-grained mesocumulate wehrlite–peridotite within a package of
371 adcumulates and mesocumulates of clinopyroxenite and melagabbronite. Many of the
372 ultramafic layers have sharp contacts and have been interpreted as sills (Ballhaus and Glikson,
373 1995). We collected only a small number of gabbroic and ultramafic samples because the body is
374 culturally sensitive. However, abundant mineral compositional data are given in Ballhaus and
375 Glikson (1995).

376

377 **6.3. Predominantly mafic intrusions**

378 **6.3.1. Latitude Hill – Michael Hills**

379 Latitude Hill is located 5–10 km to the east of The Wart and the Bell Rock intrusions (Fig. 2).

380 Ballhaus and Glikson (1995) proposed that it may be a folded segment of the 8000m thick Michael

381 Hills gabbro, whereas Pascoe (2012) suggested that Latitude Hill is in faulted contact with Bell
382 Rock. Latitude Hill has an intrusive contact with granulite analogous to Wingellina Hills and The
383 Wart (Ballhaus and Glikson, 1995; present work). It contains numerous layers and lenses of olivine
384 gabbronorite and olivine pyroxenite, as well as rare peridotite, but the dip direction of the layers
385 remains uncertain.

386

387 **6.3.2. Morgan Range**

388 The Morgan Range is located ~ 10 km north of Blackstone Community (Fig. 2). It measures
389 approximately 10 km x 5 km (~50 km²) in size, striking broadly 120° and with a stratigraphic
390 thickness of >1 km. The intrusion consists mostly of relatively unaltered olivine gabbronorites and
391 troctolites that display modal layering on a scale of centimetres to metres. It forms a boat-shaped
392 structure with steep dips (up to 80°) to the interior, except for the centre where the dip is sub-
393 horizontal. The syncline plunges at a relatively shallow angle towards the southeast and
394 compositional data suggest that the intrusion is not overturned.

395 At the northeastern tip of the intrusion is a relatively small (~300 m x 300 m) interlayered
396 mafic-ultramafic block consisting of dunite, troctolite and melagabbronorite. The rocks strike
397 ~100° and dip steeply (80°) to the north. They show certain compositional similarities to
398 Wingellina Hills and Pirntirri Mulari, but because the contact with the main Morgan Range is
399 concealed by regolith it is presently unclear whether this segment forms an integral part of the
400 Morgan Range intrusion, or whether it is a fragment of another intrusion that was tectonically
401 adjoined to the Morgan Range.

402

403 **6.3.4. Hinckley Range**

404 This is a large (~30 km x 10 km), highly deformed, relatively poorly layered body that has a
405 stratigraphic thickness of about 5800 m, strikes about 100°, and dips steeply to the north at 70–
406 80°. The rocks consist of (olivine) gabbros, troctolites, and microgabbros, with a few layers or
407 lenses of anorthosite, and minor pyroxenite. Abundant pseudotachylite are related to the
408 Petermann Orogeny. Many of our samples have relatively high concentrations of K₂O and
409 incompatible trace elements, suggesting that the mafic magma assimilated, or mixed with, a
410 granitic component. Our most primitive samples were collected at the southern edge of the
411 intrusion, suggesting that the intrusion youngs towards the northeast (cf Ballhaus and Glickson,
412 1995).

413 Where the Hinckley Range intrusion is in contact with the West Hinckley Range intrusion
414 (MGA 472843E 7118953N), mingling textures between gabbro and granite are prominent. These
415 have been described in Howard et al. (2011b) and Maier et al. (2014), from whom the following
416 description is taken. The gabbro belongs to the unlayered G2 variety which tends to crosscut,
417 engulf, and post-date the layered G1 intrusions. The G2 gabbros locally form agmatites or injection
418 migmatites (Fig. 8), and angular blocks engulfed by granite. Brittle fractures in gabbro may be
419 infilled by granite veins. This indicates that the granite has intruded partially solidified gabbro.
420 Contacts between the felsic and mafic portions may also be cusped, indicating that the mafic
421 component behaved in a ductile manner in the presence of the felsic material. Similar
422 relationships occur some 20 km to the northwest, at Amy Giles Hill (Fig. 2). A leucogranite showing
423 well-developed co-mingling textures with gabbro (Howard et al., 2006a) has been dated at 1074 ±
424 3 Ma (GSWA 174589; Bodorkos and Wingate, 2008b).

425 In the West Hinckley Range, mingled gabbro forms a kilometre-scale fold with a steep
426 northwest-trending axial plane that has been intruded by syndeformational leucogranite. A strong
427 'gneissic' fabric has locally formed in mixed or agmatitic rocks as the axial planar fabric continued

428 to develop, and this has been again engulfed within subsequent injections of leucogranite. A
429 sample of undeformed leucogranite from within one of these axial planar zones, approximately 2
430 km south of the mingled gabbro–granite described above, yielded a crystallization date of $1075 \pm$
431 7 Ma (GSWA 174761; Kirkland et al., 2008e). Syn-mylonitic leucogranite has also pooled into
432 boudin necks in a northwest-trending mylonite directly south of Charnockite Flats (~2.5 km
433 northwest of the West Hinckley Range intrusion), and has been dated at 1075 ± 2 Ma (GSWA
434 185509; Kirkland et al., 2008f). The combined data define a very narrow period of intrusion of
435 massive G2 gabbro and multi-phase intrusion of leucogranites (1078–1074 Ma), northwest-
436 directed folding, and northwest-trending shearing. The relationships between gabbro and granite
437 in the Hinckley Range intrusion confirm earlier suggestions by Clarke et al. (1995b) that substantial
438 deformation occurred during the Giles Event.

439

440 **6.3.5. Murray Range**

441 The gabbroic Murray Range comprises two distinct segments, (i) a layered portion of $>25 \text{ km}^2$
442 consisting of gabbronorite or olivine gabbronorite, with the most primitive rocks occurring in the
443 centre of the intrusion, and (ii) extensive areas covered by massive gabbro, particularly to the east
444 and northeast of Pirntirri Mulari. The strike of the layering is mostly $50\text{--}70^\circ$, and the dip is sub-
445 vertical. The intrusion contains abundant stratiform layers and lenses of microgabbro and cross-
446 bedded medium-grained gabbronorites. Due to its location at the contact between the Tjuni
447 Purlka and Walpa Pulka Zones the intrusion was tectonically dismembered, obscuring the true
448 stratigraphic thickness and structure of the intrusion. Like the Hinckley Range, the Murray Range is
449 one of the G1 intrusions that was substantially intruded by G2 gabbro, consistent with the model
450 that during the emplacement of the G2 gabbros, the contact between the Tjuni Purlka and Walpa

451 Pulka Zones was a syn-magmatic shear zone (Evins et al., 2010a,b). It is thus not surprising that
452 deformation and alteration are commonly more pronounced than in the other Giles intrusions.

453

454 **6.3.6. Cavenagh Range**

455 The Cavenagh Range remains one of the least known amongst the Giles intrusions because access
456 to it is restricted on cultural grounds. The intrusion is located 10 km south of the Blackstone Range
457 (Fig. 2) and is defined by several circular remnant magnetic highs occurring over an area of
458 approximately 22 km x 18 km. The southern portion forms a syncline with a stratigraphic thickness
459 of about 1 km. It consists predominantly of olivine gabbronorites, olivine gabbros, troctolites, and
460 norites. Websterites, anorthosites and microgabbros form bands and discontinuous pods,
461 schlieren, and autoliths. Xenoliths of basement gneiss have been encountered near the
462 southeastern edge. An east-trending fault separates the southern segment from the northern
463 portion of the intrusion which dips at about 15–30° to the northeast and has a stratigraphic
464 thickness of about 2–4 km (barring structural duplication). It consists dominantly of poorly layered
465 olivine gabbronorite, troctolite, and magnetite-bearing olivine gabbronorite.

466 Layering in the Cavenagh intrusion is defined predominantly by modal variation between
467 olivine, pyroxene, and plagioclase. Boundaries between layers are mostly gradational, but
468 pyroxenites, anorthosites, and many microgabbros tend to have sharp contacts. The latter rocks
469 also tend to form lenses, schlieren, and fragments within gabbronorite and troctolite.

470 The southern to central portion of the Cavenagh intrusion is the least chemically evolved.
471 The northern portion shows a subtle trend toward more differentiated compositions with height.
472 Simultaneously, the concentrations of PGE increase, reaching 80–100 ppb PGE in two samples.
473 The microgabbros are interlayered with medium-grained gabbronorite and may contain autoliths
474 of anorthosite and thin bands, irregular clasts, and circular concretions of granular websterite and

475 clinopyroxenite adcumulate. These field relationships suggest that the microgabbros and the
476 associated medium-grained rocks intruded contemporaneously. The microgabbros tend to have
477 equigranular textures with 120° grain boundaries. In places, olivine and pyroxene grains may form
478 strings oriented in a radial configuration that are here interpreted to have resulted from crystal
479 growth in a flowing, supercooled magma. Large olivine oikocrysts can form wispy crystals that are
480 surrounded by rims of anorthosite. The latter may have formed when rapid crystallization due to
481 supercooling or degassing led to depletion in the olivine component within a boundary layer
482 surrounding the growing crystals. The microgabbros may also contain clinopyroxene oikocrysts
483 with abundant inclusions of irregular and rounded exsolved oxide grains.

484 The microgabbros have variable compositions, broadly overlapping with the medium grained
485 host rocks. Many of the samples contain a cumulus component, as indicated by whole-rock
486 chromium concentrations of up to 1900 ppm, positive strontium anomalies on multi-element
487 variation diagrams, and depleted incompatible trace element levels (e.g., <5 ppm Zr).
488 Microgabbros in the Giles intrusions and elsewhere were previously explained by intraplutonic
489 quenching (Ballhaus and Glikson, 1989, Tegner et al., 1993).

490

491 **6.3.7. Lehmann Hills, Mt Muir, and other small intrusions north of the Blackstone and Wingellina**

492 **Communities**

493 The intrusions outcrop over relatively small areas and consist of gabbronorites, olivine
494 gabbronorites and troctolites (Fig. 9a). The rocks may have a distinct flow structure, containing
495 elongated lenses and schlieren of anorthosite in pyroxenite, and autoliths of pyroxenite within
496 anorthosite. At Lehmann Hills, sulfides (up to 1% combined pyrrhotite and chalcopyrite) are
497 relatively abundant.

498 A number of small mafic bodies occur up to 20 km to the north of the Hinckley Range, in the
499 Mt Gosse – Mt Daisy Bates area (termed 'Northeast' in Table 1, data repository). These bodies
500 consist of metagabbros and metagabbro-norites. They tend to show partial granoblastic textures,
501 with garnet forming fine-grained rims around pyroxene and magnetite, or more rarely
502 porphyroblasts. Pyroxene is commonly replaced by hornblende, and biotite is also common.

503

504 **6.3.8. Mantamaru**

505 Mantamaru is the Ngaanyatjarra name for the community of Jameson. The intrusion forms one of
506 the world's largest layered igneous bodies, with an original size of at least 3400 km². This body was
507 dismembered during the Petermann orogeny resulting in the Jameson-Finlayson, Blackstone and
508 Bell Rock ranges.

509

510 Jameson–Finlayson Range

511 The Jameson–Finlayson Range extends for 66 km along a strike of ~120° and is ~ 30 km wide (Fig.
512 2). Layering is in normal orientation and dips at about 20° to 30° to the southwest, implying a
513 stratigraphic thickness of up to about 10 km. Several layer-parallel mylonitic zones occurring near
514 the base of the intrusion could account for limited structural repetition, but we argue that this
515 does not significantly affect the overall thickness estimates.

516 The bottom and top contacts of the intrusion are not exposed. At the base is magnetite–
517 ilmenite-bearing lherzolite; the rocks contain 20–50 vol.% opaques that have estimated V₂O₅
518 contents of about 1.4 wt% (Daniels, 1974). This is overlain by rhythmically layered troctolite and
519 olivine gabbro-norite (Fig. 9b). At the top of the intrusion, in the southwest, there is a layered
520 succession of troctolite, olivine gabbro, and olivine gabbro-norite, containing at least 11 major

521 titaniferous magnetite seams (Fig. 9c). However, due to poor outcrop, the thickness and contact
522 relationships of most of the seams remain poorly known.

523 The magnetite seams mostly form bands of rubble. They appear to be separated by silicate
524 intervals several hundreds of metres thick (Fig. 10). All layers that have been sampled consist of
525 massive iron oxide, with <5% silicates. Grain sizes are relatively coarse (≤ 3 mm), possibly reflecting
526 sintering (Reynolds, 1985). The main minerals are magnetite, granular ilmenite, fine ilmenite
527 lamellae, abundant hematite replacement patches and lamellae (≤ 20 vol.%), as well as goethite.
528 Layers enriched in apatite were not encountered.

529 The most reliable observations have been made on the basal magnetite layer. It has been
530 traced along a strike of about 19 km forming an aeromagnetic anomaly with sporadic broken
531 outcrop (Fig. 10). The seam may reach a thickness of 50 m, with up to three sub-seams locally
532 developed. Whether the sub-seams formed due to primary magmatic processes or structural
533 duplication is uncertain. The contacts of the seam with the magnetite-bearing leucotroctolite and
534 anorthosite hanging and footwall are sharp. The host rocks show evidence for deformation. The
535 mineralogy of the seam consists of magnetite and granular ilmenite, as well as fine ilmenite
536 lamellae within magnetite. Hematite and goethite replacement is locally abundant.

537 Concentrations of Pt, Pd, and Au have been analysed in 39 samples of the layer along strike
538 and in 3 traverses across the layer (Traka Resources, unpublished report). In addition, we analysed
539 three samples of the layer for the complete PGE spectrum (Table 1, data repository). The PGE
540 concentrations reach approximately 2 ppm Pt+Pd+Au (Fig. 11). The seam is thus markedly PGE
541 enriched relative to the Bushveld Main Magnetite Layer (Fig. 12), but has significantly lower PGE
542 contents than the main PGE mineralized magnetite layer of the Stella intrusion of South Africa
543 (Maier et al., 2003b). The metal concentration patterns of the Jameson seam show depletion in
544 Os–Ir–Ru relative to Rh–Pt–Pd, characteristic of evolved magmatic rocks (Fig. 12). The layer has

545 relatively constant vanadium concentrations throughout (up to 7400 ppm V, 1.35 wt% V₂O₅; Fig.
546 11), whereas the PGE tend to be markedly elevated at the base. Sulfur concentrations are mostly
547 100–150 ppm, locally reaching 700 ppm. Sulfur concentrations do not correlate with Cu and PGE
548 (Fig. 11), suggesting some of the sulfur could have been lost in response to equilibration of sulfide
549 with magnetite (Naldrett and Lehmann, 1988), or in response to low-grade metamorphic
550 devolatilisation. The average Pd/Ir ratio is relatively low (34), consistent with a magmatic origin of
551 the PGE mineralization. Platinum/Pd ratios are above unity, analogous to other PGE mineralized
552 magnetites and magnetite gabbros (see Maier, 2005 and references therein).

553 Magnetite layers 2, 3, 6 and 7 are not exposed, but their presence is suggested by prominent
554 magnetic anomalies (Fig. 10). Layers 4 and 5 are pervasively altered to goethite and hematite,
555 although they also contain abundant granular ilmenite. Layer 8 may be a plug-like body. Layer 11
556 is partly exposed and forms a massive, well-layered, possibly rotated layer dipping about 30° to
557 the southwest. The compositional variation in the upper seams is poorly understood, although it is
558 evident that the vanadium and noble metal contents are much lower than in layer 1, whereas iron,
559 chromium, and phosphorus concentrations increase with height (Table 2, supplementary data).

560

561 Blackstone Range

562 The body is ~50 km long and up to 5km wide. It strikes about 90° and layering dips at between 70
563 and 80° to the south and is not overturned. The exposed stratigraphic thickness is about 4 km. The
564 body is interpreted to be the exposed northern limb of an upright west-trending structural
565 syncline (the Blackstone Syncline). Relics of its north-dipping southern limb are sporadically
566 exposed 20 km to the south, directly north of the Cavenagh intrusion. The size of the combined
567 body is about 1400 km². It is conformably overlain by felsic volcanic rocks of the Tollu Group
568 (Bentley Supergroup).

569 The rocks are relatively unaltered and undeformed. Layering can be pronounced where
570 defined by thin (centimetre-scale) magnetite layers. Most of the rocks are (olivine) gabbronorites
571 and troctolites, each constituting approximately 50% of the total mass of the intrusion. Troctolitic
572 rocks occupy the central and southern portions of the intrusion. The troctolites contain less than
573 20% olivine, with the exception of two relatively olivine-rich layers (40% olivine, 100–150 m
574 thickness) that can be traced along much of the intrusion, one in the centre (sample GSWA
575 155669) and another near the southern edge. These two layers are also present in the Bell Rock
576 intrusion where they contain up to 80% olivine. The occurrence of these layers in both intrusions is
577 consistent with an interpretation whereby the two intrusions are fragments of a large, tectonically
578 dismembered proto intrusion (Nesbit and Talbot, 1966; Glikson, 1995). At the southern margin of
579 the intrusion occurs a ~ 1 m-thick magnetite layer that contains 1.5% V₂O₅. Relatively elevated Cu
580 concentrations of up to 250 ppm suggest the presence of minor sulfides, common to all
581 magnetite-rich rocks in the upper portions of the Blackstone intrusion. Whether this layer can be
582 correlated to magnetite layer 1 in the Jameson–Finlayson intrusion is uncertain, as it is highly PGE
583 depleted whereas the Jameson basal layer is relatively PGE rich.

584

585 Bell Rock Range

586 This body extends for ~ 50 km along a strike of ~ 120°. The exposed width is ~ 5–6 km, and the
587 rocks dip at 70° to the southwest. Field exposures of graded and cross-bedded layers, as well as
588 whole rock and mineral compositional data indicate younging of the intrusion to the southwest. A
589 detailed compositional study of the Bell Rock intrusion was conducted by Ballhaus and Glikson
590 (1995), from which some of the following information is taken. Our own sample base is relatively
591 small, comprising 14 samples taken across the body.

592 The exposed stratigraphic thickness is about 3800 m, but since the contacts of the body are
593 not exposed, this is a minimum estimate. The top of the intrusion is likely in contact with volcanic
594 rocks of the Bentley Supergroup; This implies that either the intrusion has been deeply eroded
595 after its emplacement or the top contact is faulted. The basal rocks consist of medium- to coarse-
596 grained troctolites and gabbros, whereas the central portion consists of magnetite-bearing
597 troctolite and the upper portion contains centimetre- to tens of centimetre-thick magnetite
598 seams, dunitic layers, numerous microgabbro sills, and a few anorthosite layers. Modal cyclicity
599 occurs on a centimetre to metre scale. A recent drillhole collared at the western edge of the
600 Latitude Hill intrusion with a dip of 70° toward the southwest (MDDH0001, drilled by Anglo
601 American Exploration (Australia) Pty Ltd as part of the Department of Mines and Petroleum's
602 Exploration Incentive Scheme), has intersected deformed magnetite-enriched troctolites that are
603 interpreted to belong to the Bell Rock intrusion (Pascoe, 2012). This suggests that magnetite
604 seams could be present below cover.

605

606 **6.3.9. Alcurra Dolerite suite**

607 The components of the Alcurra Dolerite suite comprise dolerite dykes and sills that constitute the
608 bulk of the 1078-1073 Ma Warakurna Large Igneous Province (Wingate et al., 2004), small basic
609 and intermediate bodies and dykes emplaced near the margins of older G1 layered mafic
610 intrusions, G2 massive gabbro, and comingled gabbro–granite.

611 Contact relationships constrain the emplacement dates of the Alcurra suite to <1078 Ma,
612 and dating of some of the intrusions indicates that magmatism continued to at least c. 1067 Ma
613 (Howard et al., 2009). However, geochemical data suggest that Alcurra-type rocks were likely
614 formed over a much longer period, including lavas of the Bentley Supergroup until at least 1047

615 Ma (Howard et al., 2009, 2011a; Smithies et al., 2013). The Alcurra Dolerite suite thus reflects
616 relatively long-lived mantle melting (Smithies et al., 2013).

617 The c. 1076 Ma mafic to intermediate rocks forming part of the Alcurra Dolerite suite
618 typically consist of fine- to medium-grained olivine gabbro, olivine norite, ferronorite, and
619 ferrodiorite. The latter rocks have evolved and Fe-rich tholeiitic compositions, resulting in a
620 pronounced aeromagnetic signature and high specific gravity.

621 Rocks of the Alcurra Dolerite Suite occur throughout the West Musgrave Province, including
622 the Blackstone Syncline, within the marginal zones of the G1 Jameson Range and Murray Range
623 where they intruded along the layer contacts and between the intrusions and their country rocks.
624 The northeast-trending, coarse-grained ferrogabbro dykes that crosscut the G1 Jameson intrusion,
625 and which also occur throughout the northern parts of the COOPER map sheet, used fractures and
626 faults related to the earlier Musgrave Orogeny.

627

628 **6.3.10. Saturn**

629 The Saturn intrusion defines an elliptical aeromagnetic anomaly with a diameter of approximately
630 10 km, located between the Cavenagh and Blackstone ranges (Figs. 2 and 13). Cross cutting field
631 relationships indicate that Saturn is younger than the Blackstone Range. However, a date of 1072
632 \pm 8 Ma determined by using the U–Pb method on baddeleyite in olivine gabbro (Redstone
633 Resources Ltd, 2007, written comm.) is within error of the c. 1078 to 1075 Ma range for both the
634 G1 and G2 mafic phases of the Giles Event. The concentric magnetic pattern implies zones of
635 magnetite enrichment, but massive magnetite layers have not been found on the surface.
636 Possibly, this is due to the very poor outcrop. The only exposed rocks consist of scattered massive,
637 medium-grained, leucocratic olivine gabbros, typically containing elevated concentrations of

638 biotite and up to 5% magnetite. The rocks are massive or show flow-banded textures, defined by
639 schlieren of fine-grained gabbro-norite (for example, at the 'Camp' site, Fig. 13).

640 The rocks have relatively evolved compositions, overlapping with Blackstone and Bell Rock.
641 Samples collected along the Phoebe traverse (Fig. 13, 14) contain up to 6.7% TiO₂ and 800 ppm V,
642 comparable to magnetite gabbros from the Jameson and Blackstone ranges. The rocks in the
643 centre of the Saturn intrusion are somewhat more primitive than those at the margin, having
644 higher Mg# and lower Ti concentrations (Fig. 14), consistent with a dome-like structure. Sulfide
645 contents are around 1%, higher than in most other Giles intrusions. Copper and PGE
646 concentrations are mostly relatively low, but they increase approximately halfway up the
647 magmatic stratigraphy (Fig. 14). Based on its age, the crosscutting relationships with rocks in the
648 Blackstone syncline, and compositional characteristics such as the enrichment in mica and sulfide,
649 the Saturn intrusion may be of transitional composition between the Alcurra Dolerite Suite and
650 the G1/G2 intrusive phase.

651

652 **6.3.11. Intrusions in the Halleys – Helena – DB Hill area**

653 The mafic rocks to the northeast and south of the Saturn intrusion were explored by Redstone
654 Resources at the Halleys, Halleys NW, Helena, and DB Hill prospects (Report at General Meeting,
655 2008b; Fig. 13). The intrusion(s) lack the strong remanent magnetic signature of the Cavenagh
656 intrusion and are thus interpreted to be distinct bodies. The crosscutting magnetic patterns
657 suggest that they intruded into the G1 intrusions as well as the volcanic, volcanoclastic, and clastic
658 rocks of the Kunmarnara and lower Tollu Groups.

659 Most of the rocks are medium-grained, leucocratic ferrogabbros or ferronorites. They have
660 up to 20% intercumulus or oikocrystic magnetite, up to 5% biotite, and several percent sulfide
661 minerals. The rocks tend to be massive, or show a weak, west-trending magmatic foliation.

662 Although whole-rock and mineral compositions are slightly more differentiated than in the upper
663 portions of the Blackstone or Cavenagh intrusions, Cr/V ratios are locally elevated due to
664 chromium enrichment (≤ 1.6 wt%) within magnetite. The rocks have markedly higher
665 concentrations of mica, sulfide, and incompatible trace elements, and Au/PGE ratios than the
666 Cavenagh and Blackstone intrusions. Instead, the Halleys rocks have distinct chemical and
667 petrographic affinities with the Alcurra Dolerite suite.

668 Drilling delineated a pipe-like PGE enriched body, with up to 0.33 wt% Cu and 0.24 ppm PGEs
669 over 74 m, and 0.5 wt% Cu and 0.53 ppm PGE over 16 m (Redstone Resources, 2008, comm. at
670 Annual General Meeting, 27 November). The noble metal patterns are less fractionated than in
671 the magnetite seams of the Bushveld Complex or the Stella, Jameson, and Saturn intrusions,
672 having lower PPGE/IPGE ratios. This is consistent with a magmatic origin of the Halleys
673 mineralization.

674

675 **6.3.12. Nebo–Babel**

676 The 1068 \pm 4 Ma Nebo–Babel intrusion is located about 25 km south of Jameson Community (Fig.
677 2). The Nebo–Babel Ni–Cu–PGE deposit was studied in detail by Seat et al. (2007, 2009) and Seat
678 (2008), and the following section has been compiled mostly from their work.

679 The intrusion has a tubular ('chonolithic') shape traceable for about 5km and trending north-
680 northeast to east. It is 1 km wide and has a stratigraphic thickness of approximately 0.5 km. The
681 chonolith plunges gently to the west-southwest and dips to the south at about 15°. It is offset by
682 the Jameson Fault, dividing it into 2 portions, the Nebo section in the east and the Babel section in
683 the west. Geochemical data indicate that the body is overturned. It was emplaced along a
684 shearzone in felsic orthogneiss of the Pitjantjatjara Supersuite. The magma flow direction was

685 proposed to have been towards the northeast because some of the units thin in this direction and
686 becomes progressively more fractionated.

687 At the stratigraphic base of the intrusion is a breccia zone (MBZ), overlain by a chilled margin
688 (7–9% MgO), variably textured leucogabbronorite (VLGN), melagabbronorite (mela-GN) and
689 barren gabbronorite (BGN), which in the Nebo sector is associated with oxide–apatite
690 gabbronorite (OAGN). The latter constitutes about 20–30% of the intrusion and is characterized by
691 oxide-rich layers that are 5–30 cm thick, with gradational bases and sharp upper contacts. The
692 Babel segment additionally contains the key mineralized gabbronorite unit (MGN) and a 15m thick
693 massive and coarse-grained troctolite unit located between VLGN and BGN in the upper part of
694 the intrusion.

695 In April 2002, Western Mining Corporation announced a drill intersection of 26 m containing
696 2.45% Ni, 1.78% Cu, and 0.09% Co at the Nebo–Babel prospect. The resource estimates are 392 Mt
697 at 0.30% Ni and 0.33% Cu, based on 90 drillholes (Seat et al., 2007). The sulfides consist of
698 monoclinic pyrrhotite, pentlandite, chalcopyrite, and pyrite and occur as massive ores with
699 associated sulfide breccias and stringers, and as disseminated ores, typically forming interstitial
700 blebs in the gabbronorite unit (MGN). Sulfur isotopic data show a remarkably narrow range of $\delta^{34}\text{S}$
701 values from 0 to +0.8‰. The massive sulfides formed through fractional crystallization of a sulfide
702 liquid, resulting in a cumulate of monosulfide solid solution relatively enriched in Os, Ir, Ru, and Rh
703 and depleted in Pt, Pd, and Au.

704

705 **6.3.13. Dyke suites**

706 The dyke suites in the west Musgrave Province have been studied by a number of authors,
707 including Nesbitt et al. (1970), Zhao and McCulloch (1994), Clarke et al. (1995b), Glikson et al.
708 (1996), and Scrimgeour and Close (1999). Howard et al. (2006b) identified seven distinct suites of

709 dykes. The oldest dyke suite (c. 1170 Ma, ~8% MgO) forms part of the Pitjantjatjara Supersuite.
710 Dykes associated with the Giles Event include those belonging to the Alcurra Dolerite suite (6–9%
711 MgO; Zhao et al., 1994; Edgoose et al., 2004) and further include unnamed plagioclase-rich
712 dolerites (~8% MgO) that clearly post-date the G1 intrusions, but may be synchronous with the G2
713 intrusions. Post-Giles dolerites comprise a suite of unnamed olivine- and plagioclase-porphyratic
714 dykes at c. 1000 Ma (~8% MgO), 825 Ma quartz dolerite dykes of the Gairdner-Willouran LIP (~8%
715 MgO), and c. 800 Ma dykes of the Amata Dolerite (Zhao et al., 1994; Glikson et al., 1996; Wingate
716 et al., 1998). A further suite of dykes (~ 9.5% MgO) may be of broadly coeval, or younger, than the
717 Gairdner Dyke Swarm.

718 Godel et al. (2011) distinguished five distinct dyke suites (NB1–5) in the Nebo–Babel area.
719 Types NB1–3 are low-Ti basalts with 5–20% MgO, postulated to be derived from the sub-
720 continental lithospheric mantle (SCLM). Types NB4 and NB5 are high-Ti basalts with 5–14% MgO
721 interpreted to be sourced from a mantle plume. The NB1 type is of approximately similar
722 composition to the plagioclase-phyric dykes of Howard et al. (2006b), containing about 10–13%
723 MgO. NB4 was proposed to represent the Alcurra Dolerite suite.

724

725 **7. Geochemistry of the intrusions**

726 **7.1. Lithophile elements and Nd–Sr–S isotopes**

727 The concentration of the major elements in the G1 layered intrusions is largely controlled by
728 variation in the proportions of olivine, orthopyroxene, clinopyroxene, plagioclase, and magnetite.
729 The modal proportion of plagioclase in the ultramafic rocks is mostly <10%, resulting in relatively
730 high MgO and FeO contents and low Al₂O₃ contents (Fig. 15a). Most of the remaining samples
731 collected are gabbronorites or troctolites with <15% MgO and >10% Al₂O₃. Application of the lever
732 rule indicates that the modal proportion of plagioclase in the latter rocks is typically >50%, with

733 the Cavenagh intrusion being least feldspathic (Fig. 15a,b). Titanomagnetite is an important phase
734 in the Mantamaru, Halleys, and Saturn intrusions, as indicated by high TiO_2 at low MgO and Al_2O_3
735 concentrations (Fig. 15b). Elevated K_2O concentrations in the Hinckley and Murray Range
736 intrusions are likely the result of relatively enhanced crustal contamination (Fig. 15c), whereas the
737 elevated K_2O levels at the Halleys prospect possibly result from advanced fractionation since the
738 country rocks are K-poor mafic intrusive rocks. Relatively high P_2O_5 levels at the Halleys prospect
739 and the Saturn, Jameson, and Blackstone intrusions suggest the presence of apatite (Fig. 15d).
740 Relatively P enriched rocks have also been intersected by drilling along the eastern edge of the Bell
741 Rock intrusion (Pacoe, 2012).

742 The ultramafic intrusions are characterized by relatively high Cr and Ni contents, controlled
743 mainly by olivine, clinopyroxene, and orthopyroxene (Fig. 16). The basal rocks at Wingellina Hills
744 plot along a trend from olivine to orthopyroxene, reflecting their harzburgitic and olivine-
745 orthopyroxenitic composition. The remainder of the ultramafic rocks at Wingellina Hills and
746 Pirntirri Mulari are wehrlites and websterites. Cumulus chromite is largely confined to Wingellina
747 Hills where Cr contents exceed the levels that can be hosted in pyroxene. Most of the gabbroic
748 intrusions contain <1000 ppm Cr, with the exception of samples from the Wingellina Hills and
749 Halleys intrusions. The elevated Cr concentrations at Halleys are the result of abundant Cr-bearing
750 magnetite.

751 In most intrusions, Ni shows a good positive correlation with MgO (Fig. 16b). Samples with
752 significant amounts of olivine are mostly confined to the Wingellina Hills intrusion which have
753 $\text{MgO} > 30$ wt% and $\text{Ni} > 1000$ ppm. The trend of the Wingellina Hills samples in Ni-MgO space can
754 be extrapolated to a Ni concentration in olivine of about 2500–3000 ppm, broadly overlapping
755 with measured olivine compositions from Pirntirri Mulari (~3000 ppm Ni). Two ultramafic samples
756 from Pirntirri Mulari and one from the Morgan Range have distinctly higher Ni contents than the

757 other ultramafic rocks. This is possibly a result of alteration, in view of their relatively high loss-on-
758 ignition (LOI) values. The presence of sulfide and magnetite could explain the elevated Ni levels in
759 the Halleys and Blackstone intrusions (up to ~1500 ppm Ni), compared to the other gabbroic-
760 troctolitic intrusions, which tend to contain <500 ppm Ni. Even higher Ni contents occur at the
761 Nebo Babel Ni-Cu sulfide deposit which have a tenor of 5-6% Ni in the sulfides.

762 The state of differentiation of the intrusions can be compared in a plot of Cr/V ratio vs Mg#
763 (Fig. 17). Wingellina Hills, Pirntirri Mulari, The Wart, and Morgan Range are least evolved, showing
764 some overlap with the Lower Zone of the Bushveld Complex, except that the Bushveld has higher
765 Cr/V ratios due to higher chromite contents. Intrusions with intermediate compositions include
766 Cavenagh, Murray Range, Hinckley Range, the massive G2 gabbros, and the slightly more
767 differentiated Mt. Muir together with the intrusive fragments to the north of Mt Muir and
768 Hinckley Range ('North' and 'Northeast' in Table 1, supplementary data). The most evolved
769 intrusions are Mantamaru (although Bell Rock contains some relatively unevolved samples),
770 Saturn, Halleys, and dykes belonging to the Alcurra Dolerite suite. However, in the Alcurra Dolerite
771 suite there are also relatively unevolved samples that have up to 9 wt% MgO.

772 The mafic intrusions show fractionated lithophile multi-element (spider) patterns, with
773 relative enrichments in the most incompatible elements, but negative Nb anomalies (Maier et al.,
774 2014). Positive Ti anomalies are found at Mantamaru, Halleys, and Lehman Hills and in many
775 samples from Saturn, reflecting the presence of magnetite. The Saturn and Halleys intrusions have
776 distinctly elevated incompatible trace element contents relative to most other intrusions,
777 including Cavenagh and Blackstone, consistent with a distinct magmatic lineage. Notably, the trace
778 element patterns of gabbros from Wingellina Hills are identical to those of Wingellina Hills
779 pyroxenites, indicating crystallization from magmas of broadly similar composition.

780 Data from sample suites that contain a higher liquid component are plotted in Fig. 18. The
781 Alcurra Dolerite suite (Fig. 18a), G2 gabbros (Fig. 18b), NB1 dykes, and the fine grained marginal
782 rocks from Nebo–Babel show considerable similarity including pronounced negative Nb and Ta
783 anomalies and, in the case of the G2 gabbros and Nebo–Babel chilled margins, negative Ti
784 anomalies. In all suites, the incompatible trace element contents are typically higher than in the
785 G1 intrusions, but the shapes of the multi-element patterns for all mafic intrusives are remarkably
786 similar. Microgabbros from Cavenagh have less fractionated, but more ‘spiky’ patterns than the
787 other liquid-rich mafic rocks (Fig. 18d). This reflects the elevated cumulate component in most
788 microgabbros.

789 Mantamaru has systematically higher ϵNd (0 to +2) and lower Ce/Nb ratios (mostly 2–7) than
790 the other intrusions (Fig. 19). Some of the least radiogenic Nd isotope compositions occur in the
791 G2 gabbros and the G1 Cavanagh and Morgan Range intrusions ($\epsilon\text{Nd} = -1$ to -4 , Ce/Nb = 3–13),
792 and in the Kalka intrusion in South Australia (Wade, 2012). Rocks of the Alcurra Dolerite suite have
793 intermediate compositions ($\epsilon\text{Nd} = -1$ to +2, Ce/Nb \sim 3–5). Most Nebo–Babel samples have
794 compositions overlapping with the Alcurra Dolerite suite ($\epsilon\text{Nd} = -1.7$ to 0.3, Ce/Nb = 5–7, except
795 for the marginal rocks, which show lithological evidence for crustal assimilation and have ϵNd as
796 low as -3). The data of Seat et al. (2011) indicates that basement rocks at Nebo–Babel have ϵNd
797 values of -4.5 to -5 and Ce/Nb ratios of 9, whereas the regionally extensive Pitjantjatjara granite
798 suite has ϵNd of -2 to -4 and highly variable Ce/Nb (5 to >20).

799 *In situ* Sr isotope analyses on plagioclase (Maier et al., 2014) are consistent with these
800 results in that the least-radiogenic Sr isotopic values are found in the Mantamaru and Halleys
801 intrusions, whereas Morgan Range, Lehman Hills, and Cavenagh have higher initial Sr isotope
802 ratios. Microgabbros from Cavenagh have less radiogenic Sr isotope ratios ($\text{Sr}_i = (87\text{Sr}/86\text{Sr})_i =$
803 $0.7042 - 0.7057$) than associated medium-grained gabbros ($\text{Sr}_i = 0.7052 - 0.7068$), and the

804 medium-grained samples show greater isotopic heterogeneity. This is interpreted to reflect lining of
805 the magma conduits by early, relatively contaminated magma pulses, allowing the relatively late-
806 stage microgabbro magma to be emplaced while undergoing relatively less crustal interaction. Of
807 note is that almost the entire range in Sr isotopic ratios seen within the west Musgrave mafic-
808 ultramafic intrusions is present within the Kalka intrusion in South Australia, and in both cases, the
809 ultramafic rocks have the highest initial Sr isotope ratios, whereas anorthosites, leucogabbros, and
810 troctolites have relatively more mantle-like compositions.

811 Whole-rock sulfur isotope data have been generated for rocks of the Jameson intrusion, the
812 G2 gabbros, and the Alcurra Dolerite suite (Table 3, supplementary data). In addition, we collected
813 sulfur isotope data for sulfide-bearing rhyolites from the Bentley Supergroup in the Palgrave area
814 (Fig. 2), as well as *in situ* (laser ablation ICP-MS) sulfur isotope data from Halleys. These data can
815 be compared to those from Nebo-Babel (Seat et al., 2009). All mafic intrusive rocks plot near the
816 composition of the mantle. By contrast, rhyolites of the Mount Palgrave Group have a much wider
817 sulfur isotopic range of $\delta^{34}\text{S}$, from +3.2 to +7. The data could either suggest that the sulfides in the
818 mafic rocks are of mantle derivation, or that any crustal sulfides were juvenile or underwent no
819 sulfur isotopic fractionation, or that assimilated crustal sulfides equilibrated with the magma at
820 high R-factors (mass ratio of silicate melt to sulfide melt; Campbell and Naldrett, 1979; Leshner and
821 Burnham, 2001). Notably, recent data from the Manchego prospect (Karykowski, 2014) show
822 strong negative $\delta^{34}\text{S}$, representing the only igneous suite amongst the Giles intrusions with
823 markedly non-magmatic S isotope signatures.

824

825 **7.2. Sulfur and chalcophile elements**

826 Most of the Giles intrusions have relatively low sulfur concentrations, at <200 ppm (Fig. 20a).

827 Slightly higher sulfur concentrations are present at Wingellina Hills, The Wart, Murray Range,

828 Hinckley Range, the 'North' and 'Northeast' intrusive fragments, some of the G2 massive gabbros,
829 and the Alcurra Dolerite suite (up to 2000 ppm). Even higher sulfide contents (in places > 1 vol. %)
830 occur at Saturn, and in olivine gabbro and olivine gabbro-norite of the upper Jameson intrusion,
831 with Cu concentrations up to 860 ppm (at 0.12 wt% SO₃). The relatively high sulfide and Cu
832 contents could be due to protracted fractionation and resulting saturation in sulfide liquid in the
833 magma, analogous to the troctolitic Kiglapait intrusion in Labrador, where sulfur saturation is
834 reached after 93% fractionation (Morse, 1981). Alternatively, the upper stratigraphic portions of
835 the Jameson Range could have undergone incipient hydrothermal alteration and addition of Cu
836 and S, possibly related to the voluminous volcanic activity that formed the Mount Palgrave Group,
837 directly to the southwest. However, sulfur isotopic data for two troctolitic samples (GSWA 189475
838 and 189478) indicate $\delta^{34}\text{S}$ of between +2.1 and +2.8, broadly consistent with a magmatic origin.

839 No sulfur data are available for Halleys, but petrographic examination indicates locally
840 several percent sulfides, consistent with Cu concentrations of >4000 ppm in some samples.
841 Sulfide-rich mafic intrusive rocks, locally containing net textured and massive sulfides were
842 recently intersected at the Manchego Prospect (Phosphate Australia, 2014; Karykowski, 2014). The
843 highest sulfide contents among the Giles intrusions occur at Nebo–Babel, including thick intervals
844 of massive and disseminated sulfides (Seat et al., 2007).

845 The mafic-ultramafic intrusions are typically relatively Cu poor (<200 ppm Cu, Fig. 20b). The
846 relatively unevolved (ultramafic) rocks have particularly low Cu contents, whereas Cu contents
847 progressively increase in the evolved (mafic) rocks, consistent with incompatible behavior of Cu in
848 fractionating sulfur-undersaturated magma. The highest Cu contents are found in the Nebo Babel
849 (sulfide tenor of 2-8% Cu), Halleys and Manchego intrusions (Seat et al., 2007, Karykowski, 2014).
850 Slightly lower Cu contents in some samples from Saturn, and the uppermost portions of the
851 Jameson intrusion, where some of the massive magnetite layers contain up to 700 ppm Cu.

852 Elevated Cu concentrations are also found in the Wingellina Hills PGE reefs (up to 500 ppm Cu),
853 and in a pyroxenite from the upper portion of Pirntirri Mulari that has 350 ppm Cu.

854 The majority of the Giles intrusions have low PGE contents (<30 ppb Pt+Pd, Fig. 20c). Higher
855 values occur in the PGE reefs of the Wingellina Hills intrusion, containing up to several ppm PGE,
856 the pyroxenite in the upper portion of Pirntirri Mulari (200 ppb PGE, not shown in Fig. 20c due to
857 lack of major element data), and in samples from Halleys (up to 200 ppb PGE). Other PGE-rich
858 rocks not plotted include those from Nebo–Babel (up to 0.5 ppm PGE in whole rocks, up to ~7 ppm
859 PGE in sulfide) and Manchego (up to approximately 1 ppm PGE in sulfide, Karykowski, 2014). The
860 lowermost magnetite layer in the Jameson intrusion has up to about 2 ppm PGE, at very low
861 sulfide contents. Scattered PGE enrichment, not accompanied by sulfide enrichment, occurs in the
862 Morgan Range (≤ 80 ppb), The Wart (one sample with 120 ppb), and Cavenagh (three samples with
863 75–100 ppb).

864 In almost all intrusions, including the sulfide-bearing Nebo Babel, Halleys and Manchego
865 intrusions, Cu/Pd ratios are above the range of the primitive mantle (~7000); thus, they are PGE-
866 depleted relative to mantle (Fig. 20d). This could suggest that the magma had equilibrated with
867 sulfide prior to final emplacement, in the mantle or the crust, or that it assimilated Cu-rich crust,
868 or that the mantle source was relatively enriched in Cu. Cu/Pd ratios progressively increase with
869 decreasing Mg# and samples with Cu/Pd < primitive mantle are mostly confined to Wingellina
870 Hills, Pirntirri Mulari, and the Morgan Range, as well as some Cavenagh samples, that is, rocks with
871 Mg# mostly >60. We interpret this to reflect sulfide liquid saturation in response to primarily
872 magmatic fractionation rather than contamination. We do not consider it likely that the PGE
873 depletion is due to a relatively small degree of mantle melting, as NB1 is strongly S undersaturated
874 and has lower TiO₂ contents (0.8%) than typical MORB (>1%, Gale et al., 2013).

875 Most of the sample suites that represent liquids rather than cumulates (NB1 dykes, the
876 Alcurra Dolerite suite, and Nebo–Babel chilled margins) are PGE depleted, with Cu/Pd ratios above
877 primitive mantle values. The main exceptions are the unevolved G2 gabbros, which contain up to
878 10–15 ppb Pt and Pd each and have Cu/Pd ratios overlapping with primitive mantle. Notably, Nebo
879 Babel chilled margins too have PGE concentrations typical of basaltic magmas (~10–20 ppb Pt and
880 Pd each), but Cu/Pd > 10 000.

881

882 **7.3. Mineral chemistry**

883 We determined the compositions of olivine, orthopyroxene, clinopyroxene, and plagioclase in
884 more than 50 rock samples from the Giles intrusions, excluding the the Alcurra Dolerite suite and
885 the G2 massive gabbros. A summary of some key compositions are given in Table 1, and a detailed
886 discussion of the data can be found in Maier et al. (2014). In the present paper, we focus on
887 discussing olivine compositions. The mineral has between 40 and 87 mole % Fo (Fig. 21). Olivine
888 from the Blackstone intrusion and parts of Cavenagh show the lowest Fo values, whereas the
889 highest Fo contents occur at Wingellina Hills, Pirntirri Mulari, The Wart, and Morgan Range.
890 Samples from Jameson, Saturn, Hinckley Range, Murray Range, Nebo–Babel, Lehman Hills, and
891 Latitude Hill contain olivine with intermediate Fo contents.

892 Olivine from the mafic and mafic-ultramafic Giles intrusions has up to 3500 ppm Ni (Fig.
893 21a). Ni contents show a positive correlation with Fo content and are higher than in olivines of
894 comparable Fo content from many basic magmas globally, although relatively high Ni
895 concentrations appear to be characteristic of many layered intrusions (Fig. 21b). The highest Ni
896 contents in magmatic olivine identified so far occurs in the Kevitsa intrusion of Finland (with up to
897 1.5 wt% Ni; Yang et al., 2013).

898 For the origin of the Ni enrichment in olivine of the layered intrusions several models may be
899 considered. (i) Equilibration of olivine with trapped melt could lower the Fo content without
900 significantly affecting Ni concentrations. This model would be consistent with the observed
901 decoupling of Fo from Ni contents in olivine, and from An contents of plagioclase. However, Godel
902 et al. (2011) found Ni enrichment in olivines from the NB1 dykes. Thus, the observed Ni
903 enrichment must, at least in part, reflect an early magmatic process. (ii) Equilibration of olivine
904 with percolating sulfide liquid. However, there is little evidence for sulfide in most intrusive rocks
905 related to the Giles Event. (iii) Assimilation of Ni-rich sulfide before final magma emplacement.
906 This model is presently also considered unlikely as the West Musgrave crust is relatively poor in
907 magmatic Ni sulfides. (iv) Magma derivation from a pyroxenitic–eclogitic mantle source (Sobolev
908 et al., 2011). This model is equally rejected as it implies that most layered intrusions globally are
909 derived from pyroxenitic mantle sources, for which there is presently no evidence. (v)
910 Contamination leading to magma reduction and relatively low Fo contents. However, the oxygen
911 fugacity of the Giles layered intrusions is thought to be similar to most other mafic-ultramafic
912 crustal rocks, at around the quartz-fayalite-magnetite buffer (Staubman, 2010). (vi) Polybaric
913 fractionation, with initial high-P crystallization of pyroxene at depth leading to depletion of the
914 magma in MgO relative to Ni, followed by ascent and final emplacement of the magma at low
915 pressure conditions where olivine may become stable. Such a magma, and any olivine crystallising
916 from it, could have relatively high Ni to MgO ratios (Maier et al., 2013). In addition, the D_{Ni} into
917 olivine increases with falling pressure (Li and Ripley, 2010). The model would be consistent with
918 the distinctly lower Cr/Al ratios of pyroxenes in the Giles intrusions relative to the Bushveld
919 Complex (Maier et al., 2014), potentially reflecting high-P pyroxene crystallization. Ballhaus and
920 Glickson (1995) noted that olivine in the Giles intrusions shows a continuous range of Fo contents,
921 in contrast to other layered intrusions such as the Bushveld Complex which lack olivines of

922 composition Fo 60–80, normally interpreted to be the result of the peritectic reaction of olivine to
923 pyroxene, temporarily destabilizing olivine. The authors proposed that the lack of an “olivine gap”
924 in the Giles intrusions is due to polybaric fractionation.

925

926 **8. Discussion**

927 **8.1. Nature of parent magmas to the Giles intrusions**

928 Knowledge of the composition of the parent magma to cumulate rocks is of considerable interest
929 to petrologists and economic geologists because this potentially allows to constrain the nature of
930 the mantle source, the degree of crustal contamination of the magma, the crystallization history of
931 an intrusion, and the prospectivity for magmatic mineral deposits. One of the most common
932 approaches to estimate the parent magma composition is based on the study of chilled contact
933 rocks of intrusions. However, such rocks are commonly contaminated and thus not necessarily
934 representative of the primary magma. Another approach could be to examine microgabbroic rocks
935 that are abundant within several Giles intrusions, but many of these rocks contain a cumulate
936 component.

937 A technique that has been successfully applied in the Bushveld Complex and more recently
938 in the Giles intrusions comprises the study of fine-grained sills or dykes associated with the
939 intrusion (Sharpe, 1981; Barnes et al., 2010). For the ultramafic segments of the G1 intrusions, a
940 suitable parent magma candidate would be the fine-grained low-Ti tholeiitic ‘plagioclase-rich
941 dykes’ initially documented in the Bell Rock Range area by Howard et al. (2007). These dykes are
942 compositionally equivalent to the NB1 dyke type of Godel et al. (2011). A further suitable parent
943 magma type could be the unevolved members of the massive G2 gabbros. They contain ~ 10–13
944 wt% MgO, 12 wt% FeO_T, 350 ppm Ni, up to 700 ppm Cr, and 10–15 ppb Pt and Pd each.

945 Crystallization of the Wingellina Hills intrusion from either NB1- or primitive G2-type magma is

946 consistent with the relatively low Ti content of these magmas (Fig. 53 in Maier et al., 2014).
947 Modeling using the PELE software (Boudreau, 1999) indicates that at low to intermediate pressure
948 NB1 has a crystallization order of chromite > olivine+chromite > olivine+chromite+clinopyroxene >
949 chromite+clinopyroxene+plagioclase+orthopyroxene. This is broadly consistent with petrographic
950 observations on the Pirntirri Mulari and Wingellina Hills intrusions, namely the occasional
951 occurrence of chromite grains within olivine. The modeled Fo content at 1–5 kb is 87 mol.%,
952 consistent with analyses. Furthermore, calculations by Godel et al. (2011) indicate that NB1
953 reaches sulfur saturation after 30% crystallization, at about the time plagioclase appears on the
954 liquidus, consistent with the stratigraphic position of the PGE reefs at the top of the ultramafic
955 zone in the Wingellina Hills intrusion. None of the other c. 1070 Ma dyke suites analysed by
956 Howard et al. (2006b) or Godel et al. (2011) provides a suitable fit for the ultramafic intrusions.
957 Considering the parental magmas to the gabbroic intrusions, possible candidates are the Nebo–
958 Babel chilled margin (7–9 wt% MgO, Mg# of 51–61) or the compositionally similar NB3 dyke type.
959 For the Halleys and Saturn intrusions, the Alcurra Dolerite suite (or the compositionally equivalent
960 NB4 dyke type of Godel et al., 2011) would be a potential parent magma, based on similarities in
961 incompatible trace element and noble metal ratios (e.g., high Cu/Pd) and common enrichment in
962 mica and sulfide (Howard et al., 2009). The most primitive members of the Alcurra Dolerite suite
963 have 8–9 wt% MgO and Cr/V ratios of 2–4 (Table 1, supplementary data).

964

965 **8.2. Mantle sources of the magmas**

966 Godel et al. (2011) proposed that NB 1 magmas were derived from the sub-continental
967 lithospheric mantle (SCLM). Their model was based on the relatively low Ti contents and MREE to
968 HREE ratios in the magmas, implying the presence of amphibole in the source. However, several
969 arguments can be made against this model: (i) Magmas believed to be derived from the SCLM, e.g.

970 Bushveld B1 magmas, have much lower S contents (400 ppm, Barnes et al., 2010) than NB1, which
971 has S contents typical of many other global basalts (~1000 ppm; Godel et al., 2011). (ii) SCLM
972 derived magmas may have high Pt/Pd above unity (Barnes et al., 2010) whereas NB1 has Pt/Pd
973 below unity, in the range of most other basalts. (iii) Between c. 1220 and c. 1120 Ma, the west
974 Musgrave Province experienced ultra-high temperature (UHT) metamorphism in the middle crust
975 (Kelsey et al., 2009). Smithies et al. (2010, 2011) argued that this requires removal of the regional
976 SCLM, consistent with a dramatic lowering in the pressure of crustal melting at the beginning of
977 the Musgrave Orogeny. If there was SCLM at the beginning of the Giles Event, it must have formed
978 after the Musgrave Orogeny, which means it would have been still young, hot, and weak, and have
979 too radiogenic Nd isotopic compositions to be parental to the NB1 dykes ($\epsilon_{Nd} = -2$). Based on
980 these data, we do not subscribe to the model of Godel et al. (2011). The evidence for an
981 asthenospheric mantle source to NB1 is more persuasive, e.g., the resemblance of Yb–Ti–Zr–Nb
982 concentrations of NB1 to those in MORB. The enrichment in LILE and LREE within NB1 could be
983 modeled by low degree (<<5%) crustal contamination of asthenospheric high-Mg basalt.
984 The origin of the magma forming the Alcurra Dolerite suite also remains unresolved. Godel et al.
985 (2011) proposed that the NB4 dykes, which they considered analogues of the Alcurra Dolerite
986 suite, were plume melts. The main argument rests on the relatively high Ti contents and
987 MREE/HREE ratios of NB4, ostensibly requiring the presence of residual garnet in a deep mantle
988 source. However, the high La/Yb ratios of the Alcurra-type rocks (5 - 9.7) are not accompanied by
989 depletions in Yb (average 3.34 ppm) with respect to MORB and are thus better explained through
990 minor contamination (<<1%) with HFSE-rich crust. Crustal contamination could potentially also
991 explain the relatively high Cu/Pd and Au/PGE ratios of the Alcurra-type magmas, whereas the
992 origin of the elevated Pt/Pd remains presently poorly understood.

993

994 **8.3. Crustal contamination of the magmas**

995 The Giles intrusions have ϵNd from +2 to -5 (Fig. 19 and Table 4, supplementary data). These data
996 could be explained by variable crustal contamination, or melting of compositionally diverse mantle
997 sources, or both. At least some degree of *in situ* crustal contamination is suggested by the field
998 evidence, e.g. the mingling of the G2 gabbros with granite and abundant xenoliths in several G1
999 intrusions such as Hinckley Range (Fig. 6) and Kalka (Gray and Goode, 1989).

1000 In the ultramafic intrusions, crustal contamination is suggested by the existence of 2 distinct
1001 crystallization paths. The basal portions of the Wingellina Hills and Kalka (South Australia)
1002 intrusions, and most of the central portions of Pirntirri Mulari, have a crystallization sequence of
1003 olivine > orthopyroxene+olivine+chromite > orthopyroxene+clinopyroxene, in contrast to the
1004 central portion of the Wingellina Hills intrusion (Ballhaus and Glickson, 1995) which has a similar
1005 crystallization order as that modeled (using PELE) for NB1, i.e. chromite - olivine+chromite -
1006 olivine+chromite+clinopyroxene - chromite+clinopyroxene+plagioclase+orthopyroxene. Two
1007 distinct liquid lines of descent have also been found at the Muskox intrusion, Canada, where the
1008 basal rocks show a crystallization sequence of olivine > clinopyroxene+olivine, whereas the upper
1009 units show olivine > olivine+orthopyroxene, interpreted to result from contamination with partial
1010 melts of the roof (Irvine, 1970).

1011 Among the gabbroic G1 intrusions, Cavenagh has the lowest ϵNd (ϵNd as low as -5),
1012 overlapping with those of basalts of the Mummawarrawarra and Glyde Formations of the Bentley
1013 Supergroup (and Warakurna Supersuite). As there are no Musgrave crustal rocks currently known
1014 to have ϵNd values below -6, the required degree of contamination of the Cavenagh magma may
1015 seem unrealistically high. However, granites of the regionally occurring Pitjantjatjara Supersuite
1016 are extremely rich in HFSE, which may greatly reduce the required amounts of contamination
1017 (Kirkland et al., 2013). Relatively strong contamination of the Cavenagh intrusion is consistent with

1018 the high Ce/Sm and Ce/Nb ratios (Fig. 22). In contrast, Nd and Sr isotopic data for the troctolitic
1019 intrusions (namely Mantamaru) approximate the chondritic uniform reservoir (CHUR) (ϵ_{Nd} mostly
1020 from 0 to +2, $I_{\text{Sr}} \sim 0.704$), have markedly higher ϵ_{Nd} or lower I_{Sr} than any Musgrave crust present
1021 at the time and relatively low Ce/Sm and Ce/Nb ratios (Fig. 22). These data indicate very minor
1022 (<5%) crustal contamination in most troctolitic intrusions.

1023 The Alcurra Dolerite suite has slightly lower ϵ_{Nd} (+1 to -1) than the Mantamaru intrusion.
1024 Smithies et al. (2013) showed that less than 10% bulk contamination of the most primitive samples
1025 of the Alcurra Dolerite suite with average Pitjantjatjara Supersuite granite can explain the entire
1026 isotopic variation and much of the highly incompatible trace element variation within the Alcurra
1027 Dolerite suite. If one assumes as contaminant a low-degree (20%) partial melt of average
1028 Pitjantjatjara Supersuite granite, the required assimilation is <4%. Such low degrees of
1029 contamination would produce only slight shifts to lower ϵ_{Nd} isotope values, and it would thus
1030 appear unlikely that the mantle source was strongly depleted. Thus, the unevolved magmas of the
1031 Alcurra Dolerite suite were likely derived by relatively shallow melting (<80 km) of weakly
1032 depleted mantle, followed by early and very minor (<4%) contamination with highly enriched
1033 crustal material, and then closure of the continuously fractionating system to further
1034 contamination. The latter process can be applied to all Giles intrusions, consistent with the broad
1035 similarity in ϵ_{Nd} within individual bodies, at variable Ce/Nb and La/Sm.

1036

1037 **8.4. Magma emplacement**

1038 The emplacement conditions of the Giles intrusions were discussed by Maier et al. (2014), from
1039 which the following summary has been compiled. Field relationships and compositional data
1040 indicate that the depth of emplacement varied considerably between intrusions. In the case of
1041 Mantamaru, country-rock inclusions and intrusive contacts indicate that the body was emplaced

1042 at the stratigraphic level of the Mummawarrawarra Basalt (Kunmarnara Group). Based on the low
1043 metamorphic grade (greenschist facies) of the basalts, it is argued that the Mantamaru intrusion
1044 crystallised in an upper crustal, extensional environment. Constraints on its crystallization age are
1045 the minimum depositional age of the Kunmarnara Group (defined by intrusion of granite at c.
1046 1078 Ma; Sun et al., 1996; Howard et al., 2011b), and a direct U–Pb zircon date of 1076 ± 4 Ma
1047 (Kirkland et al., 2011; GSWA 194762).

1048 Particularly in the Hinckley Range, in the eastern part of the study area, massive G2 gabbro
1049 cuts the layered G1 intrusions and tends to show evidence of comingling with leucogranite. The
1050 latter may form pluton-scale bodies in the basement, for example the Tollu pluton. In the vicinity
1051 of major shear zones this bimodal gabbroic-granitic magmatism was accompanied by shearing and
1052 west-northwest folding, at between 1078 ± 3 and 1074 ± 3 Ma (Howard et al., 2011b). These dates
1053 overlap with the crystallization ages of the layered (G1) intrusions, but field relationships indicate
1054 that the G2 intrusions always post-date G1.

1055 In the Blackstone Sub-basin, to the south of Blackstone Community, rhyolites of the Smoke
1056 Hill Volcanics directly overlie the G1 Blackstone Range without an obvious fault. Crystallization
1057 ages for the rhyolites [1071 ± 8 Ma (GSWA 191728; Coleman, 2009); 1073 ± 7 Ma (GSWA 191706;
1058 Coleman, 2009), and 1073 ± 8 Ma (GSWA 189561; C Kirkland, 2014, written comm.) are within
1059 analytical error of the emplacement date of the G1 and G2 intrusions, and the composition of the
1060 rhyolites resembles that of leucogranites associated with the G2 intrusions. In addition, several
1061 field exposures indicate that the Blackstone intrusion was emplaced into the lower basaltic
1062 portions of the Kunmarnara Group (Bentley Supergroup). This requires extensive and rapid crustal
1063 uplift, erosion, and exhumation of the layered G1 intrusions, immediately followed by felsic
1064 volcanism.

1065 Previous authors have proposed that some of the ultramafic Giles intrusions have been
1066 emplaced at relatively high pressure (Goode and Moore, 1975; Ballhaus and Berry, 1991), based
1067 on high Al concentrations of pyroxene, spinel exsolution in pyroxene and plagioclase, rutile
1068 exsolution in pyroxene, antiperthitic exsolution in plagioclase, and orthopyroxene–clinopyroxene–
1069 spinel–albite coronas between olivine and plagioclase. For example, Goode and Moore (1975)
1070 suggested that the Ewarara intrusion was emplaced at a pressure of 10–12 kbar. If the thickness of
1071 the crust at the end of the Musgrave Orogeny, just 40my before the Giles event, was 35 km
1072 (Smithies et al., 2011), the Ewarara intrusion would have intruded near the base of the crust.
1073 Ballhaus and Glikson (1989) proposed an emplacement depth of 6.5kbar (~20km) for the
1074 Wingellina Hills and Pirntirri Mulari intrusions. However, because the coeval gabbroic intrusions
1075 are up to 10 km thick, the emplacement depth of the ultramafic bodies may have been as shallow
1076 as 10 km.

1077 Emplacement into relatively deep crustal levels could explain why the ultramafic intrusions
1078 are less abundant than the gabbroic and troctolitic bodies, and why the former are proportionally
1079 more abundant in South Australia than in Western Australia - the South Australian crust is exposed
1080 at a deeper level (Goode, 2002). Equally consistent with this model is the observation that the
1081 ultramafic intrusions tend to be exposed in the cores of regional folds (i.e., the anticline north of
1082 Blackstone Community which hosts the Pirntirri Mulari and Morgan Range intrusions), or along
1083 faults.

1084 Field relationships also allow to place some constraints on emplacement dynamics. For
1085 example, the close spatial association of many microgabbros with fragments and schlieren of
1086 pyroxenite suggests that the emplacement of the microgabbros was accompanied by
1087 disaggregation of semi-consolidated pyroxene-rich cumulate slurries. This points to a semi-
1088 consolidated magma chamber that was frequently replenished by unevolved magma.

1089

1090 **8.5. Fragmentation of intrusions**

1091 The idea that some or all of the Giles layered intrusions could be tectonically dismembered
1092 remnants of a much larger body can be traced back to Sprigg and Wilson (1959) and Nesbitt and
1093 Talbot (1966). Other authors who favoured this model include Glikson (1995), Smithies et al.
1094 (2009), Howard et al. (2011b) and Aitken et al. (2013). Geophysical, lithological, and compositional
1095 data reported in the present study indicate that the Jameson–Finlayson, Blackstone, and Bell Rock
1096 intrusions are fragments of an originally contiguous body, named the Mantamaru intrusion by
1097 Maier et al. (2014). This has a minimum preserved size of 3400 km², in the same range as the
1098 Great Dyke, Stillwater, Sept Iles, and Dufek intrusions which measure between 3000–5000 km². It
1099 has been further proposed that the Cavenagh intrusion may form the southern limb of the
1100 synclinal Blackstone intrusion (Nesbitt and Talbot, 1966; Aitken et al., 2013), potentially adding at
1101 least another 540 km² to the size of the Mantamaru intrusion. However, the Blackstone intrusion
1102 is much more differentiated than the Cavenagh intrusion, and it contains a massive magnetite
1103 layer that appears to be absent from the Cavenagh intrusion. Furthermore, the Cavenagh intrusion
1104 shows elevated PGE concentrations in its upper portion whereas the Blackstone intrusion is
1105 uniformly PGE depleted. Finally, the Blackstone intrusion has distinctly higher εNd values than the
1106 Cavenagh intrusion. These data place some doubts on a possible connection between the
1107 Cavanagh and Blackstone intrusions, although other layered intrusions such as Bushveld and Kalka
1108 can have a wide range of trace element and isotopic compositions.

1109 Based largely on compositional data, Seat (2009) proposed that the Nebo and Babel intrusive blocks
1110 are overturned fragments of an originally contiguous intrusion. This model is consistent with field evidence
1111 from, e.g., the western end of the Hinckley intrusion and the NW edge of the Blackstone Range, that
1112 deformation accompanied magmatism, locally developing open to tight folds.

1113 It is tempting to speculate that all the ultramafic intrusions located in the Tjuni Purlka Zone
1114 (Wingellina Hills, Pirntirri Mulari, The Wart, Morgan Range, Kalka, Gosse Pile, and Ewarara)
1115 originally formed a single body. This would imply lateral movement of up to 50 km within the Tjuni
1116 Purlka Zone. The intrusions share some similarities (e.g., olivine and pyroxene compositions,
1117 stratigraphic position of the Wingellina PGE reef and the Cu–PGE-rich horizon at Pirntirri Mulari)
1118 but they also show differences (e.g., thicker olivine and chromite rich segments and higher PGE
1119 concentrations at Wingellina Hills than in the other bodies, variation in plagioclase composition
1120 between Wingellina Hills and The Wart vs. Pirntirri Mulari). Further work is required to resolve this
1121 question.

1122

1123 **8.6. Comparison to other large layered intrusions**

1124 In order to better understand the petrogenesis and prospectivity of the Giles intrusions, it is useful
1125 to draw comparisons with other well-characterized and mineralized layered intrusions, e.g. the
1126 Bushveld Complex (Fig. 23). The Pirntirri Mulari, Wingellina Hills, and The Wart intrusions are the
1127 approximate stratigraphic and compositional equivalents of the Lower and Critical Zones of the
1128 Bushveld Complex. They have broadly similar olivine compositions and they show basal
1129 compositional reversals, thick ultramafic portions, and a number of ultramafic–mafic cyclic units.
1130 Reef-style PGE enrichments have been identified in the Wingellina Hills intrusion, and there are
1131 some indications that a similar horizon may exist in the Pirntirri Mulari intrusion. The equivalent
1132 prospective horizon of The Wart, Kalka, Ewarara, Gosse Pile, Ngulana, and Alvey Hills remains
1133 poorly studied, partly due to restricted access.

1134 A major difference between the Giles ultramafic intrusions and the Bushveld Complex is that
1135 the former appear to lack chromitite seams. If this is due to the early crystallization of Cr-rich

1136 clinopyroxene, as suggested by Ballhaus and Glickson (1995), this would imply a low prospectivity
1137 for chromite deposits in the Giles intrusions.

1138 The gabbroic Morgan Range intrusion contains a lens of ultramafic rocks at its northern edge
1139 and thus could represent the stratigraphic equivalent to the Upper Critical Zone - Main Zone
1140 transition of the Bushveld Complex, unless the ultramafic lens was tectonically adjoined to the
1141 Morgan Range. The lens does have some potential to host a PGE reef analogous to Wingellina
1142 Hills. In contrast, the Cavenagh, Michael Hills, Latitude Hill, Hinckley Range, Murray Range,
1143 Lehman Hills, and Mt Muir intrusions have lower PGE prospectivity as they appear to be
1144 stratigraphic equivalents of the Bushveld Main Zone, sharing intermediate compositions and
1145 relatively subdued layering.

1146 The Mantamaru intrusion is stratigraphically approximately equivalent to the upper Main
1147 Zone and Upper Zone of the Bushveld Complex. Both intrusions contain several magnetite layers
1148 (~25 in the Bushveld, at least 11 at Jameson) that can reach a thickness of more than 10 m (i.e.,
1149 layer 1 at Jameson, magnetite layer 21 in Bushveld). In addition, in both intrusions the vanadium
1150 concentration progressively decreases from the basal to the upper magnetite layers (Fig. 19), and
1151 also within individual layers. However, the vanadium contents of the Bushveld Main Magnetite
1152 Layer are twice as high as those at Jameson (13000 vs 7500 ppm). Other differences between the
1153 two intrusions include significantly higher PGE contents in the Jameson basal magnetite seam and
1154 the apparent absence of an apatite-rich layer at Jameson.

1155

1156 **8.7. Tectonic setting**

1157 The Musgrave Province is located between the lithospheric keels of the West Australian, South
1158 Australian, and North Australian Cratons. Channeling of mantle plumes along the cratonic keels
1159 could have resulted in strong adiabatic mantle partial melting and mafic magmatism (Begg et al.,

1160 2009). However, Smithies et al. (2012) argued that the >200Ma time span of continuing mantle
1161 magmatism and UHT metamorphism is inconsistent with a mantle plume. The Musgrave Province
1162 instead represented a stationary zone of mantle upwelling, resulting in a persisting hot zone.
1163 Magmatism may have been driven by processes such as plate motions, lithospheric delamination,
1164 volatile transfer from the SCLM or the crust to the convecting mantle, or mantle flow along the
1165 irregular base of the lithosphere (Silver et al., 2006). Similar scenarios were envisaged by Silver et
1166 al., (2006) and Foulger (2010) for the Ventersdorp, Great Dyke, Bushveld, and Soutpansberg
1167 continental magmatic events in southern Africa.

1168 The stage was set during and after the 1345–1293 Ma Mount West Orogeny which resulted
1169 in crustal thickening, partial melting, and densification of lower crust. The REE geochemistry of
1170 Musgrave granites suggests that at the beginning of the 1220–1150 Ma Musgrave Orogeny the
1171 depth of crustal melting changed from relatively deep to shallow, caused by delamination of
1172 residual lower crust and the underlying lithospheric mantle. The ensuing UHT metamorphism from
1173 1220 to 1120 Ma testifies to a sustained regime of highly thinned crust and mantle lithosphere.
1174 Any magmatism was predominantly felsic because the lower crust had become a zone of melting,
1175 assimilation, storage, and homogenization (MASH), inhibiting ascent of relatively dense mafic
1176 magmas. This crustal thermal structure strongly influenced conditions at the beginning of the Giles
1177 Event (Smithies et al., 2015, in press). The latter was triggered by far-field forces acting on the
1178 margins of the West Australian Craton (Evins et al., 2010b; Smithies et al., 2015, in press). Initial
1179 subsidence and deposition of the Kunmarnara Group was followed by draining of melts ponded at
1180 the base of the crust (G1, G2, Alcurra Dolerite suite). The relatively early G1 and G2 magmas were
1181 variably contaminated during ascent into the crust (Fig. 24). Subsequent magmas of the Alcurra
1182 Dolerite suite underwent relatively little contamination suggesting that the crust had become
1183 more refractory. At the same time, the Alcurra magmas are more differentiated because the crust

1184 thickened during the Giles event, allowing more intra-crustal ponding and fractional crystallization.
1185 The relatively low PGE concentrations, high Cu/Pd, Pt/Pd, and Au/PGE ratios of the Alcurra
1186 magmas could be explained by melting of hybrid crust-rich mantle, in response to foundering of
1187 crust and new SCLM (Fig. 24a).

1188

1189 **8.8. Origin of mineralization**

1190 **8.8.1. PGE reefs within the Wingellina Hills layered intrusion**

1191 The bulk of the world's PGE resources occur in the form of stratiform layers or so-called reefs
1192 hosted by layered mafic-ultramafic intrusions. Economic deposits are presently confined to just
1193 three intrusions, namely the Bushveld Complex of South Africa, the Stillwater Complex in
1194 Montana, USA, and the Great Dyke of Zimbabwe, but sub-economic deposits that may be mined in
1195 the future occur in many other intrusions. The reefs consist of relatively narrow (<1-2m), but
1196 laterally extensive layers of ultramafic or mafic rocks that typically contain <1-3 % sulfides. The
1197 host intrusions are relatively sulfur poor, and most reefs show mantle-like sulfur isotopic
1198 signatures (Liebenberg, 1970; Li et al., 2008) suggesting that saturation of the magma in sulfide
1199 melt was reached due to fractionation rather than contamination. At least in the case of the
1200 Bushveld Complex, mixing between compositionally different magmas was probably not
1201 instrumental in reef formation because the magmas were highly sulfur undersaturated (Barnes et
1202 al., 2010). The concentration of the sulfides to form the reefs was possibly aided by hydrodynamic
1203 cumulate sorting in response to syn-magmatic subsidence of the intrusions (Maier et al., 2013b).

1204 The main PGE reef of the Wingellina Hills intrusion shows certain similarities to the Great
1205 Dyke and Munni Munni PGE reefs, including the stratiform nature, the stratigraphic position
1206 towards the top of the ultramafic zone of the intrusion (Fig. 25), and the offset stratigraphic
1207 positions of the various chalcophile elements. The bulk PGE content in the Wingellina Hills reef is

1208 in the same range as that in the Main Sulfide Zone of the Great Dyke. In both intrusions, there is
1209 no marked variation in trace element ratios across the reef (unpublished data of Maier). The origin
1210 of the Wingellina Hills main PGE reef can be explained by a model of sulfide saturation from
1211 fractionating NB1-type magma. The intrusion features several additional layers of PGE enrichment
1212 above the main PGE reef, but these have not been studied by us. They could reflect magma
1213 replenishments to the chamber, as the resident magma was likely relatively PGE depleted after
1214 the formation of the main reef.

1215 The low PGE grade of the Wingellina Hills PGE reef relative to the Bushveld and Great Dyke
1216 reefs could reflect less-efficient metal concentration due to faster cooling rates in the relatively
1217 small Wingellina Hills intrusion, whereas the low sulfide contents may reflect metamorphic sulfur
1218 loss, consistent with sub-cotectic sulfide proportions in most Wingellina Hills rocks.

1219

1220 **8.8.2. Cu–Ni–PGE–Au mineralization at Halleys**

1221 Reef-style PGE–Cu–Au mineralization is typical of the upper portions of many layered intrusions
1222 (Maier, 2005). The enrichment of magnetite together with sulfide at the Halleys prospect could
1223 thus suggest that the Halleys intrusive body represents the evolved portion of an adjacent large
1224 layered intrusion, possibly the Blackstone Range of the Mantamaru intrusion, the Cavanagh Range,
1225 or the Saturn intrusion. A petrogenetic link to Blackstone and Cavanagh is presently considered
1226 unlikely as Halleys has much higher gold and sulfur concentrations as well as mica contents than
1227 Blackstone and Cavanagh. Furthermore, field relationships indicate that Halleys crosscuts the
1228 southern segment of the Blackstone Range. A petrogenetic relationship between Halleys and the
1229 Saturn intrusion is more plausible as both share the compositional features of the Alcurra Dolerite
1230 suite. However, a direct connection between Halleys and Saturn is inconsistent with the distinct
1231 magnetic signature of the intrusions.

1232 An alternative model could be that the Halleys mineralization represents contact-style
1233 mineralization at the base or sidewall of a layered intrusion analogous to, e.g., the Platreef of the
1234 Bushveld Complex or the Suhanko deposit of the Portimo Complex, Finland. These deposits are
1235 considered to have formed through sulfide liquid saturation in response to fractional
1236 crystallization accompanied by floor contamination. The proximity of the floor and the resulting
1237 high cooling rate of the magma produced wide, disseminated mineralization rather than narrow
1238 reefs. Sulfides at Halleys have $\delta^{34}\text{S}$ of -0.9 , providing little added constraints on the nature of the
1239 sulfur source. More work is clearly required to further constrain the petrogenesis of the intrusion
1240 and its mineralization.

1241

1242 **8.8.3. Vanadium and PGE mineralization in magnetite seams of the Jameson Range, Mantamaru** 1243 **intrusion**

1244 Advanced fractional crystallization of basaltic magmas leads to cotectic crystallization of magnetite
1245 with silicates, resulting in magnetite-bearing gabbroic and dioritic rocks. The formation of massive
1246 oxide layers requires that magnetite is effectively separated from the silicate minerals as the
1247 cotectic proportions of magnetite and silicates are between 5 and 30% (Toplis and Carroll, 1996).
1248 The mechanism of oxide fractionation has been debated for decades. One of the main problems
1249 has been to explain the knife-sharp contacts of many magnetite layers, requiring extremely
1250 effective separation of oxide crystals from silicate magma. The many structural similarities
1251 between layered cumulates and certain types of sedimentary rocks has led Irvine et al. (1998) to
1252 propose a mechanism of density currents sweeping down along the walls of magma chambers.
1253 Maier et al. (2013b) rejected this model for the Bushveld Complex because density currents would
1254 not preserve the abundant, highly elongated, sub-horizontally oriented anorthosite autoliths
1255 within many oxide seams. The authors instead suggested that oxide-silicate slurries were

1256 mobilized and sorted during subsidence of the Bushveld chamber. This process would be less
1257 turbulent and may preserve some of the compositional layering of the cumulates. The slurries
1258 could be injected into the semi-consolidated crystal pile, and locally form transgressive pipes.

1259 Other models for the formation of the oxide seams proposed in the past include shifts in
1260 phase stability fields of oxides caused by changes in pressure (Cameron, 1980; Lipin, 1993),
1261 temporary supersaturation in magnetite triggered by an increase in the oxygen fugacity of the
1262 magma in response to contamination (Ulmer, 1969), or a combination of magma mixing, pressure
1263 change, and oxidation in response to magma replenishment. However, one of the most popular
1264 models for the formation of massive magnetite layers remains iron oxide liquid immiscibility,
1265 originally advanced by Philpotts (1967). Experiments have produced immiscible iron oxide liquids
1266 in silicate liquids (Freestone, 1978; Roedder, 1978; Naslund, 1983), but whether the silicate melts
1267 used in the experiments are good analogues to natural magmas remains debated. Toplis and
1268 Carrol (1995, 1996) and Tollari et al. (2006, 2008) showed that using silicate melt compositions
1269 close to natural basalts and diorites, magnetite or ilmenite crystallize before the magmas become
1270 saturated with iron oxide liquid.

1271 The potential for apatite deposits in the Giles intrusions remains unknown. Most of the
1272 samples analysed here are apatite free, although Traka Resources (2013) found somewhat
1273 elevated phosphorus concentrations (up to 250 ppm P) in seam 5 of the Jameson intrusion. Anglo
1274 American intersected elevated phosphorus concentrations (up to 8000 ppm P) in magnetite-rich
1275 rocks to the south of the Bell Rock intrusion, suggesting there could be apatite potential in
1276 unexposed magnetite seams. Minor amounts of apatite were also described from Kalka (Goode,
1277 2002). Based on analogy with the Bushveld Complex where nelsonite forms the uppermost of the
1278 magnetite seams, the target horizon for apatite rich layers is at the very top of the intrusions.

1279

1280 **8.8.4. Nebo–Babel Ni–Cu deposit**

1281 Analogous to most other significant Ni-Cu deposits globally, Nebo–Babel is hosted by a tubular
1282 (chonolithic) body interpreted as a magma feeder conduit (Seat et al., 2007, 2009). The chilled
1283 margin contains sulfides, which led Seat et al. (2007) to propose that some of the magmas
1284 entrained sulfide liquid. The authors argued that the concentration of the entrained sulfides was
1285 in part controlled by changes in magma flow velocity, in turn related to changes in the shape of
1286 the conduit. However, in contrast to many other deposits elsewhere that are interpreted to have
1287 formed by addition of external sulfur to the magma, sulfide liquid saturation in the Nebo–Babel
1288 conduit was interpreted to have been triggered by magma mixing (of Alcurra-type magma with
1289 NB1-type magma) and contamination with orthogneiss, i.e. without addition of external sulfur
1290 (Seat et al., 2007; Godel et al., 2011). The orthomagmatic model is feasible in terms of sulfur mass
1291 balance: extracting 100 ppm S from 1 km³ of magma can produce a massive sulfide lens 1 km long,
1292 10 m high, and 20 m wide. However, orthomagmatic derivation of the Nebo Babel sulfides would
1293 require an extremely effective concentration mechanism as the cotectic proportion of sulfide
1294 precipitating from sulfur-saturated troctolitic–gabbro-noritic magma is very small (perhaps as little
1295 as 0.1 wt%).

1296 Seat et al. (2007) based their model on the observation that the sulfides at Nebo Babel have
1297 mantle-like sulfur isotopic compositions, and that the rocks of the Pitjantjatjara Supersuite, which
1298 forms the immediate country rock to the deposit, tend to be sulfur-poor. However, mapping by
1299 GSWA has revealed that the Pitjantjatjara granites are only a very minor lithological component
1300 within this part of the Mamutjarra Zone. More abundant are rocks of the Wirku Metamorphics,
1301 Winburn Granite, and Bentley Supergroup. Amongst these, the Winburn Granite and volcanic and
1302 volcanoclastic rocks of the Bentley Supergroup typically contain visible pyrite and can locally be
1303 sulfide-rich, with up to 3000 ppm S in some samples of ignimbrite and rhyolite (see also Fig. 79 in

1304 Howard et al., 2011, showing sulfide enrichment in drill hole WA02, Strzelecki Metals). The
1305 chamber system required for the Bentley volcanics must have been enormous - and even at the current
1306 level of exposure, Nebo Babel is located only ~ 1km north of unexposed (i.e., interpreted) areas of Bentley
1307 volcanics, and ~ 7 km east of the Winburn granite with is part of the felsic chamber system. It is possible
1308 that at a slightly deeper level, the NB magmas may have intruded into the Bentley volcanic system.

1309 Deposition of the lower portion of the Bentley Supersuite (the Mount Palgrave Group and
1310 much of the Kaarnka Group), pre-dates intrusion of the Nebo–Babel gabbro (Smithies et al., 2013).
1311 The presently available sulfur isotopic data for the units of the Bentley Supergroup indicate a
1312 range of compositions, with those from the lower part having $\delta^{34}\text{S}$ between +1.8 and +7,
1313 potentially representing a suitable external sulfur source for the Nebo–Babel deposit. We thus
1314 argue that addition of external sulfur to the Nebo–Babel magma remains a possibility. The
1315 contamination model would be consistent with the relatively high Cu/Pd and Cu/Ni ratios (possibly
1316 reflecting addition of crustal Cu) and high Au/Pd ratios (possibly due to addition of crustal Au) (see
1317 data of Seat et al., 2007).

1318 The question arises as to why only one major Ni–Cu deposit has so far been found in the
1319 Musgrave Province. Mineral deposits tend to form clusters, suggesting that more Ni-Cu deposits
1320 should occur in the Musgrave Province. Exploration during the last decade has identified several
1321 low-grade deposits (Halley's, Manchego, Succoth) suggesting the true potential of the west
1322 Musgrave Province for Ni–Cu sulfide deposits remains unrealized.

1323

1324 **9. Conclusions**

1325 The Musgrave Province was the focus of long-lived mantle upwelling producing large volumes of
1326 magnesian basaltic to tholeiitic magma and their felsic derivatives. Magmatism led to crustal
1327 melting, lithospheric delamination, and a high crustal heat flux over >200 m.y. The Province

1328 contains one of the greatest concentrations of mafic-ultramafic layered intrusions globally,
1329 amongst them Mantamaru which is one of the world's largest layered intrusions. These data
1330 illustrate that large layered intrusions are not confined to cratons. What is required is a stable
1331 tectonic environment where magmas can ascend in locally extensional, possibly transpressional
1332 zones allowing the formation of thick sill-like bodies.

1333 Due to the large size of the Giles intrusions, cooling rates were relatively slow. This led to
1334 crustal loading, subsidence of magma chambers, and sagging of cumulates prior to complete
1335 solidification. The mobilized cumulates unmixed and formed lenses and layers of peridotite and
1336 magnetite that are locally enriched in PGE. Syn- to post-magmatic tectonism led to
1337 fragmentation of many of the intrusions. The degree of crustal contamination was mostly
1338 relatively minor (<5%), although locally, basaltic magmas mingled with coeval granitic magmas.
1339 The mineralization potential of the Giles intrusions and their host rocks is considerable. Large
1340 magmatic events, particularly those dominated by mafic-ultramafic magmas may cause increased
1341 heat flux into the crust, triggering crustal melting, devolatilization, and large-scale fluid flow.
1342 Deposit types favored by such regimes include magmatic PGE–Cr–V–Fe–P deposits in large layered
1343 intrusions, Ni–Cu sulfide deposits in magma feeder conduits or at the base of layered intrusions,
1344 and hydrothermal deposits of variable style, notably in the roof and sidewalls of the largest
1345 intrusions.

1346

1347 **10. Acknowledgements**

1348 Much of the work documented here arose through a regional-scale geological mapping project
1349 jointly coordinated through the Ngaanyatjarra Council and the Geological Survey of Western
1350 Australia (GSWA), and guided by the local Indigenous people in the region between the

1351 communities of Wingellina and Warburton. The authors thank GSWA for permission to publish the
1352 data. Richard Ernst and Steve Barnes are thanked for their constructive reviews.

1353

1354 **References**

1355 Ahmat A. L. 1986. Petrology, structure, regional geology and age of the gabbroic Windimurra
1356 complex, Western Australia. PhD thesis, University of Western Australia, Perth (unpubl.).

1357 Aitken, AR, Dentith, MC, Evans, SF, Gallardo, LA, Joly, A, Thiel, S, Smithies, RH and Tyler, IM 2013,
1358 Imaging crustal structure in the west Musgrave Province from magnetotelluric and potential field
1359 data: Geological Survey of Western Australia, Report 114, 81p.

1360 Ballhaus, C and Berry, RF 1991, Crystallization pressure and cooling history of the Giles Layered
1361 Igneous Complex, central Australia: *Journal of Petrology*, v. 32, p. 1–28.

1362 Ballhaus, C and Glikson, AY 1989, Magma mixing and intraplutonic quenching in the Wingellina
1363 Hills intrusion, Giles Complex, central Australia: *Journal of Petrology*, v. 30, p. 1443–1469.

1364 Ballhaus, C and Glikson, AY 1995, The petrology of layered mafic-ultramafic intrusions in the Giles
1365 Complex, western Musgrave block, central Australia: *AGSO Journal*, v. 16, p. 69–89.

1366 Barnes, SJ 1993, Partitioning of the PGE and gold between silicate and sulphide magmas in the
1367 Munni Munni Complex, Western Australia: *Geochimica et Cosmochimica Acta*, v. 57, p. 1277–
1368 1290.

1369 Barnes, S-J and Maier, WD 1999, The fractionation of Ni, Cu and the noble metals in silicate and
1370 sulphide liquids, in *Dynamic processes in magmatic ore deposits and their application to mineral*

1371 exploration edited by RR Keays, CM Lesher, PC Lightfoot, and CEG Farrow: Geological Association
1372 of Canada; Short Course Notes 13, p. 69–106.

1373 Barnes, S-J, Maier, WD and Curl, E 2010, Composition of the marginal rocks and sills of the
1374 Rustenburg Layered Suite, Bushveld Complex, South Africa: implications for the formation of the
1375 PGE deposits: *Economic Geology*, v. 105, p.1491–1511.

1376 Becker, H, Horan, MF, Walker, RJ, Gao, S, Lorand J-P and Rudnick, RL 2006, Highly siderophile
1377 element composition of the Earth's primitive mantle: constraints from new data on peridotite
1378 massifs and xenoliths: *Geochimica et Cosmochimica Acta*, v. 70, p. 4528–4550.

1379 Begg, GC, Griffin, WL, Natapov, LM, O'Reilly, SY, Grand, SP, O'Neill, CJ, Hronsky, JMA, Poudjom
1380 Djomani, Y, Swain, CJ, Deen, T and Bowden, P 2009, The lithospheric architecture of Africa: seismic
1381 tomography, mantle petrology, and tectonic evolution: *Geosphere*, v. 5, p. 23–50.

1382 Bodorkos, S, and Wingate, MTD 2008a, 174594: metamorphosed leucogabbro, Mirturtu Camp;
1383 Geochronology Record 716: Geological Survey of Western Australia, 4p.

1384 Bodorkos, S and Wingate, MTD 2008b, 174589: quartz syenite dyke, Amy Giles Hill; Geochronology
1385 Record 715: Geological Survey of Western Australia, 4p.

1386 Bodorkos, S, Wingate, MTD and Kirkland, CL 2008a, 174538: metamonzogranite, Mount Daisy
1387 Bates; Geochronology Record 712: Geological Survey of Western Australia, 4p.

1388 Bodorkos, S, Wingate, MTD and Kirkland, CL 2008b, 174558: metamorphosed quartz diorite,
1389 Mount Fanny; Geochronology Record 713: Geological Survey of Western Australia, 4p.

1390 Bodorkos, S, Wingate, MTD and Kirkland, CL 2008c, 174736: granofelsic metasyenogranite, Mount
1391 Fanny; Geochronology Record 717: Geological Survey of Western Australia, 4p.

- 1392 Bodorkos, S, Wingate, MTD and Kirkland, CL 2008d, 174737: foliated metamonzogranite, Mount
1393 Fanny; Geochronology Record 718: Geological Survey of Western Australia, 5p.
- 1394 Bodorkos, S, Wingate, MTD and Kirkland, CL 2008e, 174747: metagabbro, Mount Fanny;
1395 Geochronology Record 719: Geological Survey of Western Australia, 4p.
- 1396 Boudreau, AE 1999, PELE—a version of the MELTS software programme for the PC platform:
1397 Computers and Geosciences, v. 25, p. 201–203.
- 1398 Cameron, EN 1980, Evolution of the Lower Critical Zone, central sector, eastern Bushveld Complex:
1399 Economic Geology, v. 75, p. 845–871.
- 1400 Campbell, IH and Naldrett, AJ 1979, The influence of silicate : sulfide ratios on the geochemistry of
1401 magmatic sulfides: Economic Geology, v. 74, p. 1503–1505.
- 1402 Cawood, PA and Korsch RJ 2008, Assembling Australia: Proterozoic building of a continent.
1403 Precambrian Research, v 166, p. 1-38.
- 1404 Clarke, GL, Buick, IS, Glikson, AY and Stewart, AJ 1995b, Structural and pressure-temperature
1405 evolution of host rocks of the Giles Complex, western Musgrave Block, central Australia: evidence
1406 for multiple high-pressure events: AGSO Journal of Australian Geology and Geophysics, v. 16, p
1407 127–146.
- 1408 Clarke, GL, Sun, S-S and White, RW 1995a, Grenville age belts and associated older terranes in
1409 Australia and Antarctica: AGSO Journal of Australian Geology and Geophysics, v. 16, p. 25–39.
- 1410 Coleman, P 2009, Intracontinental orogenesis in the heart of Australia: structure, provenance and
1411 tectonic significance of the Bentley Supergroup, western Musgrave Block, Western Australia:
1412 Geological Survey of Western Australia, Record 2009/23, 48p.

1413 Daniels, JL 1974, The Geology of the Blackstone region, Western Australia: Geological Survey of
1414 Western Australia, Bulletin 123, 257p.

1415 Edgoose, CJ, Scrimgeour, IR and Close, DF 2004, Geology of the Musgrave Block, Northern
1416 Territory: Northern Territory Geological Survey, Report 15, 48p.

1417 Eiggins, SM, Woodhead, JD, Kinsley, LPJ, Mortimer, GE, Sylvester, P, McCulloch, MT, Hergt, JM and
1418 Handler, MR 1997, A simple method for the precise determination of >40 trace elements in
1419 geological samples by ICPMS using enriched isotope internal standardisation: Chemical Geology, v.
1420 134, p. 311–326.

1421 Evins, PM, Smithies, RH, Howard, HM, Kirkland, CL, Wingate, MTD and Bodorkos, S 2010a,
1422 Redefining the Giles Event within the setting of the 1120–1020 Ma Ngaanyatjarra Rift, west
1423 Musgrave Province, Central Australia: Geological Survey of Western Australia, Record 2010/6, 36p.

1424 Evins, PM, Smithies, RH, Howard, HM, Kirkland, CL, Wingate, MTD and Bodorkos, S 2010b, Devil in
1425 the detail: the 1150–1000 Ma magmatic and structural evolution of the Ngaanyatjarra Rift, west
1426 Musgrave Province, Central Australia: Precambrian Research, v. 183, p. 572–588.

1427 Evins, PM, Smithies, RH, Howard, HM and Maier, WD 2009, Holt, WA Sheet 4546: Geological
1428 Survey of Western Australia, 1:100 000 Geological Series.

1429 Foulger, GR 2010, Plates vs plumes: Wiley-Blackwell, Chichester, 328p.

1430 Freestone, IC 1978, Liquid immiscibility in alkali-rich magmas: Chemical Geology, v. 23, p. 116–123.

1431 Gale A, Dalton CA, Langmuir CH, Su Y, Schilling J-G (2013) The mean composition of ocean ridge
1432 basalts. *Geochemistry, geophysics, geosystems*, Volume 14, Number 3,
1433 doi:10.1029/2012GC004334

1434

1435 Glikson, AY (editor) 1995, The Giles mafic-ultramafic complex and environs, western Musgrave
1436 Block, central Australia: Thematic issue: AGSO Journal of Geology and Geophysics, v. 16, no. 1–2,
1437 193p.

1438 Glikson, AY, Stewart, AT, Ballhaus, GL, Clarke, GL, Feeken, EHT, Level, JH, Sheraton, JW and Sun, S-
1439 S 1996, Geology of the western Musgrave Block, central Australia, with reference to the mafic-
1440 ultramafic Giles Complex: Australian Geological Survey Organisation, Bulletin 239, 206p.

1441 Godel, B, Seat, Z, Maier, WD and Barnes, S-J 2011, The Nebo–Babel Ni-Cu-PGE sulfide deposit
1442 (West Musgrave Block, Australia): part 2 — constraints on parental magma and processes, with
1443 implications for mineral exploration: *Economic Geology*, v. 106, p. 557–584.

1444 Goode, ADT 1970, The petrology and structure of the Kalka and Ewarara layered basic intrusions,
1445 Giles Complex, central Australia: The University of Adelaide, Adelaide, PhD thesis (unpublished).

1446 Goode, ADT 1976a, Small scale primary cumulus igneous layering in the Kalka layered intrusion,
1447 Giles Complex, central Australia: *Journal of Petrology*, v. 17, p. 379–397.

1448 Goode, ADT 1976b, Sedimentary structures and magma current velocities in the Kalka layered
1449 intrusion, central Australia: *Journal of Petrology*, v. 17, p. 546–558.

1450 Goode, ADT 1977a, Vertical igneous layering in the Ewarara layered intrusion, central Australia:
1451 *Geological Magazine*, v. 114, p. 215–218.

1452 Goode, ADT 1977b, Flotation and remelting of plagioclase in the Kalka Intrusion, central Australia:
1453 petrological implications for anorthosite genesis: *Earth and Planetary Science Letters*, v. 34, p.
1454 375–380.

1455 Goode, ADT 1977c, Intercumulus igneous layering in the Kalka layered intrusion, central Australia:
1456 Geological Magazine, v. 114, p. 215–218.

1457 Goode, ADT 1978, High temperature, high strain rate deformation in the lower crustal Kalka
1458 Intrusion, central Australia: Contributions to Mineralogy and Petrology, v. 66, p. 137–148.

1459 Goode, ADT 2002, The Western Musgrave Block – Australia: Data Metallogenica, District Overview.
1460 42p.

1461 Goode, ADT and Krieg, GW 1967, The geology of Ewarara Intrusion, Giles Complex, central
1462 Australia: Journal of the Geological Society of Australia, v. 14, p. 185–194.

1463 Goode, ADT and Moore, AC 1975, High pressure crystallisation of the Ewarara, Kalka and Gosse
1464 Pile intrusions, Giles Complex, central Australia: Contributions to Mineralogy and Petrology, v. 51,
1465 p. 77–97.

1466 Gray, CM 1971, Strontium isotope studies on granulites: The Australian National University,
1467 Canberra, PhD thesis (unpublished), 242p.

1468 Gray, CM, Cliff, RA and Goode, ADT, 1981, Neodymium–strontium isotopic evidence for extreme
1469 contamination in a layered basic intrusion: Earth and Planetary Science Letters, v. 56, p. 189–198.

1470 Gray, CM and Goode, ADT 1989, The Kalka layered intrusion, central Australia: a strontium isotopic
1471 history of contamination and magma dynamics: Contributions to Mineralogy and Petrology, v. 103,
1472 p. 35–43.

1473 Howard, HM, Smithies, RH, Kirkland, CL, Evins, PM and Wingate, MTD 2009, Age and geochemistry
1474 of the Alcurra Suite in the west Musgrave Province and implications for orthomagmatic Ni-Cu-PGE

1475 mineralization during the Giles Event: Geological Survey of Western Australia, Record 2009/16,
1476 16pp.

1477 Howard, HM, Smithies, RH and Pirajno F, 2006b, Geochemical and Nd isotopic signatures of mafic
1478 dykes in the western Musgrave Complex: Geological Survey of Western Australia Annual Review
1479 2005–06, p. 64–71.

1480 Howard, HM, Smithies, RH, Pirajno, F and Skwarnecki, MS 2006a, Bates, WA Sheet 4646:
1481 Geological Survey of Western Australia, 1:100 000 Geological Series.

1482 Howard, HM, Smithies, RH, Pirajno, F and Skwarnecki, MS 2007, Bell Rock, WA Sheet 4645:
1483 Geological Survey of Western Australia, 1:100 000 Geological Series.

1484 Howard, HM, Smithies, RH, Werner, M, Kirkland, CL and Wingate, MTD 2011a, Geochemical
1485 characteristics of the Alcurra Dolerite (Giles Event) and its extrusive equivalents in the Bentley
1486 Supergroup: Geological Survey of Western Australia, Record 2011/2, p. 27–30.

1487 Howard, HM, Werner, M, Smithies, RH, Kirkland, CL, Kelsey, DL, Hand, M, Collins, A, Pirajno, F,
1488 Wingate, MTD, Maier, WD and Raimondo, T 2011b, The geology of the west Musgrave Province
1489 and the Bentley Supergroup — a field guide: Geological Survey of Western Australia, Record
1490 2011/4, 119p.

1491 Howard HM, Smithies RH, Kirkland CL, Kelsey DE, Aitken A, Wingate MTD, Quentin de
1492 Gromard R, Spaggiari CV, Maier, WD 2015, The burning heart — The Proterozoic geology and
1493 geological evolution of the west Musgrave Region, central Australia, Gondwana Research, 27, 64-
1494 94

1495 Irvine, TN 1970, Crystallization sequences in magmas of the Muskox intrusion and some other
1496 layered intrusions: Geological Society of South Africa, Special Publication 1, p. 441–476.

1497 Irvine, TN, Andersen, JCO and Brooks, CK 1998, Included blocks (and blocks within blocks) in the
1498 Skaergaard Intrusion: Geological relations and the origins of rhythmic modally graded layers:
1499 Geological Society of America, Bulletin 110, p. 1398–1447.

1500 Kelsey, DE, Hand, M, Evins, P, Clark, C and Smithies, H 2009, High temperature, high geothermal
1501 gradient metamorphism in the Musgrave Province, central Australia: potential constraints on
1502 tectonic setting, in Biennial conference of the Specialist Group for Geochemistry, Mineralogy and
1503 Petrology, Kangaroo Island November 2009 edited by NE Timms, J Foden, K Evans and C Clark:
1504 Geological Society of Australia Abstracts No. 96, p. 28.

1505 Kelsey, DE, Hand, M, Smithies, H, Evins, P, Clark, C and Kirkland, CL 2010, What is the tectonic
1506 setting of long-lived Grenvillian-aged ultrahigh temperature, high geothermal gradient
1507 metamorphism in the Musgrave Province, central Australia?: Geological Society of America,
1508 Abstracts with Programs, 42(5), p. 516.

1509 King, RJ 2008, Using calculated pseudosections in the system NCKFMASHTO and SHRIMP II U–Pb
1510 zircon dating to constrain the metamorphic evolution of paragneisses in the Latitude Hills, West
1511 Musgrave Province, Western Australia: Geological Survey of Western Australia, Record 2009/15,
1512 67p.

1513 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008a, 183496: orthogneiss, Mount West;
1514 Geochronology Record 747: Geological Survey of Western Australia, 5p.

1515 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008b, 183459: charnockite, Latitude Hill,
1516 Geochronology Record 722: Geological Survey of Western Australia, 5p.

1517 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008c, 183509: leucogranite dyke, Mount West;
1518 Geochronology Record 724: Geological Survey of Western Australia, 4p.

1519 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008d, 193850: leucogranite dyke, Mount Fanny;
1520 Geochronology Record 748: Geological Survey of Western Australia, 4p.

1521 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008e, 174761: porphyritic granite dyke, Bell Rock;
1522 Geochronology Record 721: Geological Survey of Western Australia, 4p.

1523 Kirkland, CL, Wingate, MTD and Bodorkos, S 2008f, 185509: leucogranite, Mount Aloysius;
1524 Geochronology Record 725: Geological Survey of Western Australia, 4p.

1525 Kirkland, CL, Wingate, MTD and Smithies, RH 2011, 194762: leucogabbro, Mount Finlayson;
1526 Geochronology Record 966: Geological Survey of Western Australia, 4p.

1527 Kirkland, CL, Smithies, RH, Woodhouse, AJ, Howard, HM, Wingate, MTD, Belousova, EA, Cliff, JB,
1528 Murphy, RC and Spaggiari, CV 2013, Constraints and deception in the isotopic record; the crustal
1529 evolution of the west Musgrave Province, central Australia: Gondwana Research, v. 23, p. 759–
1530 781.

1531 Latypov R, Hanski E, Lavrenchuk A, Huhma H, and Havela T 2011, A ‘Three-Increase Model’ for the
1532 Origin of the Marginal Reversal of the Koitelainen Layered Intrusion, Finland. *Journal of Petrology*,
1533 v. 52, p. 733-764.

1534 Leshner, CM and Burnham, OM 2001, Multicomponent elemental and isotopic mixing in Ni–Cu–
1535 (PGE) ores at Kambalda, Western Australia: *Canadian Mineralogist*, v. 39, p. 421–446.

1536 Li, ZX 2000, Palaeomagnetic evidence for unification of the North and West Australian craton by
1537 ca. 1.7 Ga: new results from the Kimberley Basin of northwestern Australia: *Geophysical Journal
1538 International*, v. 142, p. 173-180.

1539 Li, C and Ripley, EM 2010, The relative effects of composition and temperature on olivine-liquid
1540 partitioning: statistical deconvolution and implications for petrologic modelling: *Chemical Geology*,
1541 v. 275, p. 95–104.

1542 Li, C, Ripley, EM, Oberthür, T, Miller, JD Jr and Joslin, GD 2008, Textural, mineralogical and stable
1543 isotope studies of hydrothermal alteration in the main sulfide zone of the Great Dyke, Zimbabwe
1544 and the precious metals zone of the Sonju Lake Intrusion, Minnesota, USA: *Mineralium Deposita*,
1545 v. 43, p. 97–110.

1546 Liebenberg, L 1970, The sulfides in the layered sequence of the Bushveld igneous Complex:
1547 *Geological Society of South Africa, Special Publication*, v. 1, p. 108–208.

1548 Lightfoot, PC and Naldrett, AJ 1983, The geology of the Tabankulu section of the Insizwa Complex,
1549 Transkei, Southern Africa, with reference to the nickel sulphide potential: *Transactions of the*
1550 *Geological Society of South Africa*, v. 86, p. 169–187.

1551 Lipin, BR 1993, Pressure increases, the formation of chromite seams, and the development of the
1552 Ultramafic series in the Stillwater Complex, Montana: *Journal of Petrology*, v. 34, p. 955–976.

1553 Maier, WD 2005, Platinum-group element (PGE) deposits and occurrences: mineralization styles,
1554 genetic concepts and exploration criteria: *Journal of African Earth Sciences*, v. 41, p. 165–191.

1555 Maier, WD, Barnes, SJ, Gartz, V and Andrews, G 2003b, Pt–Pd reefs in magnetitites of the Stella
1556 layered intrusion, South Africa: a world of new exploration opportunities for platinum group
1557 elements: *Geology*, v. 31, p. 885–888.

1558 Maier, WD, Barnes, S-J and Groves, DI 2013b, The Bushveld Complex, South Africa: Formation of
1559 platinum–palladium, chrome and vanadium- rich layers via hydrodynamic sorting of a mobilized

1560 cumulate slurry in a large, relatively slowly cooling, subsiding magma chamber: *Mineralium*
1561 *Deposita*, v. 48, p. 1–56.

1562 Maier, WD and Eales, HV 1997, Correlation within the UG2 – Merensky Reef interval of the
1563 Western Bushveld Complex, based on geochemical, mineralogical and petrological data.
1564 Geological Survey of South Africa, Bulletin 120, 56p.

1565 Maier, WD, Rasmussen, B, Li, C, Barnes, S-J and Huhma, H 2013a, The Kunene anorthosite
1566 complex, Namibia, and its satellite bodies: geochemistry, geochronology and economic potential:
1567 *Economic Geology*, v. 108, p. 953–986.

1568 Metals X Ltd 2013, Globally significant nickel project, viewed 8 April 2014,
1569 <http://www.metalsx.com.au/system/assets/26/original/Nickel_Division.pdf>.

1570 Morris, PA and Pirajno, F 2005, Geology, geochemistry, and mineralization potential of
1571 Mesoproterozoic sill complexes of the Bangemall Supergroup, Western Australia: Geological
1572 Survey of Western Australia of Western Australia, Report 99, 72p.

1573 Morse, SA 1981, Kiglapait geochemistry IV: the major elements: *Geochimica et Cosmochimica*
1574 *Acta*, v. 45, p. 461–479.

1575 Naldrett, AJ and Lehmann, J 1988, Spinel nonstoichiometry as the explanation for Ni-, Cu-, and
1576 PGE-enriched sulphides in chromitites, in *Geo-Platinum 87* edited by HM Prichard et al.: Elsevier,
1577 Barking, p. 93–109.

1578 Naslund, HR 1983, The effect of oxygen fugacity on liquid immiscibility in iron-bearing silicate
1579 melts: *American Journal of Science*, v. 283, p. 1034–1059.

1580 Nesbitt, RW, Goode, ADT, Moore, AC and Hopwood, TP 1970, The Giles Complex, central Australia:
1581 a stratified sequence of mafic and ultramafic intrusions: Special Publications of the Geological
1582 Society of South Africa, v. 1, p. 547–564.

1583 Nesbitt, RW and Talbot, JL 1966, The layered basic and ultrabasic intrusives of the Giles Complex,
1584 central Australia: Contributions to Mineralogy and Petrology, v. 13, p. 1–11.

1585 Norrish, K and Chappell, BW 1977, X-ray fluorescence spectrometry, in Physical Methods in
1586 Determinative Mineralogy, 2nd Edition edited by J Zussman: Academic Press, London, p. 201–272.

1587 Pascoe, A 2012, The geochemistry and petrogenesis of the mafic and ultramafic giles intrusions at
1588 Latitude Hill, west Musgrave Province, Western Australia: The University of Tasmania, Hobart, BSc
1589 Honours thesis (unpublished).

1590 Patino Douce, A.E. & Beard, J.S. (1995). Dehydration-melting of Biotite Gneiss and Quartz
1591 Amphibolite from 3 to 15 kbar. Journal of Petrology v. 36, 707-738.

1592 Philpotts, AR 1967, Origin of certain iron–titanium oxide and apatite rocks: Economic Geology, v.
1593 62, p. 303–315.

1594 Phosphate Australia Ltd 2014, Manhego Prospect: Musgrave Project, Western Australia, viewed 6
1595 May 2014, at <<http://www.asx.com.au/asxpdf/20140103/pdf/42lz643zrngqwx.pdf>>.

1596 Raedeke, LD 1982, Petrogenesis of the Stillwater Complex: The University of Washington, Seattle,
1597 PhD thesis (unpublished).

1598 Redstone Resources Ltd 2008a, Quarterly report for the period ending June 30th 2008, viewed 6
1599 May 2014, at
1600 <http://www.redstone.com.au/investor_centre_asx_announcements_files/2008_07_31_01.pdf>.

1601 Redstone Resources Ltd 2008b, Report of the Annual General Meeting, viewed 8 April 2014, at
1602 <<http://www.asx.com.au/asxpdf/20081127/pdf/31dvzvjbkp7f5b.pdf>>.

1603 Reynolds, IM 1985, The nature and origin of titaniferous magnetite-rich layers in the Upper Zone
1604 of the Bushveld Complex: a review and synthesis: *Economic Geology*, v. 80, p. 1089–1108.

1605 Rollinson HR (2013) *Using geochemical data: evaluation, presentation, interpretation*, Routledge,
1606 Oxon, UK.

1607 Scrimgeour, IR and Close, DF 1999, Regional high pressure metamorphism during intracratonic
1608 deformation: the Petermann orogeny, central Australia: *Journal of Metamorphic Geology*, v. 17, p.
1609 557–572.

1610 Seat, Z 2008, *Geology, petrology, mineral and whole-rock chemistry, stable and radiogenic isotope*
1611 *systematics and Ni–Cu–PGE mineralisation of the Nebo–Babel intrusion, west Musgrave, Western*
1612 *Australia: The University of Western Australia, Perth, PhD thesis (unpublished).*

1613 Seat, Z, Beresford, SW, Grguric, BA, Gee, MA and Grassineau, NV 2009, Reevaluation of the role of
1614 external sulfur addition in the genesis of Ni–Cu–PGE deposits: evidence from the Nebo–Babel Ni–
1615 Cu–PGE deposit: West Musgrave, Western Australia: *Economic Geology*, v. 104, p. 521–538.

1616 Seat, Z, Beresford, SW, Grguric, BA, Waugh, RS, Hronsky, JMA, Gee, MMA, Groves, DI and
1617 Mathison, CI 2007, Architecture and emplacement of the Nebo–Babel gabbro-norite-hosted
1618 magmatic Ni–Cu–PGE sulfide deposit, West Musgrave, Western Australia: *Mineralium Deposita*, v.
1619 42, p. 551–582.

1620 Sharpe, MR 1981, The chronology of magma influxes to the eastern compartment of the Bushveld
1621 Complex, as exemplified by its marginal border group: *Journal of the Geological Society of London*,
1622 v. 138, p. 307–326.

1623 Sheraton, JW and Sun, S-s 1995, Geochemistry and origin of felsic igneous rocks of the western
1624 Musgrave Block: AGSO Journal, v. 16, p. 107–125.

1625 Silver, PG, Behn, MD, Kelley, K, Schmitz, M and Savage, B 2006, Understanding cratonic flood
1626 basalts: Earth and Planetary Science Letters, v. 245, p. 190–201.

1627 Smithies, RH, Howard, HM, Evins, PM, Kirkland, CL, Bodorkos, S and Wingate, MTD 2009, The west
1628 Musgrave Complex — some new geological insights from recent mapping, geochronology, and
1629 geochemical studies: Geological Survey of Western Australia, Record 2008/19, 20p.

1630 Smithies, RH, Howard, HM, Evins, PM, Kirkland, CL, Kelsey, DE, Hand, M, Wingate, MTD, Collins,
1631 AS, Belousova, E and Allchurch, S 2010, Geochemistry, geochronology and petrogenesis of
1632 Mesoproterozoic felsic rocks in the western Musgrave Province of central Australia, and
1633 implication for the Mesoproterozoic tectonic evolution of the region: Geological Survey of
1634 Western Australia, Report 106, 73p.

1635 Smithies, RH, Howard, HM, Evins, PM, Kirkland, CL, Kelsey, DE, Hand, M, Wingate, MTD, Collins, AS
1636 and Belousova, E 2011, Mesoproterozoic high temperature granite magmatism, crust–mantle
1637 interaction and the intracontinental evolution of the Musgrave Province: Journal of Petrology,
1638 doi:10.1093/petrology/egr010.

1639 Smithies, RH, Howard, HM, Kirkland, CL, Werner, M, Medlin, CC, Wingate, MTD and Cliff, JB 2013,
1640 Geochemical evolution of rhyolites of the Talbot Sub-basin and associated felsic units of the
1641 Warakurna Supersuite: Geological Survey of Western Australia, Report 118, 74p.

1642 Smithies, RH, Kirkland, CL, Korhonen, FJ, Aitken, ARA, Howard, HM, Maier, WD, Wingate, MTD,
1643 Quentin de Gromard, R and Gessner, K 2015, The Mesoproterozoic thermal evolution of the

1644 Musgrave Province in central Australia – Plume vs. the geological record: *Gondwana Research*, v.
1645 27, p. 64-94

1646 Smithies, RH, Howard, HM, Kirkland, CL, Korhonen, FJ, Medlin, CC, Maier, WD, Quentin de
1647 Gromard, R, and Wingate, MTD, in press. Piggyback Supervolcanoes - Long-Lived, Voluminous,
1648 Juvenile Rhyolite Volcanism in Mesoproterozoic Central Australia: *Journal of Petrology*, in press.

1649 Sobolev, SV, Sobolev, AV, Kuzmin, DV, Krivolutsкая, NA, Petrunin, AG, Arndt, NT, Radko, VA and
1650 Vasiliev, YR 2011, Linking mantle plumes, large igneous provinces and environmental
1651 catastrophes: *Nature*, v. 477, p. 312–316.

1652 Sprigg, RC and Wilson, RB 1959, The Musgrave mountain belt in South Australia: *Geologische*
1653 *Rundschau*, v.c47, p. 531–542.

1654 Staubmann, M 2010, The petrogenesis and economic potential of the Southern Cavenagh Range
1655 Intrusion (West Musgraves), Western Australia: The University of Tasmania, Hobart, BSc Honours
1656 thesis (unpublished).

1657 Stewart, AJ 1995, Resolution of conflicting structures and deformation history of the Mount
1658 Aloysius granulite massif, western Musgrave Block, central Australia: *AGSO Journal of Australian*
1659 *Geology and Geophysics*, v. 16, p. 91–105.

1660 Sun, S.-s, and McDonough, WF 1989, Chemical and isotopic systematics of oceanic basalts:
1661 implications for mantle composition and processes, in *Magmatism in the Ocean Basins* edited by
1662 AD Saunders and MJ Norry: *Geological Society Special Publication*, 42, p. 313–345.

1663 Sun, S-S, Sheraton, JW, Glikson, AY and Stewart, AJ 1996, A major magmatic event during 1050–
1664 1080 Ma in central Australia, and an emplacement age for the Giles Complex: *AGSO Journal of*
1665 *Australian Geology and Geophysics*, 24, p. 13–15.

1666 Tegner, C, Wilson, JR and Brooks, CK 1993, Intraplutonic quench zones in the Kap Edvard Holm
1667 layered gabbro complex, east Greenland: *Journal of Petrology*, v. 34, p. 681–710.

1668 Teigler, B and Eales, HV 1996, The Lower and Critical Zones of the western limb of the Bushveld
1669 Complex, as indicated by the Nooitgedacht boreholes: Geological Survey of South Africa, *Bulletin*
1670 111, 126p.

1671 Tollari, N, Baker, D and Barnes, S-J 2008, Experimental effects of pressure and fluorine on apatite
1672 saturation in mafic magmas, with reference to layered intrusions and massif anorthosites:
1673 *Contributions to Mineralogy and Petrology*, v. 156, p. 161–175.

1674 Tollari, N, Toplis, MJ and Barnes, S-J 2006, Predicting phosphate saturation in silicate magmas: an
1675 experimental study of the effects of melt composition and temperature: *Geochimica et*
1676 *Cosmochimica Acta*, v. 70, p. 1518–1536.

1677 Toplis, MJ and Carroll, MR 1995, An experimental-study of the influence of oxygen fugacity on Fe–
1678 Ti oxide stability, phase-relations, and mineral–melt equilibria in ferro-basaltic systems: *Journal of*
1679 *Petrology*, v. 36, p. 1137–1170.

1680 Toplis, MJ and Carroll, MR 1996, Differentiation of ferro-basaltic magmas under conditions open
1681 and closed to oxygen: implications for the Skaergaard intrusion and other natural systems: *Journal*
1682 *of Petrology*, v. 37, p. 837–858.

1683 Traka Resources Ltd 2011, Quarterly Activities Report for the three months ended 30 June 2011.

1684 Traka Resources Ltd 2013, Musgrave Project, viewed 29 August 2013, at
1685 <<http://www.trakaresources.com.au>>.

1686 Ulmer, GC 1969, Experimental investigation of chromite spinels: *Economic Geology*, Monograph 4,
1687 p 114–131.

1688 Wade, BP 2006, Unravelling the tectonic framework of the Musgrave Province, central Australia:
1689 The University of Adelaide, Adelaide, PhD thesis (unpublished).

1690 Waight, TE, Maas, R and Nicholls, IA 2000, Fingerprinting feldspar phenocrysts using crystal
1691 isotopic composition stratigraphy: implications for crystal transfer and magma mingling in S-type
1692 granites: *Contributions to Mineralogy and Petrology*, v. 139, p. 227–239.

1693 White, RW, Clarke, GL and Nelson, DR 1999, SHRIMP U–Pb zircon dating of Grenville-age events in
1694 the western part of the Musgrave Block, central Australia: *Journal of Metamorphic Geology*, v. 17,
1695 p. 465–481.

1696 Wilson, AH, Naldrett, AJ and Tredoux, M 1989, Distribution and controls of platinum group
1697 element and base metal mineralization in the Darwendale subchamber of the Great Dyke,
1698 Zimbabwe: *Geology*, v. 17, p. 649–652.

1699 Wingate, MTD and Evans, DAD 2003, Palaeomagnetic constraints on the Proterozoic tectonic
1700 evolution of Australia: Yoshida, M, Windley, BF and Dasgupta, S (eds), *Proterozoic East Gondwana:
1701 Supercontinent Assembly and Breakup*. Geological Society, London, Special Publications, v. 206, p.
1702 77-91.

1703 Wingate, MTD, Pirajno, F and Morris, PA 2004, Warakurna large igneous province: a new
1704 Mesoproterozoic large igneous province in west-central Australia: *Geology*, v. 32, p. 105–108.


1705 Yang, S, Maier, WD, Hanski, E, Lappalainen, M, Santaguida, F and Määttä, S 2013a, Origin of ultra-
1706 nickeliferous olivine in the Kevitsa Ni–Cu–PGE mineralized intrusion, Lapland, Finland:
1707 *Contributions to Mineralogy and Petrology*, v. 166, p. 81–95.

1708 Yang S, Maier WD, Lahaye Y, O'Brien, H 2013b, Strontium isotope disequilibrium of plagioclase in
 1709 the Upper Critical Zone of the Bushveld Complex: evidence for mixing of crystal slurries, Cont Min
 1710 Petrol DOI 10.1007/s00410-013-0903-4

1711 Zhao, J-X, McCulloch, MT and Korsch, RJ 1994, Characterisation of a plume-related ~800 Ma
 1712 magmatic event and its implications for basin formation in central-southern Australia: Earth and
 1713 Planetary Science Letters, v. 121, p. 349–367.

1714

1715 **Figure captions**




1716


1717 Figure 1. Simplified geological map of the Musgrave Province, with mafic-ultramafic intrusions

1718 highlighted in bottom panel. From Maier et al. (2014).

1719




1720 Figure 2. Interpreted bedrock geology map of the west Musgrave Province. Location of Nebo-
1721 Babel deposits is indicated by black star. From Maier et al. (2014).




1722

1723 Figure 3. Interpreted bedrock geological map of the Pirntirri Mulari intrusion, showing sample
 1724 localities (yellow circles). Stippled line indicates position of postulated reef horizon. Light shaded
 1725 areas indicate regions of regolith cover. Modified from Maier et al. (2014).




1726

1727 Figure 4. Textures of rocks in the Pirntirri Mulari intrusion. a) Centimetre-scale interlayering of
 1728 pyroxenite and gabbronorite showing sharp bottom contact and upward grading; note small
 1729 lenses and schlieren of pyroxenite within gabbronorite (near GSWA 189359); b) pegmatoidal layer
 1730 within medium-grained pyroxenite (GSWA 189360); c) contact between pyroxenite and overlying
 1731 gabbronorite; finger-like structure of pyroxenite is interpreted as injection of pyroxenite mush into
 1732 gabbronorite (near GSWA 189374); d) schlieren of pyroxenite within leucogabbronorite (near
 1733 GSWA 189358). Adapted from Maier et al. (2014).




1734

1735 Figure 5. Compositional variation with stratigraphic height in the Pirntirri Mulari intrusion. Red
 1736 diamonds indicate the platinum group element – rich sample analysed by Redstone Resources Ltd,
 1737 and horizontal shaded bar indicates postulated position of platinum group element reef. Range of
 1738 primitive mantle composition (for Cu/Pd) is based on Barnes and Maier (1999) and Becker et al.
 1739 (2006). Figure from Maier et al. (2014).



1740


1741 Figure 6. Simplified geological map of Wingellina Hills intrusion, showing location of boreholes
 1742 where platinum group element mineralization has been intersected. Figure provided by Metals X
 1743 Ltd, with permission. Figure from Maier et al. (2014).



1744


1745 Figure 7. Log of percussion drillhole WPRC0-064, Wingellina Hills intrusion (Web = websterite,

1746 Wehr = wehlite). From Maier et al. (2014).



1747

1748 Figure 8. a,b) Examples of mingling textures between G2 gabbro and granite in the West Hinckley
1749 Range. Figure from Maier et al. (2014).




1750

1751 Figure 9. Field photographs of samples from the Lehman Hills, Jameson and Blackstone intrusions:
1752 a) banded horizon containing schlieren, lenses and fragments of fine-grained gabbronorite, and
1753 medium- to coarse-grained pyroxenite (Lehman Hills, near GSWA 189310); c) basal magnetite
1754 layer, Jameson Range; note shallow dip of layer to the right (locality GSWA 194642); e) modally
1755 graded layering in olivine gabbronorite at Jameson Range (MGA 363526E 7149428N). f) steeply
1756 south-dipping magnetite layer, southern edge of Blackstone Range (GSWA 194679). Adapted from
1757 Maier et al. (2014).

1758


1759

1760




1761

1762 Figure 10. Aeromagnetic total magnetic intensity (TMI) image of the area to the northwest of
 1763 Jameson, showing interpreted trend of magnetite layers (aeromagnetic highs) within the Jameson
 1764 intrusion. Yellow circles are GSWA sample sites. Figure from Maier et al. (2014).




1765

1766 Figure 11. Composition of basal magnetite layer within the Jameson intrusion, based on 32
 1767 samples collected along strike by Traka Resources (Traka Resources Ltd, 2011, written comm., 21
 1768 October): a) Pt+Pd vs V; b) Pt+Pd vs S. Adapted from Maier et al. (2014).




1769

1770 Figure 12. Metal patterns of the basal magnetite layer of the Jameson intrusion (shaded field),
 1771 compared to basal magnetite layer in Upper Zone of Bushveld Complex (blue line) and PGE-rich
 1772 magnetite layers of Stella intrusion, South Africa (red lines). Adapted from Maier et al. (2014).




1773

1774 Figure 13. Aeromagnetic image of the Saturn intrusion, between the Cavenagh intrusion (lower
 1775 left) and the Blackstone intrusion (to the north of the image). Red circles are sampling points for
 1776 various transverses (named and enclosed in ellipses). Note the concentric pattern defining the
 1777 Saturn intrusion. Figure from Maier et al. (2014).



1778

1779 Figure 14. Compositional traverse (west to east) across the Saturn intrusion at Phoebe (see Fig. 27
 1780 for sample localities). Note increase in PGE and Cu concentrations approximately halfway along
 1781 the traverse. Plots of: a) TiO₂; b) Pt+Pd; c) Cu vs Eastings. Adapted from Maier et al. (2014).



1782

1783 Figure 15. Binary variation diagrams vs MgO of selected major elements in the Giles intrusions: a)

1784 Al₂O₃; b) TiO₂; c) K₂O; d) P₂O₅. Coloured fields in (a) denote cumulates whose compositions are


1785 principally controlled by variation in modal proportions of plagioclase and olivine (blue) and

1786 plagioclase+pyroxene (grey). Yellow field indicates rocks that contain significant magnetite.


1787 Vectors in b)–d) indicate that some cumulates contain substantial magnetite, apatite, and granite

1788 components. 'Northern' intrusions include intrusive fragments to the north of Mt Muir and

1789 Hinckley Range. Figure from Maier et al. (2014).




1791 Figure 16. Rocks of the Giles intrusions plotted into binary variation diagrams vs MgO of: a) Cr; and
 1792 b) Ni. Approximate compositions of selected silicate and oxide minerals are shown in blue
 1793 lettering. Mineral abbreviations as for Figure 19. Figure from Maier et al. (2014).



1794


1795 Figure 17. Binary variation diagram of Cr/V vs Mg# for the Giles intrusions. Bushveld data are from
 1796 Maier et al. (2013b). 'Northern' intrusions include intrusive fragments North of Mt Muir and the
 1797 Hinckley Range. Cavenagh data include samples from Staubmann (2010). Figure from Maier et al.
 1798 (2014).

1799




1800

1801 Figure 18. Primitive mantle-normalized multi-element variation diagrams for rocks of the Giles
 1802 Event that may be liquids, including: a) unevolved samples of the Alcurra dolerite suite; b)
 1803 unevolved samples of the G2 gabbros; c) G1 gabbros enriched in incompatible trace elements; and
 1804 d) Cavenagh microgabbros. Normalization factors are from Sun and McDonough (1989). Adapted
 1805 from Maier et al. (2014).




1807 Figure 19. Plot of ϵ_{Nd} vs Ce/Nb for the Giles intrusions. Note that troctolitic G1 intrusions and the
 1808 Alcurra Dolerite suite plot near the mantle range, whereas the other intrusions contain an
 1809 enriched component. The compositional field of Pitjantjatjara granite contains the 10th–90th
 1810 percentile of Ce/Nb data. Data for Kalka intrusion are from Wade (2006). Figure from Maier et al.
 1811 (2014).



1812

1813 Figure 20. Binary variation diagrams vs Mg# of: a) S; b) Cu; c) Pt+Pd; d) Cu/Pd. Primitive mantle

1814 value in d) is from Barnes and Maier (1999). Adapted from Maier et al. (2014).



1815

1816 Figure 21. Plot of Ni vs Fo in olivine: a) data from the west Musgrave Province; b) global data of
 1817 layered intrusions (data compiled from Teigler and Eales, 1996; Maier and Eales, 1997; Lightfoot et
 1818 al., 1984; Raedeke, 1982; E Hanski, unpublished data). Grey shading is background data, from
 1819 Sobolev et al. (2011). Figure from Maier et al. (2014).


1820

1821

1822


1823

1824




1825

1826 Figure 22. Binary variation diagrams of La/Sm vs La/Nb: Most of the troctolitic intrusions
 1827 (Mantamaru), the Alcurra Dolerite suite, and the Halleys and Saturn intrusions have primitive
 1828 mantle-like trace element ratios, whereas many of the other intrusions (notably Cavenagh,
 1829 Hinckley Range, Murray Range, and the ultramafic intrusions) contain a crustal component,
 1830 possibly of Pitjantatjara granite (Pit granite). Solid line represents mixing line between picrite (with
 1831 trace element contents assumed to be 4x primitive mantle, i.e. equivalent to ~25% partial mantle
 1832 melting) and a contaminated magma produced by AFC ($r=0.8$, $f=0.8$) of picrite with a 17% partial
 1833 melt of Pitjantjarra granite (calculated by assuming modal proportions determined during
 1834 experimental melting of biotite gneiss at 875°, 3kbar, Patino Douce and Beard, 1995, and D values
 1835 summarized in Rollinson, 2013).



1836


1837 Figure 23. Stratigraphic comparison of Giles intrusions with Bushveld Complex. (Bushveld log and
 1838 data from Maier et al., 2013b). Low. C. Zone = Lower Critical Zone. U.C. Zone = Upper Critical Zone.
 1839 Figure from Maier et al. (2014).



1840

1841 Figure 24. Schematic model of emplacement of the Giles intrusions: a) foundering of crust and
 1842 new SCLM; b) ponding and ascent of mantle melts. Note that the horizontal dimension is greatly
 1843 compressed for added clarity. See text for discussion. SCLM = subcontinental lithospheric mantle.

1844 Figure from Maier et al. (2014).



1845

1846 Figure 25. Comparison of the positions of the PGE reef in a number of well-characterized layered

1847 intrusions. Figure from Maier et al. (2011: 4).