

LETTER TO THE EDITOR

Differential Control by 5-HT and 5-HT_{1A}, 2A, 2C Receptors of Phasic and Tonic GABA Inhibition in the Visual Thalamus

Vincenzo Crunelli^{1,2} & Giuseppe Di Giovanni^{1,2}

¹ Neuroscience Division, School of Bioscience, Cardiff University, Cardiff, UK

² Department of Physiology and Biochemistry, University of Malta, Msida, Malta

Correspondence

Professor Giuseppe Di Giovanni,

Department of Physiology and Biochemistry,

University of Malta,

Msida MSD 2080, Malta

and

Neuroscience Division, School of Biosciences,

Cardiff University,

Museum Avenue,

Cardiff CF10 3AX, UK.

Tel.: +356 23402776

Fax: +356 21310577

E-mails: giuseppe.digiovanni@um.edu.mt and

digiovannig@cardiff.ac.uk

Received 24 August 2015; revision 15

October 2015; accepted 16 October 2015

The copyright line for this article was changed on 10 February 2016 after original online publication.

doi: 10.1111/cns.12480

Thalamocortical (TC) neurons, including those of the dorsal lateral geniculate nucleus (dLGN), one of the visual sensory thalamic nuclei, exhibit two forms of GABA_A receptor-mediated inhibition: phasic or classical inhibitory postsynaptic currents (IPSCs) generated by the activation of synaptic GABA_A receptors (sGABA_AR) and tonic inhibition generated by extra- or perisynaptic GABA_A receptors (eGABA_AR) [1,2]. The source of GABA mediating tonic inhibition mostly arises from spillover out of the synaptic cleft, because tonic inhibition is blocked by TTX and removal of extracellular Ca²⁺ in adult murine dLGN TC neurons [3]. Therefore, modulation of vesicular GABA release may not only affect phasic but also tonic inhibition [1,4]. Previous work in the cat and rat dLGN has shown that several neurotransmitters, including acetylcholine, serotonin (5-HT), dopamine, and norepinephrine can modulate vesicular GABA release from inhibitory interneurons, resulting in changes in phasic inhibition (IPSC frequency), primarily through presynaptic modulation of GABA release from dendro-dendritic synapses [5]. However, except for dopamine in the somatosensory thalamus, the effect of these neurotransmitters on tonic GABA_A inhibition in TC neurons has not been examined. Here, we investigated whether 5-HT and its 5-HT_{1A}, 5-HT_{2A} and 5-HT_{2C} receptors exert a control over tonic and phasic GABA_A currents in dLGN TC neurons. We used whole

cell patch clamp recordings in coronal slices (300 μm) containing the dLGN from postnatal day 20–25 Wistar rats. Data analysis and experimental procedures were similar to those previously described [1,6] and in accordance with the Animals (Scientific Procedures) Act 1986 (UK). Focal application of gabazine (GBZ, 100 μM) was used to reveal the presence of tonic GABA_A current (Figure 1). All serotonergic drugs were dissolved in the recording solution, and their concentrations, co-administration, and effects on phasic and tonic GABA_A current are shown in Table 1 and Figure 1. We found that 5-HT enhances phasic GABA_A inhibition (i.e., spontaneous IPSCs), but has no action on tonic inhibition. These effects are identical to those observed following 5-HT_{1A/7}R activation with 8-OH-DPAT. On the other hand, α-M-5-HT and mCPP enhances and reduces, respectively, both phasic and tonic GABA_A inhibition. These effects are dependent on 5-HT_{2A}R and 5-HT_{2C}R activation, respectively, as they are blocked by co-perfusion with selective antagonists, ketanserin, and SB242084. Thus, the lack of 5-HT modulation of tonic inhibition might be explained by the counterbalance of co-activation of 5-HT_{2A}Rs and 5-HT_{2C}Rs by the endogenous ligand (Figure 1 and Table 1).

Our findings are in agreement with recent evidence in visual cortex showing that 5-HT enhances phasic inhibition by activating 5-HT_{2A}Rs (via calcium/calmodulin-dependent protein kinase II,




Figure 1 Serotonergic modulation of phasic and tonic GABA_A currents in TC neurons of the dLGN. The tonic GABA_A current was measured as the outward change in baseline current following focal (via a pipette) application of the GABA_A antagonist gabazine (100 μM, GBZ, gray bar) (holding potential: -70 mV), as previously described (1). Baseline current was measured as the averaged 20 seconds current before GBZ application, while the shift in baseline current was measured as the averaged 20 seconds current after GBZ application. Focal application of GBZ (100 μM) reveals different magnitude of tonic GABA_A current. Each 5-HT ligand was applied in the recording solution, either alone or in combination, and only one TC neuron was recorded in each slice. sIPSCs recorded in the different experimental conditions were collected before GBZ application and analyzed as previously described [6]. **(A)** Representative current traces from different TC neurons obtained under control condition, and in continuing presence of 5-HT (50 μM, black box), 8-OH-DPAT (100 μM, blue box), α-M-5-HT (100 μM, red box), 100 μM α-M-5-HT and 10 μM ketanserin (gray box), 100 μM α-M-5-HT and 10 μM SB242084 (purple box), 50 μM mCPP (green box), and 50 μM mCPP and 10 μM SB242084 (yellow box). For each current recording (top traces), a representative pre-GBZ period (2 seconds long, bottom traces) shows IPSCs elicited during different 5-HT drug treatments. Summary of the effects of the different 5-HT ligands and their combinations on tonic current **(B)** and sIPSC frequency **(C)**. **P* < 0.05, ***P* < 0.01, ****P* < 0.002 versus control group. ##*P* < 0.01 α-M-5-HT + SB242084 versus α-M-5-HT. ###*P* < 0.01 mCPP and SB242084 versus mCPP. One-way ANOVA, Dunnett's multiple comparison tests. 5-HT, serotonin; 8-OH-DPAT, (±)-2-Dipropylamino-8-hydroxy-1,2,3,4-tetrahydronaphthalene; α-M-5-HT mCPP, meta-chlorophenylpiperazine. sIPSCs, spontaneous inhibitory postsynaptic current. SB 242084, 6-Chloro-2,3-dihydro-5-methyl-N-[6-[(2-methyl-3-pyridinyl)oxy]-3-pyridinyl]-1H-indole-1-carboxamide hydrochloride.

Table 1 Tonic and phasic GABA_A inhibition in dLGN TC neurons under various experimental conditions

Experimental condition	Peak amplitude (pA)	Decay time constant (τ_{decay} , ms)	Charge transfer (fC)	Total current (pA)
Control (n = 22)	-47.0 ± 3.9	5.6 ± 0.4	-207 ± 35.5	-1.2 ± 0.3
50 μM 5-HT (n = 5)	-68.2 ± 8.6*	7.0 ± 1.1	-211 ± 126	-4.8 ± 0.7*
100 μM 8-OH-DPAT (n = 5) 5-HT _{1A/7} agonist	-67.5 ± 3.4*	9.0 ± 1.5**	-157 ± 75.6	-3.4 ± 0.6*
100 μM α -M-5-HT (n = 4) 5-HT _{2A} and 5-HT _{2C} agonist	-98.0 ± 8.9*	4.6 ± 1.1	-139 ± 30.3	-3.5 ± 1.1*
100 μM α -M-5-HT and 10 μM Ketanserin (n = 6) Ketanserin – 5-HT _{2A} and 5-HT _{2C} antagonist	-47.3 ± 9.6	6.5 ± 1.0	-140 ± 51.5	-1.6 ± 0.4
100 μM α -M-5-HT and 10 μM SB242084 (n = 6) SB 242084 – highly selective 5-HT _{2C} antagonist	-64.1 ± 10.6*	5.9 ± 1.1	-164 ± 167	-3.4 ± 0.9*
50 μM mCPP (n = 4) Preferential 5-HT _{2C} agonist	-23.0 ± 4.5*	5.3 ± 2.1	-147 ± 80.2	-0.4 ± 0.2*
50 μM mCPP and 10 μM SB242084 (n = 4)	-27.6 ± 2.8*	6.2 ± 0.7*	-209 ± 21.0	-1.2 ± 0.9

Populations of individual IPSCs in a cell were averaged as described previously [6]. Frequency (not listed Figure 1), peak amplitude, decay time constant (τ_{decay}), and charge transfer of the IPSCs were measured under control condition and during drug application. Number of recorded neurons for each condition is in parentheses. Data are expressed as mean ± SD. Data were analyzed by one-way ANOVA (GraphPad Instat 3 software) followed by *post hoc* analyses (Dunnett's and Dunn's multiple comparison tests) * $P < 0.05$, ** $P < 0.01$ versus control group.

CaMKII) [7]. However, whereas in visual cortex 5-HT decreases tonic inhibition via a 5-HT_{1A}R-dependent suppression of protein kinase A (PKA) activity [7], we could not detect any effect on the tonic current by 5-HT or 5-HT_{1A/7}R activation in the dLGN.

Moreover, our study is in agreement with previous *in vivo* studies in which stimulation of the dorsal raphe nucleus decreased TC neuron firing in the dLGN [8], suggesting 5-HT had an inhibitory action. In contrast, however, *in vitro* intracellular recordings of ferret TC cells found that 5-HT is predominantly hyperpolarizing in all thalamic nuclei tested except for the dLGN, medial geniculate and in a subset of pulvinar neurons, in which depolarizing responses were observed [9]. More recently, it has been shown that in rats 5-HT excites all TC neurons in first-order thalamic nuclei and most (85%) TC neurons in higher order nuclei, while it hyperpolarizes the remaining cells [10]. Specifically in the rat dLGN, 5-HT_{2C}R activation with α -M-5-HT and the highly selective ligand CP-809,101 produced depolarization of TC neurons shifting their firing from bursts to tonic [11]. This would suggest that serotonin has a complex modulatory effect on thalamic nuclei which appear to be nucleus-, 5-HTR subtype-, and 5-HTR synaptic localization-dependent. 5-HT_{1A/7} are present in the thalamus [12], and whereas a strong 5-HT_{2C}R immunoreactivity has been detected, although not somatically, in dLGN TC neurons [13], 5-HT_{2A}R immunostaining did not reach detectable levels [11]. Interestingly, both 5-HT_{2A}R and 5-HT_{2C}R mRNA are expressed in the dLGN GABAergic interneurons of young rats [14]. Therefore, it is likely that the increase in sIPSC frequency that we observed in our study may result from postsynaptic 5-HT₂Rs on dendritic F2 terminals of dLGN interneurons, as previously shown [14].

Nevertheless, 5-HT_{1A/7}R and 5-HT_{2A/2C}Rs might also be located postsynaptically on TC neurons, and their activation may lead to phosphorylation of different subunits of sGABA_ARs and eGABA_ARs acting to differentially modulate their function. The bind-

ing of 5-HT to 5-HT_{1A/7}R and 5-HT_{2A/2C}Rs might activate multiple signal transduction cascades, which ultimately activate different protein kinases, such as PKA or protein kinase C (PKC) and differently regulate sGABA_ARs and eGABA_ARs, as indicated by our results. On the other hand, the potential contribution of 5-HTRs on retinal and cortical terminals can be ruled out as glutamatergic function was blocked in our preparations through the addition of kynurenic acid to the recording solution.

Overall, this study is the first to show a modulation of tonic GABA_A current by 5-HTRs in the thalamus and also to highlight that phasic and tonic inhibition in the dLGN are modulated by 5-HT through different receptor subtypes, leading to a finely tuned balance of sensory information processing in the dLGN. By showing a differential modulation of phasic versus tonic GABA_A inhibition, our results demonstrate a novel mechanism by which the ascending serotonergic afferents can control the thalamic gate to the visual cortex in a behavioral state-dependent manner. Moreover, because of the putative role for thalamic tonic GABA_A inhibition in sleep regulation and pathological oscillations, such as those present in absence epilepsy, the opposing effects of 5-HT_{2A}Rs and 5-HT_{2C}Rs activation may provide suitable targets for pharmacological intervention in sleep and other CNS disorders.

Acknowledgments

Epilepsy Research UK (grant P1202 to VC and GDG), The Wellcome Trust (grant 91882 to VC), and the Malta Council of Science and Technology (grant R&I-2013-14 to GDG and VC).

Conflict of interest

The authors declare no conflict of interest.

References

- Errington AC, Di Giovanni G, Crunelli V, Cope DW. mGluR control of interneuron output regulates feedforward tonic gaba(a) inhibition in the visual thalamus. *J Neurosci* 2011;**31**:8669–8680.
- Errington AC, Di Giovanni G, Crunelli V, editors. *Extrasynaptic GABA_A Receptors*. New York, NY: Springer, 2014.
- Bright DP, Aller MI, Brickley SG. Synaptic release generates a tonic GABA(A) receptor-mediated conductance that modulates burst precision in thalamic relay neurons. *J Neurosci* 2007;**27**:2560–2569.

4. Herd MB, Brown AR, Lambert JJ, Belelli D. Extrasynaptic GABA(A) receptors couple presynaptic activity to postsynaptic inhibition in the somatosensory thalamus. *J Neurosci* 2013;**33**:14850–14868.
5. Cox CL, Sherman SM. Control of dendritic outputs of inhibitory interneurons in the lateral geniculate nucleus. *Neuron* 2000;**27**:597–610.
6. Cope DW, Di Giovanni G, Fyson SJ, et al. Enhanced tonic GABAA inhibition in typical absence epilepsy. *Nat Med* 2009;**15**:1392–1398.
7. Jang HJ, Cho KH, Joo K, Kim MJ, Rhie DJ. Differential modulation of phasic and tonic inhibition underlies serotonergic suppression of long-term potentiation in the rat visual cortex. *Neuroscience* 2015;**301**:351–362.
8. Kayama Y, Shimada S, Hishikawa Y, Ogawa T. Effects of stimulating the dorsal raphe nucleus of the rat on neuronal activity in the dorsal lateral geniculate nucleus. *Brain Res* 1989;**489**:1–11.
9. Monckton JE, McCormick DA. Neuromodulatory role of serotonin in the ferret thalamus. *J Neurophysiol* 2002;**87**:2124–2136.
10. Varela C, Sherman SM. Differences in response to serotonergic activation between first and higher order thalamic nuclei. *Cereb Cortex* 2009;**19**:1776–1786.
11. Coulon P, Kanyshkova T, Broicher T, et al. Activity modes in thalamocortical relay neurons are modulated by g(q)/g(11) family g-proteins – serotonergic and glutamatergic signaling. *Front Cell Neurosci* 2010;**4**:132.
12. Chapin EM, Andrade R. A. 5-HT(7) receptor-mediated depolarization in the anterodorsal thalamus. II. Involvement of the hyperpolarization-activated current I (h). *J Pharmacol Exp Ther* 2001;**297**:403–409.
13. Li QH, Nakadate K, Tanaka-Nakadate S, Nakatsuka D, Cui YL, Watanabe Y. Unique expression patterns of 5-HT_{2A} and 5-HT_{2C} receptors in the rat brain during postnatal development: western blot and immunohistochemical analyses. *J Comp Neurol* 2004;**469**:128–140.
14. Munsch T, Freichel M, Flockerzi V, Pape HC. Contribution of transient receptor potential channels to the control of GABA release from dendrites. *Proc Natl Acad Sci USA* 2003;**100**:16065–16070.