

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/88244/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Wright, V. Paul and Chernes, Lesley 2016. Leaving no stone unturned: the feedback between increased biotic diversity and early diagenesis during the Ordovician. *Journal of the Geological Society* 173 (2) , pp. 241-244. 10.1144/jgs2015-043 file

Publishers page: <http://dx.doi.org/10.1144/jgs2015-043> <<http://dx.doi.org/10.1144/jgs2015-043>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Leaving no stone unturned: the feedback between increased biotic diversity and early**  
2 **diagenesis during the Ordovician**

3 **V. Paul Wright<sup>1</sup> and Lesley Cherns<sup>2\*</sup>**

4 <sup>1</sup>Natural Sciences, National Museum of Wales, Cathays Park, Cardiff, CF10 3NP, UK

5 <sup>2</sup>School of Earth and Ocean Sciences, Cardiff University, Main Building, Park Place, Cardiff,  
6 CF10 3AT, UK

7 \*Corresponding author: cherns@cardiff.ac.uk

8 **(Abstract)**

9 Ordovician change in the nature of seafloor carbonates saw rapid decline of previously  
10 widespread flat pebble conglomerates and the Palaeozoic peak abundance of hardgrounds.  
11 The effective disappearance of flat pebble conglomerates, widely attributed to physical  
12 disruption of substrate by bioturbation, is re-interpreted as reflecting increased depth of  
13 carbonate precipitation below the Taphonomically Active Zone such that early lithified  
14 carbonates were less frequently reworked by scour. With deeper, more stable zones of  
15 cementation, exhumed limestones formed hardgrounds, whose mid Ordovician acme  
16 supported rapid increase in epizoan diversity. Further deepening of cementation to below  
17 normal scour accompanied post-Ordovician decline in submarine hardgrounds.

18 Supplementary material: database for Figure 1 is available at

19 <http://www.geolsoc.org.uk/SUP---->

20

21 The early Palaeozoic evolutionary and ecological development of benthic metazoans was  
22 strongly affected by changes to the nature of the sea floor environment such as increased

23 burrowing activity (e.g. Cambrian Substrate Revolution, Bottjer *et al.* 2000) and widespread  
24 development of shallow marine hard substrates (Great Ordovician Biodiversification Event  
25 (GOBE); Harper 2006). How did those changes impact upon, and relate to, carbonate  
26 systems and shallow sediment diagenesis? Here we consider two characteristic carbonate  
27 facies of early Palaeozoic shelf seas: flat pebble conglomerates (FPC) and submarine  
28 hardgrounds (carbonate cemented sea floors). Both peak in abundance before rapid decline,  
29 the former most widespread in the late Cambrian–early Ordovician, while the latter reach  
30 their Palaeozoic acme in the Mid-Late Ordovician (e.g. Taylor 2008, fig. 2).

31         Flat pebble conglomerates (locally breccias) with carbonate intraclasts (rudstones,  
32 floatstones) are a striking feature of Late Proterozoic to early Ordovician shallow marine  
33 carbonate successions. They were mostly deposited in subtidal, typically offshore settings  
34 reflecting storm or tsunami reworking of shallow cemented limestone beds (Mount & Kidder  
35 1993; Pratt 2002; Myrow *et al.* 2004; Pratt & Bordonaro 2007). The FPCs are variable in bed  
36 geometry and thickness, matrix or clast supported texture, but typically have tabular, thin  
37 (<20mm) pebble to cobble sized clasts of fine grainstone to calcimudstone (e.g. Myrow *et al.*  
38 2004). This distinctive lithofacies effectively disappears from the stratigraphic record in  
39 offshore settings after the Early Ordovician (Sepkoski 1982; Sepkoski *et al.* 1991; Liu &  
40 Zhan 2009). The accepted view (e.g. Sepkoski *et al.* 1991) has been that with an Ordovician  
41 increase in the extent and depth of burrowing (Droser & Bottjer 1989; Bottjer *et al.* 2000),  
42 biotic mixing of the sediment would have prevented early cementation and the formation of  
43 thin lithified zones, removing the source for FPCs after scouring by storms or tsunamis.

44         In the Ordovician, submarine hardgrounds become widely developed in shallow seas,  
45 colonised by an expanding diversity of encrusting and boring epizoans (e.g. Brett &  
46 Brookfield 1984) that form specialized new communities in the GOBE. They have been

47 interpreted as indicating seafloor calcite precipitation and aragonite dissolution in ‘calcite  
48 seas’ (Taylor & Wilson 2003; Palmer & Wilson 2004; Harper 2006; Taylor 2008).

49 The aim of this paper is to provide a single, geochemical explanation for both the  
50 decline of subtidal FPC lithofacies and the peak abundance of hardgrounds during the  
51 Ordovician.

52

### 53 **Sea floor diagenesis and changes in sea floor shallow geochemical profiles during the** 54 **early Palaeozoic**

55 The diagenetic processes for mobilization of calcium carbonate during shallow burial have  
56 been appreciated for some time (e.g. Sanders 2003; Berkeley *et al.* 2008; Cherns *et al.* 2011).  
57 Calcium carbonate, especially the more soluble aragonite, is dissolved in the uppermost  
58 sediment layer largely as a result of acidity caused by the oxidation of H<sub>2</sub>S, and while most  
59 back-fluxes to the water column, some is re-precipitated as calcite in the sediment column in  
60 areas of increased alkalinity such as depths where sulfate reduction takes place (e.g. Sanders  
61 2003, 2004; Fig. 1). This oxidized zone, effectively the Taphonomically Active Zone (TAZ),  
62 will be controlled by diffusion from the overlying water column, if oxygenated, and by  
63 mixing caused by bio-irrigation (mainly burrowing; e.g. Aller 1982; Aller & Aller 1998).  
64 Organic matter accumulated more among finer grained sediment sources the microbially  
65 mediated decay processes that drive skeletal carbonate dissolution and re-precipitation (e.g.  
66 Walter & Burton 1990; Walter *et al.* 1993; Hendry 1993). The importance of such processes  
67 linked to the mobilization of labile carbonates in the very shallow sediment column forms the  
68 basis for understanding the limestone-marl alternations that form a widely developed facies in  
69 Phanerozoic epeiric sea settings (Munnecke & Samtleben 1996; Westphal & Munnecke  
70 2003; Munnecke & Westphal 2005).

71           Brasier *et al.* (2011) proposed that changes in the position, relative to the sea floor, of  
72 the depth of the redox boundary during the Ediacaran–early Cambrian affected the zones of  
73 early lithification in the shallow sediment column. Late Proterozoic precipitation of calcium  
74 carbonate took place at or very close (<1cm) to the sea floor (also Peters & Gaines 2012), and  
75 though the early Cambrian advent of metazoan burrowing and biomineralization depressed  
76 the zone of cementation it remained very shallow (early-mid Cambrian subtidal burrow depth  
77 <3 cm, typically mm scale: Tarhan *et al.* 2015; Droser & Bottjer 1988). The depth of subtidal  
78 bioturbation, and by inference the zone of cementation, remained <6 cm through to the mid  
79 Ordovician, before both bioturbation depth (<30 cm) and intensity increased significantly in  
80 the Late Ordovician (Droser & Bottjer 1989).

81

## 82 **Abundance of flat pebble conglomerates and hardgrounds**

83 Subtidal FPCs are most common in the late Cambrian–Early Ordovician, before rapid  
84 decline, and notably while the TAZ remained very shallow (Fig. 2). Their temporal record,  
85 using publications by formation as a proxy for abundance, provides a direct comparison with  
86 published data for submarine hardgrounds (Fig. 2; Taylor 2008). A peak FPC distributional  
87 map illustrates the extensive occurrence (with a latitudinal control), and suggests any bias  
88 from availability of rock formations is likely not significant (Fig. 3). In shallow carbonate  
89 epeiric seas of the North China Plate, the subsequent decline of FPCs corresponds to the  
90 decrease also in subtidal microbialites and increasing intensity of bioturbation as the shallow  
91 sea floor character changed in the late Early Ordovician (early Floian) (Liu 2009; Liu & Zhan  
92 2009). Notably, in post-Ordovician times, minor occurrences of subtidal FPCs (Fig. 2)  
93 correspond to post-extinction events, when suppression of bioturbation would have led to  
94 shallowing of the TAZ (e.g. Wignall & Twitchett 1999; Calner 2005).

95           For submarine hardgrounds (carbonate cemented sea floors) the Palaeozoic peak of  
96 abundance is in the Ordovician (Taylor & Wilson 2003; Palmer & Wilson 2004; Harper  
97 2006; Taylor 2008; Fig. 2). Early, encrusting hardground faunas are described from surfaces  
98 of cemented FPC in the late Cambrian (Brett *et al.* 1983). By the mid Ordovician hard  
99 substrate morphologies are variable, some largely comprising reworked, encrusted limestone  
100 nodules but others forming beds with complex hummocky and undercut surfaces; hardground  
101 biotas are notably more diverse (Brett & Liddell 1978; Brett & Brookfield 1984; Wilson *et al.*  
102 1992). Taylor and Wilson (2003, p. 44) suggested that the “Ordovician was a golden age for  
103 epizoans on hard substrates” due in part to increased hard substrate availability. The  
104 appearance of encrusters makes hardgrounds more recognizable after the early Ordovician  
105 (Brett & Liddell 1978) and likely reflects the availability of more stable substrates as  
106 compared with the fragmented cemented layers characteristic earlier in the Palaeozoic.

107           Previously the abundance of hardgrounds had been explained by local calcite  
108 cementation sourced from carbonate released by sea-floor dissolution of aragonite in  
109 undersaturated (with respect to aragonite) Ordovician ‘calcite seas’ (Wilson *et al.* 1992;  
110 Taylor & Wilson 2003; Palmer & Wilson 2004; Harper 2006). Recent experimental data,  
111 however, indicate that aragonite precipitation continued alongside calcite during ‘calcite seas’  
112 in warm water environments (Balthasar & Cusack 2015). In a study of Ordovician  
113 hardgrounds from eastern North America, Kenyon-Roberts (1995) found no direct evidence  
114 for sea floor dissolution, but did note petrographic evidence of hardground formation in  
115 shallow sub-oxic conditions below the TAZ. Cherns & Wright (2011), when comparing the  
116 taphonomy of skeletal lagerstätten between ‘aragonite’ and ‘calcite seas’, found no  
117 differences, suggesting that ‘calcite seas’ did not increase aragonitic shell dissolution.

118

119 **Diagenetic model for Early Palaeozoic subtidal settings** (Fig. 4)

120 As the depth and intensity of burrowing increased through the early Palaeozoic, the TAZ  
121 thickened and the depth at which secondary carbonate re-precipitated also deepened. This  
122 reduced the probability that lithified carbonate would be exhumed by erosional reworking  
123 caused by wave scour. The FPCs in shallow subtidal settings formed only while the TAZ  
124 was very thin, and hence the zone of cementation was close to the sediment-water interface  
125 allowing even relatively small and frequent scour events (storms or tsunamis) to exhume the  
126 cemented horizons (Late Precambrian–late Early Ordovician; Figs 2, 4). As the TAZ  
127 thickened and the depth of carbonate precipitation increased, these horizons were less likely  
128 to be reworked by scouring. The FPC facies was replaced in shallow subtidal settings by less  
129 frequently exhumed, and hence more developed and thicker, cemented horizons, which when  
130 eventually exposed by scour formed reworked concretions and hardgrounds on which hard  
131 substrate biotas expanded (Mid–Late Ordovician; Figs. 2, 4). Development of these carbonate  
132 horizons may have been facilitated by increased skeletal input through diversification and  
133 faunal expansion during this interval (Porter 2010). In more offshore settings rarely affected  
134 by wave-related erosion, the secondary carbonate could accumulate uninterrupted to produce  
135 the nodular, diagenetic bedding of limestone-marl alternations. As the TAZ deepened  
136 through the later Ordovician the cementation zone was displaced to deeper levels where  
137 reduced likelihood of exhumation led to decline in hardground abundance and more  
138 widespread development of diagenetic bedding (Late Ordovician; Fig. 3; Westphal 2006, fig.  
139 2). An implication is that diagenetic bedding would have been preserved in shallower areas  
140 than previously, assuming that shallower sea floors were more susceptible to periodic  
141 reworking than deeper ones (Peters & Loss 2012). This hypothesis is testable if it can be  
142 demonstrated that diagenetic bedding is found in shallower settings by the Late Ordovician.

143

144 **Discussion**

145 This model proposes that a single trend, namely the deepening of the zone of cementation  
146 below the TAZ, can explain the rapid decline of FPCs, and the peak abundance of  
147 hardgrounds during the Ordovician, before their subsequent decline. That progressive change  
148 was ultimately a consequence of the previously documented increased depth and intensity of  
149 bio-irrigation. Rather than destroying the potential for rapid shallow cementation (Sepkoski  
150 1982; Sepkoski et al. 1991), lowering of the TAZ decreased the likelihood of erosional  
151 exhumation of thin cemented carbonate layers by wave scour.

152 Many variables affected early diagenesis during the early Palaeozoic, including  
153 increased nutrient-rich organic matter and oxygen levels in sea floor sediments as a result of  
154 metazoan evolution (McIlroy & Logan 1999; Bottjer *et al.* 2000, Dornbos *et al.* 2005). The  
155 amount of labile aragonite from skeletal carbonate dissolution also likely increased with  
156 biomineralization, although the skeletal contribution to carbonate deposition remained limited  
157 up to mid Ordovician times (e.g. Pruss *et al.* 2010, fig. 8). Could the thinness of limestone  
158 beds ripped up to form FPCs from latest Proterozoic to Early Ordovician be explained by  
159 lower flux of carbonate from skeletal dissolution before the GOBE rather than reflecting  
160 frequency of exhumation at shallow burial depths? Although the intensity of bio-irrigation  
161 increased markedly in the Early Cambrian, bioturbation depth (<6 cm) remained shallow  
162 through to the Late Ordovician, when both depth (<30 cm) and average ichnofabric index  
163 increased substantially (Droser & Bottjer 1989; Tarhan *et al.* 2015). If changes in the  
164 biogeochemical environment of the upper sediment layers were a consequence of that deeper  
165 and more intense bio-irrigation, accompanied by increased oxygenation of the seas and  
166 oversaturation (Pruss *et al.* 2010) that affected diagenetic carbonate precipitation, what other  
167 effects took place in terms of carbonate behaviour? The biogenically reworked mixed layer  
168 increased only slowly, from 0.2 cm in the early-mid Cambrian to 1.5 cm in the Ordovician-



169 Silurian (Tarhan *et al.* 2015). Could the presence of a thin TAZ potentially result in higher  
170 levels of acidity through sulfide oxidation and a greater degree of undersaturation with  
171 respect to aragonite, compared with today's thicker TAZ?

172         Organisms with more labile, aragonitic shells (primarily molluscs) are diverse in the  
173 early Cambrian radiation although their fossils are relatively sparse in the trilobite dominated  
174 Cambrian–Lower Ordovician skeletal record (Porter 2007; Porter *et al.* 2010). From the mid  
175 Ordovician, skeletal material was a major contributor to carbonate sediment; limestone shell  
176 beds, most commonly brachiopod-rich, increase in proportion, thickness and abundance in  
177 shallow marine settings (Kidwell and Brenchley 1994; Li & Droser 1997, 1999; Pruss *et al.*  
178 2010). Molluscs are dominant in some storm beds, most commonly representing local  
179 reworking of concentrations of dead gastropod shells accumulated in the upper sediment  
180 layers (Li & Droser 1999; Harper 2006, fig. 9). Did a thicker TAZ later in the Ordovician  
181 result in less intense sulfide oxidation, less dissolution and a longer survival time of aragonite  
182 shells in the TAZ?

183         The Palaeozoic decrease in abundance of hardgrounds after the Ordovician (Fig. 2) is  
184 here interpreted as reflecting the lowering of the TAZ and cementation zone to depths in the  
185 sediment column affected less frequently by wave (storm or tsunami) reworking. It might  
186 also imply that such reworking later rarely affected sediments much below the TAZ at 30cm.  
187 Mesozoic peaks of hardground occurrence (Fig. 2) may in part reflect large outcrop areas of  
188 marine sediments (Smith and McGowan 2007), such as the Cretaceous Chalk. The extensive  
189 hardgrounds of the middle Jurassic are hosted predominantly in very shallow, oolitic facies  
190 and are of much more diverse origins than those of the Ordovician (Kenyon-Roberts 1995).

191

192 **Conclusions**

193 The decline of flat pebble conglomerates and the peak abundance of submarine hardgrounds  
194 in the Ordovician are interpreted as reflecting the progressive deepening of the zone of  
195 carbonate precipitation below the TAZ, resulting in less frequent reworking of the upper part  
196 of the sediment column by scour. The deepening was a consequence of increased depth and  
197 intensity of bioturbation and bio-irrigation. This shift also created a range of more lithified  
198 substrates in subtidal settings, promoting a rapid expansion in epizoan diversity. Thus,  
199 changes in bioturbation affected carbonate diagenesis and the composition of the sea floor  
200 carbonates, and provided new niches for invertebrates.

201

## 202 **Acknowledgments**

203 We thank Carl Brett for discussion, and are grateful to Paul Smith and an unknown reviewer  
204 for constructive criticism of the manuscript.

205

## 206 **References**

207 Aller, R.C. 1982. Carbonate dissolution in nearshore terrigenous muds: the role of physical  
208 and biological reworking. *Journal of Geology*, **90**, 79-95.

209 Aller, R.C. & Aller, J.Y. 1998. The effect of biogenic irrigation intensity and solute exchange  
210 on diagenetic reaction rates in marine sediments. *Journal of Marine Research*, **56**,  
211 905-936.

212 Balthasar, U. & Cusack, M. 2015. Aragonite-calcite seas—Quantifying the gray area.  
213 *Geology*, **43**, 99-102.

- 214 Berkeley, A., Perry, C.T., Smithers, S.G., Horton, B.P. & Taylor, K.G. 2007. A review of the  
215 ecological and taphonomic controls on foraminiferal assemblage development in  
216 intertidal environments. *Earth-Science Reviews*, **83**, 205-230.
- 217 Bottjer, D.J., Hagadorn, J.W. & Dornbos, S.Q. 2000. The Cambrian substrate revolution.  
218 *GSA today*, **10**, 1-7.
- 219 Brasier, M.D., Antcliffe, J.B. & Callow, R.H.T. 2011. Evolutionary trends in remarkable  
220 fossil preservation across the Edicaran-Cambrian transition and the impact of  
221 metazoan mixing. *In*: Allison, P.A. & Bottjer, D.J. (eds.) *Taphonomy: process and*  
222 *bias through time*. Springer, Berlin, 519-567.
- 223 Brett, C.E. & Brookfield, M.E. 1984. Morphology, faunas and genesis of Ordovician  
224 hardgrounds from southern Ontario, Canada. *Palaeogeography, Palaeoclimatology,*  
225 *Palaeoecology*, **46**, 233-290.
- 226 Brett, C.E. & Liddell, W.D. 1978. Preservation and palaeoecology of a Middle Ordovician  
227 hardground community. *Paleobiology*, **4**, 329-348.
- 228 Brett, C.E., Liddell, W.D. & Derstler, K.L. 1983. Late Cambrian hard substrate communities  
229 from Montana/Wyoming: the oldest known hardground encrusters. *Lethaia*, **16**, 281-  
230 289.
- 231 Calner, M. 2005. A Late Silurian extinction event and anachronistic period. *Geology*, **33**,  
232 305-308.
- 233 Chrens, L., Wheeley, J.R. & Wright, V.P. 2011. Taphonomic bias in shelly faunas through  
234 time: early aragonite dissolution and its implications for the fossil record. *In*: Allison,  
235 P.A. & Bottjer, D.J. (eds.) *Taphonomy: process and bias through time*. Springer,  
236 Berlin, 79-105.

- 237 Cherns, L. & Wright, V.P. 2011. Skeletal mineralogy and biodiversity of marine  
238 invertebrates: size matters more than seawater chemistry. *Geological Society, London,*  
239 *Special Publications*, **358**, 9-17.
- 240 Dornbos, S.Q., Bottjer, D.J. & Chen, J.-Y. 2005. Paleoecology of benthic metazoans in the  
241 Early Cambrian Maotianshan Shale biota and the Middle Cambrian Burgess Shale  
242 biota: evidence for the Cambrian substrate revolution. *Palaeogeography,*  
243 *Palaeoclimatology, Palaeoecology*, **220**, 47-67.
- 244 Droser, M.L. & Bottjer, D.J. 1988. Trends in depth and extent of bioturbation in Cambrian  
245 carbonate marine environments, western United States. *Geology*, **16**, 233-236.
- 246 Droser, M.L. & Bottjer, D.J. 1989. Ordovician increase in extent and depth of bioturbation:  
247 Implications for understanding early Paleozoic ecospace utilization. *Geology*, **17**, 850-  
248 852.
- 249 Harper, D.A. 2006. The Ordovician biodiversification: setting an agenda for marine life.  
250 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **232**, 148-166.
- 251 Hendry, J.P. 1993. Calcite cementation during bacterial manganese, iron and sulphate  
252 reduction in Jurassic shallow marine carbonates. *Sedimentology*, **40**, 87-106.
- 253 Kenyon-Roberts, S.M. 1995. *The petrography and distribution of some calcite sea*  
254 *hardgrounds*. PhD thesis, University of Reading.
- 255 Kidwell, S.M. & Brenchley, P.J. 1994. Patterns in bioclastic accumulation through the  
256 Phanerozoic - changes in input or in destruction?. *Geology*, **22**, 1139-1143.

- 257 Li, X. & Droser, M.L. 1997. Nature and distribution of Cambrian shell concentrations:  
258 Evidence from the Basin and Range Province of the Western United States  
259 (California, Nevada, and Utah). *PALAIOS*, **12**, 111-126.
- 260 Li, X. & Droser, M.L. 1999. Lower and Middle Ordovician shell beds from the Basin and  
261 Range province of the western United States (California, Nevada, and Utah).  
262 *PALAIOS*, **14**, 215-233.
- 263 Liu, J. 2009. Marine sedimentary response to the Great Ordovician Biodiversification Event:  
264 examples from North China and South China. *Paleontological Research*, **13**, 9-21.
- 265 Liu, J. & Zhan, R. 2009. Temporal distribution of diagnostic biofabrics in the Lower and  
266 Middle Ordovician in North China: clues to the geobiology of the Great Ordovician  
267 Biodiversification Event. *Acta Geologica Sinica*, **83**, 513-523.
- 268 McIlroy, D. & Logan, G.A. 1999. The impact of bioturbation on infaunal ecology and  
269 evolution during the Proterozoic-Cambrian transition. *PALAIOS*, **14**, 58-72.
- 270 Mount, J.F. & Kidder, D. 1993. Combined flow origin of edgewise intraclast conglomerates:  
271 Sellick Hill Formation (Lower Cambrian), South Australia. *Sedimentology*, **40**, 315-  
272 329.
- 273 Munnecke, A. & Samtleben, C. 1996. The formation of micritic limestones and the  
274 development of limestone-marl alternations in the Silurian of Gotland, Sweden.  
275 *Facies*, **34**, 159-176.
- 276 Munnecke, A. & Westphal, H. 2005. Variations in primary aragonite, calcite, and clay in  
277 fine-grained calcareous rhythmites of Cambrian to Jurassic age — an environmental  
278 archive? *Facies*, **51**, 611-626.

- 279 Myrow, P., M. , Tice, L., Archuleta, B., Clark, B., Taylor, J.F. & Ripperdan, R.L. 2004. Flat-  
280 pebble conglomerate: its multiple origins and relationship to metre-scale depositional  
281 cycles. *Sedimentology*, **51**, 973-996.
- 282 Palmer, T. & Wilson, M. 2004. Calcite precipitation and dissolution of biogenic aragonite in  
283 shallow Ordovician calcite seas. *Lethaia*, **37**, 417-427.
- 284 Peters, S.E. & Gaines, R.R. 2012 Formation of the ‘Great Unconformity’ as a trigger for the  
285 Cambrian explosion. *Nature*, **484**, 363-366.
- 286 Peters, S.E. & Loss, D.P. 2012. Storm and fair-weather wave base: A relevant distinction?  
287 *Geology*, **40**, 511-514.
- 288 Porter, S.M. 2007. Seawater chemistry and early carbonate biomineralization. *Science*, **316**,  
289 1302-1302.
- 290 Porter, S.M. 2010. Calcite and aragonite seas and the de novo acquisition of carbonate  
291 skeletons. *Geobiology*, **8**, 256-277.
- 292 Pratt, B.R. 2002. Storms versus tsunamis: Dynamic interplay of sedimentary, diagenetic, and  
293 tectonic processes in the Cambrian of Montana. *Geology*, **30** 423-426.
- 294 Pratt, B.R. & Bordonaro, O.L. 2007. Tsunamis in a stormy sea: Middle Cambrian inner-shelf  
295 limestones of western Argentina. *Journal of Sedimentary Research*, **77**, 256-262.
- 296 Pruss, S.B., Finnegan, S., Fischer, W.W. & Knoll, A.H. 2010. Carbonates in skeleton-poor  
297 seas: new insights from Cambrian and Ordovician strata of Laurentia. *PALAIOS*, **25**,  
298 73-84.

- 299 Sanders, D. 2003. Syndepositional dissolution of calcium carbonate in neritic carbonate  
300 environments: geological recognition, processes, potential significance. *Journal of*  
301 *African Earth Sciences*, **36**, 99-134.
- 302 Sanders, D. 2004. Potential significance of syndepositional carbonate dissolution for platform  
303 banktop aggradation and sediment texture: a graphic modeling approach. *Austrian*  
304 *Journal of Earth Sciences*, **95/96**, 71-79.
- 305 Sepkoski, J.J., Jr. 1982. Flat-pebble conglomerates, storm deposits, and the Cambrian bottom  
306 fauna. In: Einsele, G. & Seilacher, A. (eds.) *Cyclic and Event Stratification*. Springer-  
307 Verlag, 371-386.
- 308 Sepkoski, J.J., Jr, Bambach, R.K. & Droser, M.L. 1991. Secular changes in Phanerozoic  
309 event bedding and the biological overprint. In: Einsele, G., Ricken, W. & Seilacher,  
310 A. (eds.) *Cycles and events in stratigraphy*. Springer-Verlag, 298-312.
- 311 Smith, A.B. & McGowan, A.J. 2007. The shape of the Phanerozoic marine palaeodiversity  
312 curve: how much can be predicted from the sedimentary rock record of Western  
313 Europe? *Palaeontology*, **50**, 765-774.
- 314 Tarhan, L.G., Droser, M.L., Planavsky, N.J. & Johnston, D.T. 2015. Protracted development  
315 of bioturbation through the early Palaeozoic Era. *Nature Geoscience*, advance online  
316 publication, doi: 10.1038/ngeo2537.
- 317
- 318 Taylor, P.D. 2008. Seawater chemistry, biomineralization and the fossil record of calcareous  
319 organisms. In: Okada, H., Mawatari, S.F., Suzuki, N. & Gautam, P. (eds.) *Origin and*  
320 *Evolution of Natural Diversity: Proceedings of International Symposium "The Origin*

321 and Evolution of Natural Diversity", 1-5 October 2007, Sapporo, Japan. Hokkaido  
322 University. 21-29.

323

324 Taylor, P.D. & Wilson, M.A. 2003. Palaeoecology and evolution of marine hard substrate  
325 communities. *Earth-Science Reviews*, **62**, 1-103.

326 Torsvik, T.H. 2009. *Bugplates software (IGCP503)*. With reconstructions of Torsvik, T. H.  
327 and Cocks, L. R. M. (2002-2009). StatoilHydro.  
328 <http://www.geodynamics.no/bugs/SoftwareManual.pdf>.

329 Walter, L.M., Bischof, S.A., Patterson, W.P. & Lyons, T.W. 1993. Dissolution and  
330 Recrystallization in Modern Shelf Carbonates - Evidence from Pore-Water and Solid-  
331 Phase Chemistry. *Philosophical Transactions of the Royal Society of London, Series*  
332 *A*, **344**, 27-36.

333 Walter, L.M. & Burton, E.A. 1990. Dissolution of platform carbonate sediments in marine  
334 pore fluids. *American Journal of Science*, **290**, 601-643.

335 Westphal, H. 2006. Limestone–marl alternations as environmental archives and the role of  
336 early diagenesis: a critical review. *International Journal of Earth Sciences*, **95**, 947-  
337 961.

338 Westphal, H. & Munnecke, A. 2003. Limestone-marl alternations: A warm-water  
339 phenomenon? *Geology*, **31**, 263-266.

340 Wignall, P.B. & Twitchett, R.J. 1999. Unusual intraclastic limestones in Lower Triassic  
341 carbonates and their bearing on the aftermath of the end-Permian mass extinction.  
342 *Sedimentology*, **46**, 303-316.



343 **Figure Captions**

344 Figure 1. Shallow burial diagenetic environment and calcium carbonate precipitation. A,  
345 labile aragonite shells (molluscs), more susceptible to early dissolution in the oxic upper  
346 sediment layers of the Taphonomically Active Zone (TAZ), release carbonate and leave  
347 moulds that are readily destroyed through bioturbation. B, calcitic shells more likely to  
348 survive early dissolution, and more rarely steinkerns of aragonitic shells; diffused carbonate  
349 precipitates in a zone of cementation in the sulfate reduction zone.

350 Figure 2. Abundance (using publications by formation as proxy) of subtidal flat pebble  
351 conglomerates (pale, points; supplemental information available online  
352 at [www.geolsoc.org.uk/SUP0xxxx](http://www.geolsoc.org.uk/SUP0xxxx)) and submarine hardgrounds (dark, histogram; based on  
353 Taylor 2008 and <http://markwilson.voices.wooster.edu/bioerosion-bibliography/>). Note:  
354 revision of Cambrian stratigraphy into four series is ongoing; divisions into Lower, Middle  
355 and Upper series follow standard usage in literature.

356 Figure 3. Palaeogeographic reconstruction for 485 Ma (Bugplates IGCP503; T.H. Torsvik  
357 2009) showing late Cambrian–early Ordovician extent of flat pebble conglomerate facies  
358 (black stars).

359 Figure 4. Diagenetic model for carbonate precipitation in subtidal Palaeozoic settings,  
360 showing early Cambrian through Ordovician changes in the depth of the Taphonomically  
361 Active Zone (TAZ) and zone of cementation, susceptibility of lithified limestone layers to  
362 scouring, and distribution of flat pebble conglomerates, submarine hardgrounds and the  
363 diagenetic nodular bedding of limestone-marl alternations. SWI sediment-water interface.

364

365 Figure 1.

366

367

368

369


370

371

372

373

374



375

376 Figure 2.

377

378

379

380

381

382

383

384


385



386

387

388

389





## 404 Flat pebble conglomerates from subtidal settings (Fig. 2)

AGE	LOCATION	FORMATION	SHELF SETTING	PROPOSED PROCESS	AUTHOR(S)
Mesoproterozoic	China, Hebei Province	Gaoyuzhuang Formation	Subtidal within storm wave base	storms	Luo <i>et al.</i> 2014
Proterozoic, Pre-Marinoan glaciation	NW Canada, Mackenzie Mountains	Keele Fm	mid to outer ramp	storms	Day <i>et al.</i> 2004
Vendian	Gourma, West Africa		slope	bottom currents'	Bertrand-Sarfati & Moussine-Pouchkine 1983
Ediacaran	South Africa	Swartpunt Fm	Low energy deeper ramp	storm	Narbonne <i>et al.</i> 1997
Ediacaran	Canadian Arctic	Gametrail Fm	subtidal ramp	storms	MacNaughton <i>et al.</i> 2008
Ediacaran	Kazakhstan	Kyrshabakty Formation	shallow carbonate platform	high energy events	Heubeck <i>et al.</i> 2013
Ediacaran - Lower Cambrian	Oman	Ara Group	carbonate platform		Grotzinger & Al-Rawahi 2014
Lower Cambrian	South Australia	Sellick Hill Formation	subtidal	storms	Mount & Kidder 1993
Lower Cambrian	W Mongolia	Bayan Gol Fm, Zavkhan Basin	shallow subtidal	storms	Kruse <i>et al.</i> 1996
Lower Cambrian	South China	Shuijingtuo Fm	subtidal		Ishikawa <i>et al.</i> 2008
Middle Cambrian	Canadian Arctic		ramp		Dewing & Nowlan 2012
Middle Cambrian	British Columbia, Canada	Jubilee Fm			Pope 1990

Middle Cambrian	Argentina	La Laja Fm	subtidal shelf	tsunamis	Pratt & Bordonaro 2007
Middle Cambrian	Australia	Ranken Lst	low energy shallow subtidal	storms	Kruse 1996
Middle Cambrian	Wyoming, USA	Upper Gros Ventre Shale			Csonka 2009
upper Middle Cambrian	W Utah, USA	upper Wheeler, Marjum fms	middle carbonate belt - subtidal shelf		Robison 1964
Middle-Upper Cambrian	NW China		Supratidal to subtidal fpc	storms	Liang <i>et al.</i> 1993
Middle Cambrian - Lower Ordovician	Siberia	Ust'-Brus, Labaz, Orakta, Kulyumbe, Ujgur and Iltyk fms			Kouchinsky <i>et al.</i> 2008
Upper Cambrian	NW Siberia	Chopko Fm, Chopka River	carbonate platform, turbidites	submarine landslides	Varlamov <i>et al.</i> 2006
Upper Cambrian	N China	Gushan, Chanshang formations	subtidal shelf	storms	Ding <i>et al.</i> 2008; Meng <i>et al.</i> 1997
Upper Cambrian	China, Shandong Province	Chaomidian Formation (Furongian)	shallow subtidal		Lee <i>et al.</i> 2010; Chen 2014; Chen <i>et al.</i> 2009, Chen <i>et al.</i> 2010; Van Loon <i>et al.</i> 2013
Upper Cambrian	S Korea	Hwajeol Formation	subtidal, relatively deep		Kim & Lee 2000
U Cambrian	Western USA		Outer detrital belt (subtidal lagoon)	Storms	Sepkoski 1982
Upper Cambrian	Rocky Mts USA	Snowy Range Fm (Sunwaptan-L Skullrockian)	Inner detrital belt, subtidal	Storms (leading to slope failure)	Brett <i>et al.</i> 1983; Myrow <i>et al.</i> 2004; Myrow <i>et al.</i> 2012
Upper Cambrian	Montana USA	Deadwood Fm	subtidal shelf	tsunamis	Pratt 2002

Upper Cambrian	Wyoming, USA	Snowy Range Fm - Upper Deadwood Formation	subtidal intrashelf basin	storms	Saltzman 1999; Myrow <i>et al.</i> 2004
Upper Cambrian	Wyoming, USA			fpc produced by dewatering	Wiison 1985; Kozub 1997
Upper Cambrian	Virginia, USA	Nolichucky Formation	shallow subtidal basin facies	storms	Markello & Read 1981, 1982
Upper Cambrian	Nevada and Utah, USA	SPICE interval	Intrashelf basin	storms	Saltzman <i>et al.</i> 1998
Upper Cambrian	Montana USA	Grove Creek, Snowy Range, Maurice formations			Dorf & Lochman 1940
Upper Cambrian	Maryland, USA	Conococheague Limestone	sand shoal environments		Demicco 1985; Demicco <i>et al.</i> 1991
Upper Cambrian	Virginia, USA	Conococheague Limestone, Copper Ridge Dolomite	Group I outer shelf	storms	Whisonant 1987
Upper Cambrian	Tennessee, USA	Maynardsville Fm, Conosauga Group	subtidal		Glumac and Walker 1997
Upper Cambrian	Wisconsin and Minnesota, USA	Tunnel City Group	shallow subtidal	storms	Eoff 2014
Upper Cambrian	California, USA	Nopah Fm (Sunwaptan); also Desert Valley Formation, Whipple Cave Formation, Notch Peak Formation, Ajax Dolomite	shallow subtidal	storms	Shapiro & Awramik 2006
Upper Cambrian	S Alberta, Canada	Bison Creek and Mistaya formations	shallow subtidal shelf	storms	Westrop 1989

Upper Cambrian - Lower Ordovician	Alberta , Canada	Survey Peak Fm; Ibexian-Tremadoc			Ji & Barnes 1996
Upper Cambrian - Lower Ordovician	Mexico	Tiñu Fm	subtidal dysoxic shelf	debris flows	Landing <i>et al.</i> 2007
Upper Cambrian - Lower Ordovician	N China	Fengshan Formation - Yeli Formation	subtidal shelf		Yang <i>et al.</i> 2002
Upper Cambrian - Lower Ordovician	China, Jilin Province	candidate GSSP Xiaoyangqiao	subtidal	storms	Chen <i>et al.</i> 1988
Upper Cambrian - Lower Ordovician	Utah, USA; Nevada USA	Notch Peak and House Limestone fms; Whipple Cove and House Limestone fms	shoals on shallow carbonate shelf		Popov <i>et al.</i> 2002; Cook & Taylor 1975, 1977
Upper Cambrian - Lower Ordovician	Colorado, USA	Dotsero Fm, Manitou Fm			Berg 1960
Lower and Upper Cambrian, Lower Ordovician	Appalachians , USA	Dunham Fm; Pine Plains Fm; Ogdenburg and Tribes Hill fms		storms	Friedman 1994
Upper Cambrian - Lower Ordovician	Siberia	Nya sequence	shallow carbonate platform		Dronov <i>et al.</i> 2009
Upper Cambrian - Lower Ordovician	Mid-East Korea	Choson Supergroup	subtidal	storm, diagenetic lts	Kwon <i>et al.</i> 2002
Lower Ordovician	Newfoundland	Watt's Bight and Boat harbour fms	deep subtidal to peritidal	storms	Pruss <i>et al.</i> 2010

Lower Ordovician	Utah, USA	Pogonip Group - Notch Peak Formation, House Limestone, Fillmore Formation, and Wah Wah Limestone	shallow subtidal to peritidal		Pruss <i>et al.</i> 2010
Lower Ordovician	S Korea	Dumugol Fm	shallow to deep ramp	storms	Lee & Kim 1992
Lower Ordovician	Korea	Mungok Fm	subtidal shelf	storms	Kim and Lee 1995; Choi, Kim & Lee 1993
Lower Ordovician	NY, USA	Tribes Hill Fm	intertidal to supratidal	desiccation and high energy events (seismic /storm /tsunami?)	Braun & Friedman 1969
Lower Ordovician (Tremadoc)	Pingquan, Hebei Province, N China		shallow subtidal to shaly basinal		Liu & Zhan 2009
Lower Ordovician	Pingquan, Hebei Province, N China and Xingshan, Hubei Province, S China		Lower Tremadoc shallow subtidal, Upper Tremadoc shallow to deep subtidal		Liu 2009
Lower Ordovician	NW Hubei	Nantsinkian-lower Dawan fms; Tremadoc - early Floian	shallow marine carbonate platform; shallow to deeper subtidal		Liu <i>et al.</i> 2011
Lower Ordovician	Nevada, USA	Ninemile Shale	within storm wave base	storms	Sprinkle & Guensburg 1995



Lower Ordovician	Utah and Nevada, USA	Kanosh Shale	Intrashelf basin	storms	Wilson <i>et al.</i> 1992
Lower Ordovician	Utah	Fillmore Formation	storm dominated shelf	storms	Sprinkle & Guensburg 1995; Dattilo 1993; Benner <i>et al.</i> 2004
Upper Silurian	Gotland, Sweden	upper Hemse-Eke fms	subtidal to very shallow, microbial shoals	anachronistic facies - suppressed burrowing	Cherns 1982, 1983; Calner 2005
Upper Silurian	Somerset Is., Arctic Canada	Reach Bay Fm	Subtidal within storm wave base	storms	Jones & Dixon 1976
Upper Devonian (Frasnian)	Holy Cross Mts, Poland		Shallow subtidal	storms or tsunamis	Kazmierczak & Goldring 1978
Lower Triassic	S Turkey	Dienerian Fm	storms affecting shallow shelf	anachronistic facies - suppressed burrowing	Pruss <i>et al.</i> 2006
Lower Triassic	South China; North Italy		storm-dominated shelf to deep basin; mid ramp carbonates	anachronistic facies - suppressed burrowing	Wignall & Twitchett 1999
Lower Triassic	SW USA	Moenkopi - Union Wash formations	subtidal to deep	anachronistic facies - suppressed burrowing	Pruss <i>et al.</i> 2005; Woods 2009
Lower Jurassic	Portugal	Achada Dolomites and Limestones	Subtidal	dip-slip movements causing tsunamis	Kullberg <i>et al.</i> 2001

405

406 **Reference list:**

407 Benner, J.S., Ekdale, A.A. & De Gibert, J.M. 2004. Macroborings (Gastrochaenolites) in

408 Lower Ordovician hardgrounds of Utah: Sedimentologic, paleoecologic, and

409 evolutionary implications. *PALAIOS*, **19**, 543-550, doi: 10.1669/0883-

410 1351(2004)019<0543:mgiloh>2.0.co;2.

- 411 Berg, R.R. 1960. Cambrian and Ordovician history of Colorado *In*: Weimar, R.J. & Haun,  
412 J.D. (eds) *Guide to the geology of Colorado*. Geological Society of America, Rocky  
413 Mountain Association of Geologists, and Colorado Scientific Society, Denver, 10-17
- 414 Bertrand-Sarfati, J. & Moussine-Pouchkine, A. 1983. Platform-to-basin facies evolution: the  
415 carbonates of Late Proterozoic (Vendian) Gourma (West Africa). *Journal of*  
416 *Sedimentary Research*, **53**, 275-293.
- 417 Braun, M. & Friedman, G.M. 1969. Carbonate lithofacies and environment of the Tribes  
418 Hill Formation (Lower Ordovician) of the Mohawk Valley, New York. *Journal of*  
419 *Sedimentary Petrology*, **39**, 113-135.
- 420 Brett, C.E., Liddell, W.D. & Derstler, K.L. 1983. Late Cambrian hard substrate  
421 communities from Montana/Wyoming: the oldest known hardground encrusters  
422 *Lethaia*, **16**, 281-289.
- 423 Calner, M. 2005. A Late Silurian extinction event and anachronistic period. *Geology*, **33**,  
424 305-308.
- 425 Chen, J. 2014. Surface and subsurface reworking by storms on a Cambrian carbonate  
426 platform: evidence from limestone breccias and conglomerates. *Geologos*, **20**, 13-  
427 23.
- 428 Chen, J., Chough, S.K., Chun, S.S. & Han, Z. 2009. Limestone pseudoconglomerates in the  
429 Late Cambrian Gushan and Chaomidian Formations (Shandong Province, China):  
430 soft-sediment deformation induced by storm-wave loading. *Sedimentology*, **56**,  
431 1174-1195.
- 432 Chen, J., Han, Z., Zhang, X., Fan, A. & Yang, R. 2010. Early diagenetic deformation  
433 structures of the Furongian ribbon rocks (Shandong Province, China) – a new  
434 perspective of the genesis of limestone conglomerates. *Science China Earth*  
435 *Sciences* **53**, 241-252.

- 436 Chen, J., Qian, Y., Zhang, J., Lin, Y., Yin, L., Wang, Z.-h., Wang, Z.-z., Yang, J. & Wang,  
437 Y. 1988. The recommended Cambrian-Ordovician global boundary stratotype of the  
438 Xiaoyangqiao section (Dayangcha, Jilin Province), China. *Geological Magazine*,  
439 **125**, 415-444.
- 440 Cherns, L. 1982. Palaeokarst, tidal erosion surfaces and stromatolites in the Silurian Eke  
441 Formation of Gotland, Sweden. *Sedimentology*, **29**, 819-833.
- 442 Cherns, L. 1983. The Hemse-Eke boundary facies relationships in the Ludlow series of  
443 Gotland, Sweden. *Sveriges Geologiska Undersökning, Series C*, **800**, 1-45.
- 444 Choi, Y.S., Kim, J.C. & Lee, Y.I. 1993. Subtidal, flat-pebble conglomerates from the Early  
445 Ordovician Mungok Formation, Korea : origin and depositional process. *Journal of*  
446 *the Geological Society, Korea*, **29**, 15-29.
- 447 Cook, H.E. & Taylor, M.E. 1975. Early Paleozoic continental margin sedimentation,  
448 trilobite biofacies, and the thermocline, western United States. *Geology*, **3**, 559-562.
- 449 Cook, H.E. & Taylor, M.E. 1977. Comparison of continental slope and shelf environments  
450 in the Upper Cambrian and lowest Ordovician of Nevada *In*: Cook, H.E. & Enos, P.  
451 (eds) *Deep-Water Carbonate Environments. Society of Economic Paleontologists*  
452 *and Mineralogists Special Publication* , **25**, 51-81.
- 453 Csonka, J.D. 2009. *Sedimentary dynamics and stratinomy of a Middle Cambrian ichnofossil*  
454 *lagerstätte, Gros Ventre Formation, Wyoming, USA*. MS thesis, Michigan State  
455 University, ProQuest, UMI 1468318, 91 p.
- 456 Dattilo, B.F. 1993. The Lower Ordovician Fillmore Formation of western Utah: storm-  
457 dominated sedimentation on a passive margin. *Brigham Young University Geology*  
458 *Studies*, **39**, 71-100.

459 Day, E.S., James, N.P., Narbonne, G.M. & Dalrymple, R. 2004. A sedimentary prelude to  
460 Marinoan glaciation, Cryogenian (Middle Neoproterozoic) Keele Formation,  
461 Mackenzie Mountains, northwestern Canada. *Precambrian Research*, **133**, 223-247.

462 Demicco, R., V., Spencer, R.J., Waters, B.B. & Cloyd, K.C. 1991. Two-dimensional models  
463 of a Cambrian carbonate shelf deposit. *Kansas Geological Survey Bulletin*, **233** 463-  
464 472.

465 Demicco, R.V. 1985. Platform and off-platform carbonates of the Upper Cambrian of  
466 western Maryland. *Sedimentology*, **32**, 1-22.

467 Dewing, K. & Nowlan, G. 2012. The Lower Cambrian to Lower Ordovician carbonate  
468 platform and shelf margin, Canadian Arctic Islands. *In: Derby, J.R., Fritz, R.D.,*  
469 *Longacre, S.A., Morgan, W.A. & Sternbach, C.A. (eds) The great American*  
470 *carbonate bank: The geology and economic resources of the Cambrian-Ordovician*  
471 *Sauk megasequence of Laurentia. AAPG Memoir*, **98**, 627-647.

472 Ding, Y., Bai, Z., Liu, J. & Han, Z. 2008. Multiple origins of flat-pebble conglomerate and  
473 sedimentary environments of the Gushan Formation at Tangwangzhai in Shandong  
474 Province. *Journal of Palaeogeography*, **10**, 125-138.

475 Dorf, E. & Lochman, C. 1940. Upper Cambrian formations in southern Montana.  
476 *Geological Society of America Bulletin*, **51**, 541-556.

477 Dronov, A., Kanygin, A., Timokhin, A., Tolmacheva, T.Y. & Gonta, T. 2009. Correlation  
478 of eustatic and biotic events in the Ordovician paleobasins of the Siberian and  
479 Russian platforms. *Paleontological Journal*, **43**, 1477-1497.

480 Eoff, J.D. 2014. Sedimentary facies of the upper Cambrian (Furongian; Jiangshanian and  
481 Sunwaptan) Tunnel City Group, Upper Mississippi Valley: New insight on the old  
482 stormy debate. *Sedimentary Geology*, **302**, 102-121, doi:  
483 <http://dx.doi.org/10.1016/j.sedgeo.2013.09.008>.

- 484 Friedman, G.M. 1994. Upper Cambrian-Lower Ordovician (Sauk) platform carbonates of  
485 the northern Appalachian (Gondwana) passive margin. *Carbonates and Evaporites*,  
486 **9**, 143-150.
- 487 Glumac, B. & Walker, K.R. 1997. Selective dolomitization of Cambrian microbial  
488 carbonate deposits: A key to mechanisms and environments of origin. *PALAIOS*, 98-  
489 110.
- 490 Grotzinger, J. & Al-Rawahi, Z. 2014. Depositional facies and platform architecture of  
491 microbialite-dominated carbonate reservoirs, Ediacaran-Cambrian Ara Group,  
492 Sultanate of Oman. *AAPG Bulletin*, **98**, 1453-1494.
- 493 Heubeck, C., Ergaliev, G. & Evseev, S. 2013. Large-scale seismogenic deformation of a  
494 carbonate platform straddling the Precambrian-Cambrian Boundary, Karatau Range,  
495 Kazakhstan. *Journal of Sedimentary Research*, **83**, 1005-1025, doi:  
496 10.2110/jsr.2013.76.
- 497 Ishikawa, T., Ueno, Y., Komiya, T., Sawaki, Y., Han, J., Shu, D., Li, Y., Maruyama, S. &  
498 Yoshida, N. 2008. Carbon isotope chemostratigraphy of a Precambrian/Cambrian  
499 boundary section in the Three Gorge area, South China: prominent global-scale  
500 isotope excursions just before the Cambrian Explosion. *Gondwana Research*, **14**,  
501 193-208.
- 502 Ji, Z. & Barnes, C.R. 1996. Uppermost Cambrian and Lower Ordovician conodont  
503 biostratigraphy of the Survey Peak Formation (Ibexian/Tremadoc), Wilcox Pass,  
504 Alberta, Canada. *Journal of Paleontology*, **70**, 871-890.
- 505 Jones, B. & Dixon, O.A. 1976. Storm deposits in the Reach Bay Formation (upper Silurian),  
506 Somerset Island, Arctic Canada (An application of Markov chain analysis). *Journal*  
507 *of Sedimentary Petrology*, **46**, 393-401.

508 Kazmierczak, J. & Goldring, R. 1978. Subtidal flat-pebble conglomerate from the Upper  
509 Devonian of Poland: a multiprovenant high-energy product. *Geological Magazine*,  
510 **115**, 359-366.

511 Kim, J.C. & Lee, Y.I. 1995. Flat-pebble conglomerate: a characteristic lithology of Upper  
512 Cambrian and Lower Ordovician shallow-water carbonate sequences. *In*: Cooper, J.  
513 D., Droser, M. L. & Finney, S. C. (eds) *Ordovician Odyssey: Short Papers for the*  
514 *Seventh International Symposium on the Ordovician System, 1995. Pacific Section*  
515 *SEPM*, **77**, 371-4.

516 Kouchinsky, A., Bengtson, S., Gallet, Y., Korovnikov, I., Pavlov, V., Runnegar, B., Shields,  
517 G., Veizer, J., Young, E. & Ziegler, K. 2008. The SPICE carbon isotope excursion in  
518 Siberia: a combined study of the upper Middle Cambrian–lowermost Ordovician  
519 Kulyumbe River section, northwestern Siberian Platform. *Geological Magazine*,  
520 **145**, 609-622.

521 Kozub, P. 1997. The origin of flat-pebble conglomerates in the Upper Cambrian of the  
522 Clarks Fork region, Park County, Wyoming. *In*: Mendelson, C.V., Maniewicz, C.M.  
523 (eds), *Tenth Keck Research Symposium in Geology Proceedings. Department of*  
524 *Geology, Beloit College, South Beloit, Illinois*, 134-7.

525 Kruse, P.D. 1996. Update on the northern Australian Cambrian sponges Rankenella,  
526 Jawonya and Wagima. *Alcheringa*, **20**, 161-178, doi: 10.1080/03115519608619188.

527 Kruse, P.D., Gandin, A., Debrenne, F. & Wood, R. 1996. Early Cambrian bioconstructions  
528 in the Zavkhan Basin of western Mongolia. *Geological Magazine*, **133**, 429-444.

529 Kullberg, J.C., OloÂAriz, F., Marques, B., Caetano, P.S. & Roch, R.B. 2001. Flat-pebble  
530 conglomerates: a local marker for Early Jurassic seismicity related to syn-rift  
531 tectonics in the Sesimbra area (Lusitanian Basin, Portugal). *Sedimentary Geology*,  
532 **139** 49-70.

- 533 Kwon, Y., Chough, S., Choi, D. & Lee, D. 2002. Origin of limestone conglomerates in the  
534 Choson Supergroup (Cambro–Ordovician), mid-east Korea. *Sedimentary Geology*,  
535 **146**, 265-283.
- 536 Landing, E., Westrop, S.R. & Keppie, J.D. 2007. Terminal Cambrian and lowest Ordovician  
537 succession of Mexican West Gondwana: biotas and sequence stratigraphy of the  
538 Tiñu Formation. *Geological Magazine*, **144**, 909-936.
- 539 Lee, J.-H., Chen, J. & Chough, S. 2010. Paleoenvironmental implications of an extensive  
540 maceriate microbialite bed in the Furongian Chaomidian Formation, Shandong  
541 Province, China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **297**, 621-  
542 632.
- 543 Lee, Y.I. & Kim, J.C. 1992. Storm-influenced siliciclastic and carbonate ramp deposits, the  
544 Lower Ordovician Dumugol Formation, South Korea. *Sedimentology*, **39**, 951-969.
- 545 Liang, C.M., Friedman, G.M. & Zheng, Z.C. 1993. Carbonate storm deposits (tempestites)  
546 of Middle to Upper Cambrian age in the Helan Mountains, northwest China.  
547 *Carbonates and Evaporites*, **8**, 181-190.
- 548 Liu, J. & Zhan, R. 2009. Temporal distribution of diagnostic biofabrics in the Lower and  
549 Middle Ordovician in North China: clues to the geobiology of the Great Ordovician  
550 Biodiversification Event. *Acta Geologica Sinica*, **83**, 513-523.
- 551 Liu, J., Zhan, R., Dai, X., Liao, H., Ezaki, Y. & Adachi, N. 2011. Demise of Early  
552 Ordovician oolites in South China: Evidence for paleoceanographic changes before  
553 the GOBE. In: Gutiérrez-Marco, J. C., Rábano, I. & Garcia-Bellido, D. (eds)  
554 *Ordovician of the World*. Cuadernos del Museo Geominero, **14**, 309-317.
- 555  
556 Luo, G., Junium, C.K., Kump, L.R., Huang, J., Li, C., Feng, Q., Shi, X., Bai, X. & Xie, S.  
557 2014. Shallow stratification prevailed for ~ 1700 to ~ 1300 Ma ocean: Evidence from

558 organic carbon isotopes in the North China Craton. *Earth and Planetary Science*  
559 *Letters*, **400**, 219-232.

560 MacNaughton, R. B., Roots, C.F. & Martel, E. 2008. Neoproterozoic-(?) Cambrian  
561 lithostratigraphy, northeast Sekwi Mountain map area, Mackenzie Mountains,  
562 Northwest Territories: new data from measured sections. *Geological Survey of*  
563 *Canada, Current Research 2008-16*, 17p.

564 Markello, J.R. & Read, J.F. 1981. Carbonate ramp-to-deeper shale shelf transitions of an  
565 Upper Cambrian intrashelf basin, Nolichucky Formation, Southwest Virginia  
566 Appalachians. *Sedimentology*, **28**, 573-597.

567 Markello, J.R. & Read, J.F. 1982. Upper Cambrian Intrashelf Basin, Nolichucky Formation,  
568 Southwest Virginia Appalachians. *AAPG Bulletin*, **66** 860-878.

569 Meng, X., Ge, M. & Tucker, M.E. 1997. Sequence Sequence stratigraphy, sea-level changes  
570 and depositional systems in the Cambro-Ordovician of the North China carbonate  
571 platform. *Sedimentary Geology*, **114**, 189-222.

572 Mount, J.F. & Kidder, D. 1993. Combined flow origin of edgewise intraclast  
573 conglomerates: Sellick Hill Formation (Lower Cambrian), South Australia.  
574 *Sedimentology*, **40**, 315-329.

575 Myrow, P., M., Tice, L., Archuleta, B., Clark, B., Taylor, J.F. & Ripperdan, R.L. 2004.  
576 Flat-pebble conglomerate: its multiple origins and relationship to metre-scale  
577 depositional cycles. *Sedimentology*, **51**, 973-996.

578 Myrow, P.M., Taylor, J.F., Runkel, A.C. & Ripperdan, R.L. 2012. Mixed siliciclastic-  
579 carbonate upward-deepening cycles of the Upper Cambrian inner detrital belt of  
580 Laurentia. *Journal of Sedimentary Research*, **82**, 216-231, doi: 10.2110/jsr.2012.20.

581 Narbonne, G.M., Saylor, B.Z. & Grotzinger, J.P. 1997. The youngest Ediacaran fossils from  
582 southern Africa. *Journal of Paleontology*, **71**, 953-967.



- 583 Pope, A. 1990. The geology and mineral deposits of the Toby-Horsethief Creek map area,  
584 northern Purcell Mountains, southeast British Columbia (82K). *British Columbia*  
585 *Geological Survey, Mineral Resources Division, Geological Survey Branch, Open*  
586 *File 1990-26*, 57p.
- 587 Popov, L.E., Holmer, L.E. & Miller, J.F. 2002. Lingulate brachiopods from the Cambrian-  
588 Ordovician boundary beds of Utah. *Journal of Paleontology*, **76**, 211-228.
- 589 Pratt, B.R. 2002. Storms versus tsunamis: Dynamic interplay of sedimentary, diagenetic,  
590 and tectonic processes in the Cambrian of Montana. *Geology*, **30** , 423-426.
- 591 Pratt, B.R. & Bordonaro, O.L. 2007. Tsunamis in a stormy sea: Middle Cambrian inner-  
592 shelf limestones of western Argentina. *Journal of Sedimentary Research*, **77**, 256-  
593 262.
- 594 Pruss, S.B., Bottjer, D.J., Corsetti, F.A. & Baud, A. 2006. A global marine sedimentary  
595 response to the end-Permian mass extinction: examples from southern Turkey and  
596 the western United States. *Earth-Science Reviews*, **78**, 193-206.
- 597 Pruss, S.B., Corsetti, F.A. & Bottjer, D.J. 2005. The unusual sedimentary rock record of the  
598 Early Triassic: a case study from the southwestern United States. *Palaeogeography*,  
599 *Palaeoclimatology, Palaeoecology*, **222**, 33-52.
- 600 Pruss, S.B., Finnegan, S., Fischer, W.W. & Knoll, A.H. 2010. Carbonates in skeleton-poor  
601 seas: new insights from Cambrian and Ordovician strata of Laurentia. *PALAIOS*, **25**,  
602 73-84, doi: <http://dx.doi.org/10.2110/palo.2009.p09-101r>.
- 603 Robison, R.A. 1964. Upper Middle Cambrian stratigraphy of western Utah. *Geological*  
604 *Society of America Bulletin*, **75**, 995-1010.
- 605 Saltzman, M.R. 1999. Upper Cambrian carbonate platform evolution, Elvinia and  
606 Taenicephalus zones (Pterocephaliid—Ptychaspid biomere boundary), Northwestern  
607 Wyoming. *Journal of Sedimentary Research*, **69** 926-938.

608 Saltzman, M.R., Runnegar, B. & Lohmann, K.C. 1998. Carbon isotope stratigraphy of  
609 Upper Cambrian (Steptoean Stage) sequences of the eastern Great Basin: Record of  
610 a global oceanographic event. *Geological Society of America Bulletin*, **110**, 285-297.

611 Sepkoski, J.J.J. 1982. Flat-pebble conglomerates, storm deposits, and the Cambrian bottom  
612 fauna. *In*: Einsele, G. & Seilacher, A. (eds) *Cyclic and Event Stratification*.  
613 Springer-Verlag, 371-386.

614 Shapiro, R.S. & Awramik, S.M. 2006. *Favosamaceria cooperi* new group and form: a  
615 widely dispersed, time-restricted thrombolite. *Journal of Paleontology*, **80**, 411-422.

616 Sprinkle, J. & Guensburg, T.E. 1995. Origin of echinoderms in the Paleozoic evolutionary  
617 fauna: The role of substrates. *PALAIOS*, **10**, 437-453.

618 Van Loon, A., Han, Z. & Han, Y. 2012. Slide origin of breccia lenses in the Cambrian of the  
619 North China Platform: new insight into mass transport in an epeiric sea. *Geologos*,  
620 **18**, 223-235.

621 Varlamov, A., Pak, K. & Rosova, A. 2006. The Upper Cambrian of the Chopko River  
622 section, Norilsk region, northwestern Siberian platform: Stratigraphy and trilobites.  
623 *Paleontological Journal*, **40**, S1-S56.

624 Westrop, S.R. 1989. Facies anatomy of an Upper Cambrian grand cycle: Bison Creek and  
625 Mistaya formations, southern Alberta. *Canadian Journal of Earth Sciences*, **26**,  
626 2292-2304.

627 Whisonant, R. 1987. Paleocurrent and petrographic analysis of imbricate intraclasts in  
628 shallow-marine carbonates, Upper Cambrian, southwestern Virginia. *Journal of*  
629 *Sedimentary Petrology*, **57**, 983-994.

630 Wignall, P.B. & Twitchett, R.J. 1999. Unusual intraclastic limestones in Lower Triassic  
631 carbonates and their bearing on the aftermath of the end-Permian mass extinction.  
632 *Sedimentology*, **46**, 303-316.

633 Wilson, M.A., Palmer, T.J., Guensburg, T.E., Finton, C.D. & Kaufman, L.E. 1992. The  
634 development of an Early Ordovician hard ground community in response to rapid  
635 sea-floor calcite precipitation. *Lethaia*, **25**, 19-34.

636 Wilson, M.D. 1985. Origin of Upper Cambrian flat pebble conglomerates in the northern  
637 Powder River Basin. *In*: Longman, M.W., Shanley, K.W., Lindsay, R.F. & Eby,  
638 D.E. (eds) *Rocky Mountain carbonate reservoirs*. *SEPM Core Workshop*, **7**, 1-50.

639 Woods, A.D. 2009. Anatomy of an anachronistic carbonate platform: Lower Triassic  
640 carbonates of the southwestern United States. *Australian Journal of Earth Sciences*,  
641 **56**, 825-839, doi: 10.1080/08120090903002649.

642 Yang, Z., Otofujii, Y.-i., Sun, Z. & Huang, B. 2002. Magnetostratigraphic constraints on the  
643 Gondwanan origin of North China: Cambrian/Ordovician boundary results.  
644 *Geophysical Journal International*, **151**, 1-10.

645

646