

# SCIENTIFIC REPORTS



OPEN

## Targeted suppression of autoreactive CD8<sup>+</sup> T-cell activation using blocking anti-CD8 antibodies

Received: 08 April 2016  
Accepted: 09 September 2016  
Published: 17 October 2016

Mathew Clement<sup>1</sup>, James A. Pearson<sup>1</sup>, Stephanie Gras<sup>2,3</sup>, Hugo A. van den Berg<sup>4</sup>, Anya Lissina<sup>5</sup>, Sian Llewellyn-Lacey<sup>1</sup>, Mark D. Willis<sup>6</sup>, Tamsin Dockree<sup>1</sup>, James E. McLaren<sup>1</sup>, Julia Ekeruche-Makinde<sup>7</sup>, Emma Gostick<sup>1</sup>, Neil P. Robertson<sup>6</sup>, Jamie Rossjohn<sup>1,2,3</sup>, Scott R. Burrows<sup>8</sup>, David A. Price<sup>1,9</sup>, F. Susan Wong<sup>1</sup>, Mark Peakman<sup>10,\*</sup>, Ania Skowera<sup>10,\*</sup> & Linda Wooldridge<sup>5,\*</sup>

CD8<sup>+</sup> T-cells play a role in the pathogenesis of autoimmune diseases such as multiple sclerosis and type 1 diabetes. However, drugs that target the entire CD8<sup>+</sup> T-cell population are not desirable because the associated lack of specificity can lead to unwanted consequences, most notably an enhanced susceptibility to infection. Here, we show that autoreactive CD8<sup>+</sup> T-cells are highly dependent on CD8 for ligand-induced activation via the T-cell receptor (TCR). In contrast, pathogen-specific CD8<sup>+</sup> T-cells are relatively CD8-independent. These generic differences relate to an intrinsic dichotomy that segregates self-derived and exogenous antigen-specific TCRs according to the monomeric interaction affinity with cognate peptide-major histocompatibility complex class I (pMHC I). As a consequence, “blocking” anti-CD8 antibodies can suppress autoreactive CD8<sup>+</sup> T-cell activation in a relatively selective manner. These findings provide a rational basis for the development and *in vivo* assessment of novel therapeutic strategies that preferentially target disease-relevant autoimmune responses within the CD8<sup>+</sup> T-cell compartment.

CD8<sup>+</sup> T-cells recognise cognate peptide-major histocompatibility complex class I (pMHC I) antigens via the clonotypically-expressed  $\alpha\beta$  T-cell receptor (TCR) and the lineage-specific CD8 coreceptor<sup>1,2</sup>. The TCR engages the  $\alpha 1/\alpha 2$  domain peptide-binding platform of pMHC I, thereby dictating antigen specificity<sup>3</sup>. In contrast, CD8 binds at a spatially distinct and largely conserved site formed by the  $\alpha 3$  domain of the MHC I heavy chain with a contribution from  $\beta 2$ -microglobulin, an event that acts functionally to enhance antigen sensitivity<sup>4,5</sup>. Several mechanisms are involved in this latter phenomenon, including: (i) stabilisation of the TCR/pMHC I interaction<sup>6,7</sup>; (ii) recruitment of essential signalling molecules to the intracellular side of the TCR/CD3/ $\zeta$  complex<sup>8–11</sup>; and (iii) localisation of the TCR/pMHC I complex within membrane micro-domains that form privileged sites for the initiation of TCR-mediated signalling<sup>12,13</sup>. This allows the CD8 coreceptor to fine-tune antigen-specific responses within the CD8<sup>+</sup> T-cell compartment.

It has become increasingly evident in recent years that CD8<sup>+</sup> T-cells play a key role in the pathogenesis of autoimmune diseases such as type 1 diabetes (T1D)<sup>14–16</sup> and multiple sclerosis (MS)<sup>17–19</sup>. As such, there is a strong rationale for developing therapeutic strategies that target the autoreactive CD8<sup>+</sup> T-cell population<sup>20–22</sup>. Previous studies have employed antibodies directed against T-cell surface markers (CD3, CD4 and

<sup>1</sup>Division of Infection and Immunity, Cardiff University, Cardiff CF14 4XN, UK. <sup>2</sup>Infection and Immunity Program and Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute, Monash University, Clayton, Victoria 3800, Australia. <sup>3</sup>Australian Research Council Centre of Excellence for Advanced Molecular Imaging, Monash University, Clayton, VIC 3800, Australia. <sup>4</sup>Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK. <sup>5</sup>Faculty of Health Sciences, University of Bristol, Bristol BS8 1TD, UK. <sup>6</sup>Division of Psychological Medicine and Clinical Neuroscience, Cardiff University, Cardiff CF14 4XN, UK. <sup>7</sup>Mucosal Infection and Immunity Group, Department of Medicine, Imperial College London, London SW7 2AZ, UK. <sup>8</sup>QIMR Berghofer Medical Research Institute, Brisbane, QLD 4029, Australia. <sup>9</sup>Vaccine Research Center, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA. <sup>10</sup>Department of Immunobiology, King's College London, London SE1 9RT, UK. \*These authors contributed equally to this work. Correspondence and requests for materials should be addressed to M.C. (email: clementm@cardiff.ac.uk)

CD8) to induce tolerance in mice<sup>20,21,23,24</sup>, although to date, it has not been possible to translate strategies using “tolerance-inducing antibodies” into humans<sup>25</sup>. However, there are important biological differences between autoreactive and pathogen-specific CD8<sup>+</sup> T-cells that may be amenable to therapeutic exploitation. Extensive biophysical analyses have shown that pathogen-specific TCRs typically engage cognate pMHCI with high monomeric affinities (range  $K_D \sim 1\text{--}50\ \mu\text{M}$ )<sup>3,26</sup>. In contrast, autoreactive TCRs that escape negative selection display markedly lower monomeric affinities for pMHCI ( $K_D > 100\ \mu\text{M}$ )<sup>27–30</sup>. Autoimmune disease-relevant TCR/pMHCI interactions may even occur at  $K_D$  values  $> 200\ \mu\text{M}$ <sup>31,32</sup>. Importantly, CD8<sup>+</sup> T-cells bearing such low affinity TCRs are highly dependent on CD8 for cognate ligand-induced activation<sup>33,34</sup>. On the basis of these observations, we hypothesised that CD8-targeted strategies could be used to inhibit autoreactive CD8<sup>+</sup> T-cells in a relatively selective manner.

Anti-CD8 monoclonal antibodies have been used widely to study the functional role of the CD8 coreceptor<sup>35,36</sup>. To phenotype such antibodies we have defined the following criteria: (1) effect on pMHCI tetramer staining, (2) effect on pMHCI specific activation; and, (3) ability to trigger non-specific activation (i.e. to elicit effector function in the absence of TCR/pMHCI engagement)<sup>37</sup>. We have observed that considerable heterogeneity exists between different anti-CD8 antibodies. In general, anti-CD8 antibodies can inhibit or enhance pMHCI tetramer binding, which is mirrored by their effect on pMHCI antigen-specific activation<sup>35</sup>. It is notable that the anti-mouse CD8 antibody YTS105.18 has been used in previous studies to reverse T1D in non-obese diabetic (NOD) mice<sup>20,21</sup>. This clone does not inhibit pMHCI binding or antigen-specific CD8<sup>+</sup> T-cell activation and therefore cannot be classified as a “blocking” antibody<sup>38</sup>. In contrast, we have selected an anti-human CD8 antibody (DK25) that exhibits a potent blocking phenotype. DK25 inhibits pMHCI tetramer binding at the cell surface, inhibits pMHCI antigen specific T-cell activation but does not trigger non-specific activation<sup>35,37</sup>.

We show that autoreactive CD8<sup>+</sup> T-cells are preferentially inhibited by blocking anti-CD8 antibodies as a consequence of low affinity TCR/pMHCI interactions that confer an intrinsic dependence on the CD8 coreceptor for ligand-induced activation via the TCR. Our findings suggest novel strategies for the treatment of autoimmune diseases without the attendant side effects that complicate generalised immunosuppression.


## Results

**Autoreactive CD8<sup>+</sup> T-cells expressing low affinity TCRs are highly CD8-dependent.** A primary aim of this study was to test the hypothesis that autoreactive CD8<sup>+</sup> T-cells are highly dependent on CD8 for ligand-induced activation via the TCR. In preliminary experiments, we made use of the well-characterised CD8<sup>+</sup> T-cell clone 1E6, which is specific for the HLA-A\*0201-restricted preproinsulin (PPI) epitope ALWGPDPAAA (ALW)<sup>39</sup>. Despite a low affinity monomeric interaction between the 1E6 TCR and cognate pMHCI ( $K_D = 278\ \mu\text{M}$ ), this clone recognises and destroys human pancreatic  $\beta$ -cells *in vitro*<sup>32,39</sup>. To examine the CD8 dependency of 1E6, we measured effector outputs in response to titrated doses of the cognate peptide ligand presented by C1R-A\*0201 or C1R-A\*0201 D227K/T228A cells. In the absence of pMHCI/CD8 engagement due to the HLA-A\*0201 D227K/T228A mutation, interferon (IFN)- $\gamma$  production by the 1E6 CD8<sup>+</sup> T-cell clone was completely abrogated (Fig. 1A). Macrophage inflammatory protein (MIP)-1 $\beta$  release was similarly compromised at physiologically relevant peptide concentrations (Fig. 1B).


Next, we performed similar experiments with the CD8<sup>+</sup> T-cell clone ALF8, which recognises the HLA-A\*0201-restricted influenza A virus matrix protein (M1) epitope GILGFVFTL. The ALF8 TCR engages cognate pMHCI with a monomeric affinity ( $K_D = 7.71\ \mu\text{M}$ ; Table 1 and Supplementary Fig. S1) that falls within the typical range for pathogen-specific TCRs ( $K_D = 1\text{--}50\ \mu\text{M}$ )<sup>3,26</sup>. ALF8 exhibited functional responses at substantially lower concentrations of exogenous peptide compared with 1E6. In the absence of an intact pMHCI/CD8 interaction, ALF8 was still capable of efficient production of IFN- $\gamma$  and MIP-1 $\beta$ , even at low peptide concentrations (Fig. 1C,D). Similar results were obtained with the CD8<sup>+</sup> T-cell clone MCNLV, which is specific for the HLA-A\*0201-restricted human cytomegalovirus (HCMV) pp65 epitope NLVPMVATV (Fig. 1E,F). These data suggest that autoreactive CD8<sup>+</sup> T-cells expressing low affinity TCRs are less antigen sensitive and more dependent on CD8 coreceptor engagement as compared to pathogen-specific CD8<sup>+</sup> T-cells.

**CD8 has a profound effect on the number of peptides that can be recognised by autoreactive CD8<sup>+</sup> T-cells.** In an earlier study, we showed that the 1E6 clone is highly cross-reactive in the context of HLA-A\*0201, recognising  $\sim 1$  million different peptide ligands at physiologically relevant concentrations<sup>40</sup>. As part of this earlier study we performed a 10mer combinatorial peptide library (CPL) scan of the 1E6 clone using C1R-A\*0201 targets (WT MHCI/CD8) to present each peptide mixture<sup>40</sup>. To extend these findings, we performed a 10mer combinatorial peptide library (CPL) scan of 1E6 using C1R-A\*0201 D227K/T228A targets (Null MHCI/CD8) (Fig. 2 and Supplementary Fig. S2). The 1E6 clone is highly promiscuous in the presence of an intact pMHCI/CD8 interaction (Fig. 2). However, in the absence of CD8 engagement, very few responses were observed (Fig. 2 and Supplementary Fig. S2). This is consistent with a previous study that demonstrates that CD8 can control levels of T-cell crossreactivity<sup>34</sup>. Abrogation of the pMHCI/CD8 interaction eliminates promiscuous pMHCI recognition of the 1E6 TCR.

**Selection of a potent anti-CD8 antibody that blocks pMHCI tetramer binding at low TCR/pMHCI affinities.** It is established that anti-CD8 antibodies display various phenotypes<sup>35,37</sup>. In particular, SK1 and DK25 can be classified as blocking anti-CD8 antibodies because they inhibit pMHCI binding at the cell surface. Consistent with this trait, both SK1 and DK25 blocked pMHCI tetramer staining of the 1E6 clone in a concentration-dependent manner (Fig. 3A,B). In contrast, pMHCI tetramer staining of the ALF8 clone was minimally disrupted by the same antibodies across an identical concentration range (Fig. 3C,D). These results suggest that anti-CD8 antibody-mediated blockade of pMHCI binding occurs selectively in the presence of a weak TCR/



**Figure 1. Autoreactive CD8<sup>+</sup> T-cells expressing low affinity TCRs are highly CD8-dependent.**  $6 \times 10^4$  C1R-A\*0201 (WT MHC1/CD8) or C1R-A\*0201 D227K/T228A (Null MHC1/CD8) target cells were pulsed with cognate peptide at the indicated concentrations in duplicate for 1 hour at 37 °C.  $3 \times 10^4$  clonal 1E6 (A,B), ALF8 (C,D) or MCNLV (E,F) CD8<sup>+</sup> T-cells were then added and the plates were incubated for 4 hours at 37 °C. Supernatants were assayed for IFN- $\gamma$  and MIP-1 $\beta$  by ELISA. The mean  $\pm$  SD of two replicate assays is shown. Data is representative of three independent experiments.




**Figure 2. CD8 has a profound effect on the number of peptides that can be recognised by the diabetogenic CD8<sup>+</sup> T-cell clone 1E6.**  $6 \times 10^4$  C1R-A\*0201 (WT MHC1/CD8) or C1R-A\*0201 D227K/T228A (Null MHC1/CD8) target cells were pulsed in duplicate with each mixture from a 10mer combinatorial peptide library (100  $\mu$ M) for 2 hours at 37 °C.  $3 \times 10^4$  clonal 1E6 CD8<sup>+</sup> T-cells were then added and the plates were incubated overnight at 37 °C. Supernatants were assayed for MIP-1 $\beta$  by ELISA. The primary screens with wildtype (WT) targets were reported previously<sup>40</sup>; the primary screens with D227K/T228A (Null) targets are shown in Supplementary Fig. S2. A box plot summary is presented with the index sequence at the bottom. Data is representative of two independent experiments.

TCR	$K_{Deq}$ ( $\mu$ M)	$k_{on}$ ( $10^4$ /Ms)	$k_{off}$ (1/s)	$K_{Dcalc}$ ( $\mu$ M)	$t_{1/2}$ (s)
ALF3	$5.84 \pm 0.10$	$2.19 \pm 0.44$	$0.128 \pm 0.007$	$7.33 \pm 0.40$	7.8
ALF8	$7.71 \pm 0.17$	$2.89 \pm 0.29$	$0.158 \pm 0.005$	$5.48 \pm 0.39$	6.3

**Table 1. Affinities and kinetics of GILGFVFTL-HLA-A\*0201 binding to the ALF3 and ALF8 TCRs.**  $K_{Deq}$  is the measured equilibrium affinity,  $k_{on}$  is the association rate,  $k_{off}$  is the dissociation rate,  $t_{1/2}$  is the half-life ( $0.693/k_{off}$ ) and  $K_{Dcalc}$  is the calculated affinity based on kinetic parameters.

pmHCI interaction. Further experiments designed to extend this observation were conducted with DK25, which exhibited more potent blocking activity in side-by-side comparisons with SK1 (Fig. 3A–D).

**DK25 blocks the activation of CD8<sup>+</sup> T-cells at low TCR/pmHCI affinities.** To test the hypothesis that DK25 can be used as a selective agent to target CD8<sup>+</sup> T-cells bearing low affinity TCRs, we again employed the




**Figure 3. Anti-CD8 antibodies can block pMHC I tetramer binding at low TCR/pMHC I affinities.**  $5 \times 10^4$  clonal 1E6 (A,B) or ALF8 (C,D) CD8<sup>+</sup> T-cells were pre-treated with or without the indicated anti-human CD8 antibodies at various concentrations for 20 minutes at 4 °C and then stained with 25 μg/ml of cognate pMHC I tetramer for 15 minutes at 37 °C. The column graphs depict percent maximal tetramer staining (A,C) or percent inhibition of tetramer staining (B,D). Data is representative of three independent experiments.

1E6 clone and examined activation in response to both the cognate ligand (ALW) and the altered peptide variant RQWGPDPAAV (RQW), which binds the 1E6 TCR with a substantially higher monomeric affinity ( $K_D = 10 \mu\text{M}$ )<sup>41</sup>. This monoclonal system allowed a rigorous evaluation of DK25-mediated effects as a function of a single variable parameter, namely the affinity of the TCR/pMHC I interaction. In cytotoxicity assays, the 1E6 clone killed target cells presenting either ALW or RQW at levels proportional to the corresponding TCR/pMHC I affinities (Fig. 4A). This is consistent with a recent publication that demonstrates a strong correlation between 1E6 T-cell antigen sensitivity and TCR/pMHC I affinity<sup>42</sup>. Interestingly, there is no correlation between ligand potency and pMHC I stability in this system ( $T_m$  for A2-ALW and A2-RQW was 60 °C and 54 °C, respectively)<sup>42</sup>. The addition of DK25 at low concentrations (0.5 or 1 μg/ml) abolished the killing response to target cells presenting ALW (Fig. 4B). In contrast, the same concentrations of DK25 failed to abolish 1E6-mediated killing of target cells presenting RQW (Fig. 4C). These data suggest that DK25 selectively inhibits CD8<sup>+</sup> T-cell activation in the presence of low affinity TCR/pMHC I interactions.

**DK25 selectively blocks autoreactive CD8<sup>+</sup> T-cell activation.** Next, we evaluated the ability of DK25 to inhibit autoreactive and pathogen-specific CD8<sup>+</sup> T-cell clones in parallel activation assays. Complete blockade of MIP-1β release by the 1E6 clone in response to the cognate peptide ALW was observed in the presence of DK25 at all concentrations tested (0.25, 0.5 and 1 μg/ml; Fig. 5A). In contrast, the same concentrations of DK25 minimally impacted the activation of ALF3, a CD8<sup>+</sup> T-cell clone specific for the HLA-A\*0201-restricted influenza A virus M1 epitope GILGFVFTL (Fig. 5D). Consistent with this observation, the ALF3 TCR engaged cognate pMHC I with a high monomeric affinity ( $K_D = 5.84 \mu\text{M}$ ; Table 1 and Supplementary Fig. S1).


We conducted similar experiments across a range of different epitope specificities and HLA class I restriction elements. The autoreactive CD8<sup>+</sup> T-cell clones 3F2 and 4C6 recognise the HLA-A\*0201-restricted PPI epitope ALW and the HLA-A\*2402-restricted PPI epitope LWMRLPLL, respectively. In both cases, MIP-1β release was completely inhibited by DK25 across all cognate peptide concentrations tested (Fig. 5B,C). The pathogen-specific CD8<sup>+</sup> T-cell clones SB27 and MCNLV recognise the HLA-B\*3508-restricted Epstein-Barr virus (EBV) BZLF1 epitope LPEPLPQGQLTAY and the HLA-A\*0201-restricted HCMV pp65 epitope NLVPMVATV, respectively. In both cases, robust peptide-specific activation was observed in the presence of DK25 (Fig. 5E,F). The SB27 TCR has been well characterised previously and binds cognate pMHC I with a high monomeric affinity ( $K_D = 9.9 \mu\text{M}$ )<sup>43,44</sup>. Comparable results were obtained for all clones in cytotoxicity assays (Fig. 6A–F). These differences in susceptibility to inhibition by DK25, comparing grouped autoreactive versus pathogen-specific CD8<sup>+</sup> T-cells, were significant at the 5% level (Wilcoxon-Mann-Whitney rank sum test). Anti-CD8 antibodies can therefore be used to block the activation of autoreactive CD8<sup>+</sup> T-cells in a selective manner regardless of epitope specificity or HLA class I restriction.

**DK25 inhibits pancreatic β-cell killing by PPI-specific CD8<sup>+</sup> T-cells.** To examine the utility of anti-CD8 antibody-mediated blockade in a more physiological setting, we tested the ability of DK25 to



**Figure 4.** DK25 blocks the activation of CD8<sup>+</sup> T-cells at low TCR/pMHCII affinities.  $2 \times 10^3$  <sup>51</sup>Cr-labeled C1R-A\*0201 target cells were peptide-pulsed with either ALWGPDPA AAA (A,B) or RQWGPDPA AV (A,C) at the indicated concentrations for 1 hour at 37 °C.  $1 \times 10^4$  clonal 1E6 CD8<sup>+</sup> T-cells were then added with or without the anti-human CD8 antibody DK25 at a final concentration of 0.5 or 1 µg/ml. Cytotoxicity was determined after 4 hours as described in the Materials and Methods. The mean  $\pm$  SD of three replicate assays is shown. Data is representative of three independent experiments.

inhibit pancreatic  $\beta$ -cell killing by the 1E6 clone. In preliminary experiments, we made use of the surrogate  $\beta$ -cell line K562-PPI-A\*0201, created by transfecting K562 leukemia cells with PPI and HLA-A\*0201. DK25 blocked 1E6-mediated killing of K562-PPI-A\*0201 cells at concentrations  $>0.2$  µg/ml (Fig. 7A), consistent with the exogenous peptide titration data (Figs 5 and 6). We then performed a similar experiment using pancreatic  $\beta$ -cells from an HLA-A\*0201<sup>+</sup> donor (Fig. 7B). In the absence of DK25, pancreatic  $\beta$ -cells were readily killed by 1E6. The CD8<sup>+</sup> T-cell clone NLV2, which was derived from a T1D patient as a control and recognises the HLA-A\*0201-restricted HCMV pp65 epitope NLVPMVATV, displayed no direct lytic activity in the same assay. Pancreatic  $\beta$ -cell killing by 1E6 was inhibited by DK25 in a dose-dependent manner, reaching background levels at antibody concentrations  $\geq 0.5$  µg/ml. In contrast, DK25 had no effect on the ability of NLV2 to kill cognate peptide-pulsed HLA-A\*0201<sup>+</sup> pancreatic  $\beta$ -cells. There was a highly significant dependence on DK25 concentration for 1E6 killing of pancreatic  $\beta$ -cells (P-value = 0.0078), whereas the dependence on DK25 concentration for NLV2 was non-significant (NLV2 vs pancreatic  $\beta$ -cells: P = 0.52 and NLV2 vs NLV-pulsed pancreatic  $\beta$ -cells:




**Figure 5. DK25 selectively blocks autoreactive CD8<sup>+</sup> T-cell activation.**  $6 \times 10^4$  C1R-A\*0201 target cells were peptide-pulsed with ALWGPDPAAA (A,B), GILGFVFTL (D) or NLVPMVATV (F) at the indicated concentrations for 1 hour at 37 °C. In parallel,  $6 \times 10^4$  K562-A\*2402 or C1R-B\*3508 target cells were peptide-pulsed similarly with LWMRLLPLL (C) or LPEPLPQGQLTAY (E), respectively.  $3 \times 10^4$  clonal 1E6 (A), 3F2 (B), 4C6 (C), ALF3 (D), SB27 (E) or MCNLV (F) CD8<sup>+</sup> T-cells pre-treated for 30 minutes at 37 °C with the anti-human CD8 antibody DK25 at the indicated concentrations were then added and the plates were incubated for 4 hours at 37 °C. Supernatants were assayed for MIP-1 $\beta$  by ELISA. The mean  $\pm$  SD of two replicate assays is shown. Data is representative of two independent experiments.

$P = 0.28$ ). Therefore, relatively low concentrations of DK25 can therefore be used to inhibit CD8<sup>+</sup> T-cell-mediated pancreatic  $\beta$ -cell destruction without impacting pathogen-specific immunity.

**DK25 efficiently blocks the activation of myelin-specific CD8<sup>+</sup> T-cells.** To extend our study to other autoimmune targets, we examined the effect of DK25 on the activation of BW58 cells expressing human CD8 $\alpha$   $\beta$  and the 2D1 TCR, which is specific for the HLA-A\*0301-restricted myelin proteolipid protein (PLP) epitope KLIETYFSK. The 2D1 TCR plays a role in the induction of a multiple sclerosis-like disease in humanised mice expressing HLA-A\*0301<sup>45</sup>. DK25 efficiently blocked peptide-induced interleukin (IL)-2 production by the BW58 cell line at concentrations of 0.5 and 1  $\mu$ g/ml (Fig. 8A,B). Incomplete but nonetheless potent suppression was observed at an antibody concentration of 0.25  $\mu$ g/ml. Anti-CD8 antibody-mediated blockade of autoreactive CD8<sup>+</sup> T-cells is therefore not restricted by tissue specificity.

**Blocking anti-mouse CD8 antibodies inhibit autoreactive CD8<sup>+</sup> T-cell activation.** Two previous studies have used anti-CD8 antibodies to prevent and reverse T1D in mice<sup>20,21</sup>. In both cases, the investigators relied principally on the anti-mouse CD8 antibody YTS105.18. Shore *et al.* demonstrated that YTS105.18 does not block pMHC I tetramer binding or CD8<sup>+</sup> T-cell activation<sup>38</sup>. We confirmed this observation by showing that




**Figure 6.** DK25 selectively blocks autoreactive CD8<sup>+</sup> T-cell killing.  $2 \times 10^5$  Cr-labeled C1R-A\*0201 (A,B,D,F), K562-A\*2402 (C) or C1R-B\*3508 (E) target cells were peptide-pulsed with ALWGPDPAAA (A,B), LWMRLPLL (C), GILGFVFTL (D), LPEPLPQGQLTAY (E) or NLVPMVATV (F) at the indicated concentrations for 1 hour at 37 °C.  $1 \times 10^4$  clonal 1E6 (A), 3F2 (B), 4C6 (C), ALF3 (D), SB27 (E) or MCNLV (F) CD8<sup>+</sup> T-cells were then added with or without the anti-human CD8 antibody DK25 at a final concentration of 0.25, 0.5 or 1 µg/ml. Cytotoxicity was determined after 4 hours as described in the Materials and Methods. The mean  $\pm$  SD of three replicate assays is shown and is representative of two independent experiments.

murine CD8<sup>+</sup> T-cells expressing the G9C8 TCR, which is specific for the H-2K<sup>d</sup>-restricted insulin B chain epitope LYLVCGERG, exhibit robust activation in the presence of YTS105.18 (Fig. 9A). The blocking approach presented in this study is therefore distinct from the phenomena described in these earlier reports.

Next, we tested the anti-mouse CD8 antibody CT-CD8a, which inhibits pMHC I tetramer binding and CD8<sup>+</sup> T-cell activation<sup>35,37</sup>. The G9C8 TCR binds cognate pMHC I with a low monomeric affinity ( $K_D = 286 \mu\text{M}$ )<sup>46</sup>. In line with our observations in human systems, CT-CD8a efficiently blocked peptide-induced activation of G9C8-expressing CD8<sup>+</sup> T-cells (Fig. 9B). However, CT-CD8a cannot be used therapeutically *in vivo* because it triggers non-specific CD8<sup>+</sup> T-cell activation with exposure times  $\geq 18$  hours<sup>37</sup>. To date, it has not been possible to identify a blocking anti-mouse CD8 antibody with a phenotype akin to DK25, which may be due to intrinsic biophysical and structural differences between mice and humans with respect to the pMHC I/CD8 interaction<sup>47</sup>.

## Discussion

Over the past decade, a substantial body of evidence has accumulated to implicate CD8<sup>+</sup> T-cells as key players in the pathogenesis of common autoimmune diseases such as T1D<sup>14–16,48,49</sup>, MS<sup>17,20,50</sup> and psoriasis<sup>51</sup>. Autoreactive TCRs typically engage cognate pMHC I with low monomeric affinities ( $K_D$ s  $>100 \mu\text{M}$ ), in contrast to pathogen-specific TCRs ( $K_D$ s  $\sim 1–50 \mu\text{M}$ )<sup>26</sup>. It is also established that CD8<sup>+</sup> T-cells expressing low affinity TCRs




**Figure 7. DK25 inhibits pancreatic  $\beta$ -cell killing by autoreactive CD8<sup>+</sup> T-cells.** (A)  $2 \times 10^3$   $^{51}\text{Cr}$ -labeled K562-PPI-A\*0201 cells were peptide-pulsed with ALWGPDPAAA at a concentration of  $10^{-5}$  M for 1 hour at 37°C. Clonal 1E6 CD8<sup>+</sup> T-cells were then added at an E:T ratio of 1:1 (upper panel) or 5:1 (lower panel) with or without the anti-human CD8 antibody DK25 at the indicated concentrations. Cytotoxicity was determined after 4 hours as described in the Materials and Methods. The mean  $\pm$  SD of three replicate assays is shown and is representative of three independent experiments. (B)  $5 \times 10^3$  HLA-A\*0201<sup>+</sup> islet cells were incubated for 4 hours with clonal 1E6 or NLV2 CD8<sup>+</sup> T-cells pre-treated for 30 minutes at 37°C with the anti-human CD8 antibody DK25 at the indicated concentrations. As a control, islet cells were peptide-pulsed with NLVPMVATV at a concentration of 5  $\mu\text{g}/\text{ml}$  for 30 minutes at 37°C. All data points represent the mean of triplicate evaluations at a final E:T ratio of 25:1. Islet cell killing was measured using a non-radioactive europium hydrophilic ligand assay as described in the Materials and Methods. Significance of the effect of DK25 was evaluated using linear regression and the t-test statistic as described in the materials and methods.

are strictly dependent on CD8 for ligand-induced activation<sup>33,34</sup>. These observations led us to hypothesise that CD8-targeted strategies could be used to block CD8<sup>+</sup> T-cell-mediated autoreactivity in a selective manner.


In preliminary experiments, we tested this possibility using the well-characterised autoreactive CD8<sup>+</sup> T-cell clone 1E6, which was isolated from a patient with T1D and displays potent reactivity against pancreatic  $\beta$ -cells<sup>39</sup>. The cognate epitope ALW is naturally processed from PPI and represents a major target for circulating effector CD8<sup>+</sup> T-cells in HLA-A\*0201<sup>+</sup> patients with T1D<sup>39</sup>. We found that peptide-induced activation of 1E6 is highly dependent on CD8, consistent with the previously reported low affinity TCR/pMHCI interaction ( $K_D = 278 \mu\text{M}$ )<sup>31,32</sup>. It is also established that 1E6 is extremely cross-reactive in the context of HLA-A\*0201, recognising  $\sim 1$  million different peptide ligands at physiologically relevant concentrations<sup>40</sup>. In line with the strict requirement for CD8, we observed a substantial winnowing of ligand cross-recognition in the absence of an intact pMHCI/CD8 interaction. Based on these findings and the fact that pathogen-specific CD8<sup>+</sup> T-cells displayed higher levels of functional sensitivity and robust activation profiles irrespective of CD8 engagement, we explored the potential utility of CD8-specific antibodies as a means to block CD8<sup>+</sup> T-cell activation selectively in response to self-derived peptides.

Previous work has shown that anti-CD8 antibodies exhibit diverse functional and phenotypic characteristics<sup>35,37</sup>. We selected the antibody clones SK1 and DK25 for further evaluation in view of the fact that they inhibit pMHCI tetramer binding and CD8<sup>+</sup> T-cell activation in the absence of non-specific stimulatory effects. The blocking effect on pMHCI tetramer binding was concentration-dependent in both cases and only apparent at low TCR/pMHCI affinities. The DK25 clone displayed greater efficacy in this regard and was therefore used in subsequent experiments.

To test the ability of DK25 to inhibit CD8<sup>+</sup> T-cell activation as a function of TCR/pMHCI affinity, we utilised a monoclonal system in which an HLA-A\*0201-restricted altered peptide ligand was used to generate a high affinity



**Figure 8. DK25 inhibits the activation of myelin-specific CD8<sup>+</sup> T-cells.**  $3 \times 10^4$  BW58 cells expressing human CD8 $\alpha\beta$  and the 2D1 TCR were stimulated overnight at 37°C in the presence of  $6 \times 10^4$  KLIETYFSK peptide-pulsed HLA-A\*0301<sup>+</sup> B-LCL target cells with or without the anti-human CD8 antibody DK25 at a final concentration of 0.25, 0.5 or 1 μg/ml (A,B). Supernatants were assayed for IL-2 by ELISA. The mean  $\pm$  SD of two replicate assays is shown in (B) and is representative of three independent experiments.



**Figure 9. CT-CD8a blocks the activation of autoreactive murine CD8<sup>+</sup> T-cells.**  $6 \times 10^4$  P815 murine mastocytoma cells were peptide-pulsed with LYLVCGERG at the indicated concentrations for 1 hour at 37°C.  $3 \times 10^4$  splenic G9C8 TCR-transgenic CD8<sup>+</sup> T-cells pre-treated for 30 minutes at 37°C with either the anti-mouse CD8 antibody YTS105.18 (A) or the anti-mouse CD8 antibody CT-CD8a (B) at the indicated concentrations were then added and the plates were incubated overnight at 37°C. Supernatants were assayed for MIP-1β by ELISA. The mean  $\pm$  SD of two replicate assays is shown and is representative of two independent experiments.

interaction with the clonotypically-expressed 1E6 TCR. DK25 efficiently blocked activation of the 1E6 clone in response to the cognate ligand ALW, whereas only a marginal effect was observed with the altered peptide ligand

**RQW**, which resembles a pathogen-derived epitope in terms of TCR/pMHCI affinity ( $K_D = 10 \mu\text{M}$ )<sup>3,41</sup>. Similar results were obtained across a panel of different autoreactive (1E6, 3F2 and 4C6) and pathogen-specific (ALF8, SB27 and MCNLV) CD8<sup>+</sup> T-cell clones. These findings suggest that blocking anti-CD8 antibodies can be used as selective agents for therapeutic purposes in the setting of autoimmune disease. In support of this notion, we found that DK25 can prevent CD8<sup>+</sup> T-cell-mediated destruction of pancreatic  $\beta$ -cells and block the activation of myelin-specific CD8<sup>+</sup> T-cells. As such, further research is warranted into the utility of this approach in the treatment of CD8<sup>+</sup> T-cell mediated autoimmunity. Although, it is important to note that more work is required to identify pMHCI antigens important in these diseases and assess the affinity of the TCRs that recognise them to understand the scope for this approach.

Previous studies have examined the ability of anti-CD8 antibodies to induce tolerance in murine models of T1D<sup>20,21</sup>. However, the YTS105.18 antibody clone used for this purpose does not inhibit either pMHCI tetramer binding or peptide-induced CD8<sup>+</sup> T-cell activation<sup>38</sup>. The tolerising effect of YTS105.18 is therefore distinct from the novel blocking strategy proposed here. In contrast, the CT-CD8a antibody displays an inhibitory phenotype in mice akin to SK1 and DK25 in humans<sup>35</sup>. Autoreactive murine CD8<sup>+</sup> T-cells were highly susceptible to CT-CD8a-mediated blockade *in vitro*, but the presence of non-specific stimulatory effects after prolonged exposure precluded confirmatory experiments *in vivo*<sup>37</sup>.

The next step will be to identify a suitable *in vivo* model to test the utility of this approach, to establish parameters such as optimal antibody concentration and assess the effect of targeting CD8 on the ability to mount an adequate pathogen specific response. The current literature suggests that during an acute response to pathogen, a broad range of TCR affinities are mobilised<sup>52</sup>. However, strong TCR ligation is required to sustain a T-cell response and as such, the TCR repertoire “matures” so that predominantly higher affinity TCRs remain<sup>52</sup>. Therefore, inhibiting the activation of low affinity pathogen-specific T-cells would not be anticipated to have a major impact on the ability to control infection as there would exist a degree of redundancy in the mobilised TCR repertoire, such that higher affinity TCRs would still be able to expand and control infection. Future *in vivo* work would also allow us to assess the possibility that autoreactive CD8<sup>+</sup> T-cells with higher than expected TCR affinities ( $K_{Ds} < 100 \mu\text{M}$ ) occur in the repertoire and the impact that this may have on the effectiveness of this approach.

The DK25 antibody targets CD8 $\alpha$ , which is expressed not only as a heterodimer with CD8 $\beta$ , exclusively on CD8<sup>+</sup> T-cells, but also in homodimeric form on dendritic cells, natural killer cells and  $\gamma\delta$  T-cells. Although the precise role of CD8 $\alpha\alpha$  is unclear, it is thought to act as a modulator of immune function<sup>53,54</sup>. The impact of chain specificity as a determinant of *in vivo* effects will therefore require further investigation as it is anticipated that antibodies which exclusively target the CD8 $\alpha\beta$  heterodimer would be more desirable and associated with a more predictable outcome. The exact mechanism of anti-CD8 antibody action is still unclear<sup>35,37</sup>. In addition, the use of monoclonal antibodies as therapeutics is often not ideal because they can be costly to produce, can mediate side effects (via FcR binding) and require intravenous administration. Although F(ab')<sub>2</sub> fragments retain the same phenotype as the parental antibody, the overall magnitude of the effect can be reduced<sup>37</sup>. Therefore, a viable alternative would be to design small molecular inhibitors that block the interaction between pMHCI and CD8 $\beta$ .

Current treatments for autoimmune and inflammatory diseases often rely on the induction of profound and/or long-lasting immunosuppression<sup>55–57</sup>. However, adverse effects are common and often relate to a negative impact on protective immune responses. Our data suggest that autoreactive CD8<sup>+</sup> T-cells can be preferentially targeted using blocking anti-CD8 antibodies. Additional modalities could also be employed based on the underlying biology, such as rationally designed small molecular inhibitors and soluble versions of the inhibitory immunoglobulin-like transcript 2 (ILT2) receptor<sup>58</sup>. These approaches hold the potential to suppress pathogenic CD8<sup>+</sup> T-cell-mediated autoimmunity without the attendant side effects that complicate standard therapeutic interventions.

## Materials and Methods

**Cells.** The following CD8<sup>+</sup> T-cell clones were used in this study: (i) 1E6 and 3F2, specific for the HLA-A\*0201-restricted PPI epitope ALWGPDAAA (residues 15–24)<sup>39</sup>; (ii) ALF3 and ALF8, specific for the HLA-A\*0201-restricted influenza A virus M1 epitope GILGFVFTL (residues 58–66)<sup>37</sup>; (iii) MCNLV and NLV2, specific for the HLA-A\*0201-restricted HCMV pp65 epitope NLVPMVATV (residues 495–503); (iv) 4C6, specific for the HLA-A\*2402-restricted PPI epitope LWMRLPLL (residues 3–11)<sup>59</sup>; and (v) SB27, specific for the HLA-B\*3508-restricted EBV BZLF1 epitope LPEPLPQGQLTAY (residues 52–64)<sup>60</sup>. Equilibrium binding affinities for the corresponding TCR/pMHCI interactions are shown in Table 2. C1R target cells expressing HLA-A\*0201 (C1R-A\*0201), HLA-A\*0201 D227K/T228A (C1R-A\*0201 D227K/T228A) or HLA-B\*3508 (C1R-B\*3508) were generated as described previously<sup>61</sup>. K562 cells expressing HLA-A\*2402 (K562-A\*2402) and BW58 cells expressing human CD8 $\alpha\beta$  and the 2D1 TCR, which is specific for the HLA-A\*0301-restricted myelin PLP epitope KLITYFSK (residues 45–53), were also described previously<sup>39,45</sup>. All CD8<sup>+</sup> T-cells were maintained in RPMI 1640 containing 100 U/ml penicillin, 100 mg/ml streptomycin, 2 mM L-glutamine and 10% heat-inactivated fetal calf serum (all Life Technologies) supplemented with 2.5% Cellkines (Helvetica Healthcare), 200 IU/ml IL-2 and 25 ng/ml IL-15 (both PeproTech).

**G9 transgenic mice.** Transgenic mice expressing the G9C8 TCR on the NOD.TCRC $\alpha^{-/-}$  genetic background were generated as described previously<sup>62,63</sup>. Mice were housed in microisolators or scintainers in the specific pathogen-free facility at Cardiff University. The G9C8 TCR is specific for the H-2K<sup>d</sup>-restricted insulin B chain epitope LYLVCGERG (residues 15–23)<sup>62,64</sup>. All procedures were performed in accordance with protocols approved by the UK Home Office.

TCR	Specificity	HLA restriction	Minimal epitope	K <sub>Deq</sub> (μM)
1E6	PPI	A*0201	ALWGPDPAAA	278 <sup>31,32</sup>
1E6	PPI	A*0201	RQWGPDPAAV	10 <sup>41</sup>
3F2	PPI	A*0201	ALWGPDPAAA	>250 <sup>73</sup>
4C6	PPI	A*2402	LWMRLPLL	100 <sup>73</sup>
SB27	EBV	B*3508	LPEPLPQGQLTAY	9.9 <sup>43,44</sup>

**Table 2.** Measured TCR/pMHCI equilibrium binding affinities (K<sub>Deq</sub>) for the CD8<sup>+</sup> T-cell clones used in this study.

**Anti-CD8 antibodies.** The following antibodies were used in this study: (i) APC-conjugated anti-human CD8α (DK25; DAKO); (ii) purified unconjugated anti-human CD8α (SK1; BD Biosciences); (iii) purified unconjugated anti-mouse CD8α (CT-CD8a; Lifespan BioScience); and (iv) purified unconjugated anti-mouse CD8α (YTS105.18; Thermo Scientific).

**CD8<sup>+</sup> T-cell effector function assays.**  $6 \times 10^4$  C1R-A\*0201, C1R-A\*0201 D227K/T228A, C1R-B\*3508, K562-A\*2402 or HLA-A\*0301<sup>+</sup> B-lymphoblastoid cell line (B-LCL) targets were pulsed with various concentrations of cognate peptide as indicated for 1 hour at 37 °C.  $3 \times 10^4$  clonal CD8<sup>+</sup> T-cells pre-treated for 30 minutes at 4 °C with various concentrations of the anti-human CD8 antibody DK25 as indicated were incubated with peptide-pulsed targets for 4 hours at 37 °C. Untreated effectors and unpulsed targets were used as controls. Supernatants were harvested and assayed for IFN-γ and MIP-1β by ELISA (R&D Systems).

**Combinatorial peptide library screens.** A 10mer CPL in positional scanning format was purchased from Pepscan. CPL scans were performed as described previously<sup>40,65</sup>. Briefly,  $6 \times 10^4$  C1R-A\*0201 or C1R-A\*0201 D227K/T228A cells were pulsed with each of the 10mer CPL mixtures (100 μM) in duplicate for 2 hours at 37 °C.  $3 \times 10^4$  clonal CD8<sup>+</sup> T-cells were then added and the plates were incubated overnight at 37 °C. Supernatants were harvested and assayed for MIP-1β by ELISA (R&D Systems).

**pMHCI tetramer staining.** Soluble biotinylated pMHCI proteins were manufactured and tetramerised with fluorochrome-labeled streptavidin as described previously<sup>6</sup>.  $5 \times 10^4$  clonal CD8<sup>+</sup> T-cells pre-treated with or without the indicated anti-human CD8 antibodies at various concentrations in 20 μl of PBS for 20 minutes at 4 °C were stained with 25 μg/ml of cognate pMHCI tetramer for 15 minutes at 37 °C. The cells were then washed and stained with LIVE/DEAD Fixable Aqua or Violet (Life Technologies) for 10 minutes at room temperature. Data were acquired using a FACSCantoII flow cytometer (BD Biosciences) and analysed with FlowJo software (TreeStar).

**Chromium release assay.**  $2 \times 10^3$  C1R-A\*0201, K562-A\*2402 or C1R-B\*3508 targets were labeled with 30 μCi of <sup>51</sup>Cr per 10<sup>6</sup> cells for 1 hour at 37 °C in RPMI 1640 containing 100 U/ml penicillin, 100 mg/ml streptomycin, 2 mM L-glutamine and 2% heat-inactivated fetal calf serum (all Life Technologies). Labeled targets were then pulsed with peptide for 1 hour at 37 °C. CD8<sup>+</sup> T-cells pre-treated with or without the indicated anti-human CD8 antibodies at various concentrations for 30 minutes at 4 °C were added at an effector-to-target (E:T) ratio of 5:1 and the plates were incubated for 4 hours at 37 °C. Spontaneous release was determined using targets incubated alone. Total release was determined using targets incubated in the presence of 5% Triton X-100 (Sigma-Aldrich). For each sample, 10 μl of supernatant was mixed with 150 μl of OptiPhase Supermix Scintillation Cocktail (Perkin Elmer) and analysed using a MicroBeta TriLux Liquid Scintillation and Luminescence Counter (Perkin Elmer). Specific lysis was calculated using the following formula:

$$(\text{experimental release} - \text{spontaneous release} / \text{total release} - \text{spontaneous release}) \times 100.$$

**Human islet killing assays.** Human islet isolations were performed as described previously using pancreata retrieved with the consent of donors' relatives and permission from the Ethical Review Committee of King's College Hospital<sup>66</sup>. Islet cells were cultured in monolayers for 16–24 hours with medium containing 16 μmol glucose plus IL-1β (50 IU/ml; Strathmann Biotec), tumor necrosis factor (TNF)-α (2500 IU/ml; Miltenyi Biotec), IFN-γ (500 IU/ml; Miltenyi Biotec) and IFN-α (1000 IU/ml; Roche Laboratories) to increase MHCI expression<sup>39</sup>. Clonal CD8<sup>+</sup> T-cells pre-treated for 30 minutes at 37 °C with various concentrations of the anti-human CD8 antibody DK25 as indicated were then incubated with  $5 \times 10^3$  islet cell targets at an E:T ratio of 25:1 for 4 hours at 37 °C. Cytotoxicity was analysed using a non-radioactive europium hydrophilic ligand assay with DELFIA Technology (Perkin Elmer)<sup>39,67</sup>.

**Protein expression, purification and surface plasmon resonance.** The ALF3 and ALF8 TCRs were expressed, refolded and purified using an engineered disulfide linkage between the α and β constant domains<sup>68</sup>. Soluble HLA-A\*0201 incorporating the influenza A virus M1 peptide GILGFVFTL was prepared as described previously<sup>69</sup>. All surface plasmon resonance experiments were conducted at 25 °C using a BIAcore 3000 instrument with HBS buffer (10 mM HEPES pH 7.4, 150 mM NaCl and 0.005% surfactant P20) containing 1% BSA to prevent non-specific binding. The human TCR-specific monoclonal antibody 12H8<sup>70</sup>, which recognises a conformation-dependent epitope, was amine-coupled to research-grade CM5 chips. Ligand binding to

solid-phase TCRs was then measured over a pMHC concentration range of 0.78–200  $\mu$ M according to established protocols<sup>68</sup>. Data were analysed using BIAevaluation software version 3.1 with the 1:1 Langmuir binding model.

**Statistical analysis.** DK25-induced shifts in functional sensitivity were quantified by calculating differences in pEC<sub>50</sub> (delta pEC<sub>50</sub>), a parameter determined as  $-\log$  [50% maximal response peptide concentration] for each condition (Supplementary Table 1)<sup>71</sup>. Differences in susceptibility to inhibition by DK25, comparing grouped autoreactive versus pathogen-specific CD8<sup>+</sup> T-cells, were assessed using the Wilcoxon-Mann-Whitney rank sum test<sup>72</sup>. The effect of DK25 in pancreatic islet experiments was assessed using linear regression and the t-test statistic with built-in tools in *Mathematica*<sup>72</sup>.

## References

- Gao, G. F. & Jakobsen, B. K. Molecular interactions of coreceptor CD8 and MHC class I: the molecular basis for functional coordination with the T-cell receptor. *Immunol Today* **21**, 630–636 (2000).
- Gao, G. F., Rao, Z. & Bell, J. I. Molecular coordination of alphabeta T-cell receptors and coreceptors CD8 and CD4 in their recognition of peptide-MHC ligands. *Trends Immunol* **23**, 408–413 (2002).
- Rosjohn, J., Gras, S., Miles, J. J., Turner, S. J., Godfrey, D. I. & McCluskey, J. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol* **33**, 169–200 (2015).
- Zamoyska, R. CD4 and CD8: modulators of T-cell receptor recognition of antigen and of immune responses? *Curr Opin Immunol* **10**, 82–87 (1998).
- Holler, P. D. & Kranz, D. M. Quantitative analysis of the contribution of TCR/pepMHC affinity and CD8 to T cell activation. *Immunity* **18**, 255–264 (2003).
- Wooldridge, L. *et al.* Interaction between the CD8 coreceptor and major histocompatibility complex class I stabilizes T cell receptor-antigen complexes at the cell surface. *J Biol Chem* **280**, 27491–27501 (2005).
- van den Berg, H. A., Wooldridge, L., Laugel, B. & Sewell, A. K. Coreceptor CD8-driven modulation of T cell antigen receptor specificity. *J Theor Biol* **249**, 395–408 (2007).
- Barber, E. K., Dasgupta, J. D., Schlossman, S. F., Trevillyan, J. M. & Rudd, C. E. The CD4 and CD8 antigens are coupled to a protein-tyrosine kinase (p56lck) that phosphorylates the CD3 complex. *Proc Natl Acad Sci USA* **86**, 3277–3281 (1989).
- Zamoyska, R. *et al.* Inability of CD8 alpha' polypeptides to associate with p56lck correlates with impaired function *in vitro* and lack of expression *in vivo*. *Nature* **342**, 278–281 (1989).
- Purbhoo, M. A. *et al.* The human CD8 coreceptor effects cytotoxic T cell activation and antigen sensitivity primarily by mediating complete phosphorylation of the T cell receptor zeta chain. *J Biol Chem* **276**, 32786–32792 (2001).
- Hutchinson, S. L. *et al.* The CD8 T cell coreceptor exhibits disproportionate biological activity at extremely low binding affinities. *J Biol Chem* **278**, 24285–24293 (2003).
- Arcaro, A. *et al.* Essential role of CD8 palmitoylation in CD8 coreceptor function. *J Immunol* **165**, 2068–2076 (2000).
- Arcaro, A. *et al.* CD8beta endows CD8 with efficient coreceptor function by coupling T cell receptor/CD3 to raft-associated CD8/p56(lck) complexes. *J Exp Med* **194**, 1485–1495 (2001).
- Peakman, M. Immunological pathways to beta-cell damage in type 1 diabetes. *Diabet Med* **30**, 147–154 (2013).
- Roep, B. O. & Tree, T. I. Immune modulation in humans: implications for type 1 diabetes mellitus. *Nat Rev Endocrinol* **10**, 229–242 (2014).
- Tsai, S., Sharni, A. & Santamaria, P. CD8+ T cells in type 1 diabetes. *Adv Immunol* **100**, 79–124 (2008).
- Friese, M. A. & Fugger, L. Autoreactive CD8+ T cells in multiple sclerosis: a new target for therapy? *Brain* **128**, 1747–1763 (2005).
- Friese, M. A. & Fugger, L. Pathogenic CD8(+) T cells in multiple sclerosis. *Ann Neurol* **66**, 132–141 (2009).
- Sinha, S., Itani, F. R. & Karandikar, N. J. Immune regulation of multiple sclerosis by CD8+ T cells. *Immunol Res* **59**, 254–265 (2014).
- Phillips, J. M. *et al.* Type 1 diabetes development requires both CD4+ and CD8+ T cells and can be reversed by non-depleting antibodies targeting both T cell populations. *Rev Diabet Stud* **6**, 97–103 (2009).
- Yi, Z. *et al.* Long-term remission of diabetes in NOD mice is induced by nondepleting anti-CD4 and anti-CD8 antibodies. *Diabetes* **61**, 2871–2880 (2012).
- von Herrath, M. G., Coon, B., Lewicki, H., Mazarguil, H., Gairin, J. E. & Oldstone, M. B. *In vivo* treatment with a MHC class I-restricted blocking peptide can prevent virus-induced autoimmune diabetes. *J Immunol* **161**, 5087–5096 (1998).
- Chatenoud, L., Thervet, E., Primo, J. & Bach, J. F. Anti-CD3 antibody induces long-term remission of overt autoimmunity in nonobese diabetic mice. *Proc Natl Acad Sci USA* **91**, 123–127 (1994).
- Chatenoud, L. & Bluestone, J. A. CD3-specific antibodies: a portal to the treatment of autoimmunity. *Nat Rev Immunol* **7**, 622–632 (2007).
- Sherry, N. *et al.* Teplizumab for treatment of type 1 diabetes (Protege study): 1-year results from a randomised, placebo-controlled trial. *Lancet* **378**, 487–497 (2011).
- Bridgeman, J. S., Sewell, A. K., Miles, J. J., Price, D. A. & Cole, D. K. Structural and biophysical determinants of alphabeta T-cell antigen recognition. *Immunology* **135**, 9–18 (2012).
- Liu, G. Y., Fairchild, P. J., Smith, R. M., Prowle, J. R., Kioussis, D. & Wraith, D. C. Low avidity recognition of self-antigen by T cells permits escape from central tolerance. *Immunity* **3**, 407–415 (1995).
- Zehn, D. & Bevan, M. J. T cells with low avidity for a tissue-restricted antigen routinely evade central and peripheral tolerance and cause autoimmunity. *Immunity* **25**, 261–270 (2006).
- Ober, B. T. *et al.* Affinity of thymic self-peptides for the TCR determines the selection of CD8(+) T lymphocytes in the thymus. *Int Immunol* **12**, 1353–1363 (2000).
- Yin, Y., Li, Y. & Mariuzza, R. A. Structural basis for self-recognition by autoimmune T-cell receptors. *Immunol Rev* **250**, 32–48 (2012).
- Cole, D. K. *et al.* Modification of MHC anchor residues generates heteroclitic peptides that alter TCR binding and T cell recognition. *J Immunol* **185**, 2600–2610 (2010).
- Bulek, A. M. *et al.* Structural basis for the killing of human beta cells by CD8(+) T cells in type 1 diabetes. *Nat Immunol* **13**, 283–289 (2012).
- Laugel, B. *et al.* Different T cell receptor affinity thresholds and CD8 coreceptor dependence govern cytotoxic T lymphocyte activation and tetramer binding properties. *J Biol Chem* **282**, 23799–23810 (2007).
- Wooldridge, L. *et al.* CD8 controls T cell cross-reactivity. *J Immunol* **185**, 4625–4632 (2010).
- Wooldridge, L. *et al.* Anti-CD8 antibodies can inhibit or enhance peptide-MHC class I (pMHC I) multimer binding: this is paralleled by their effects on CTL activation and occurs in the absence of an interaction between pMHC I and CD8 on the cell surface. *J Immunol* **171**, 6650–6660 (2003).
- Wooldridge, L. *et al.* Anti-coreceptor antibodies profoundly affect staining with peptide-MHC class I and class II tetramers. *Eur J Immunol* **36**, 1847–1855 (2006).

37. Clement, M. *et al.* Anti-CD8 antibodies can trigger CD8+ T cell effector function in the absence of TCR engagement and improve peptide-MHCI tetramer staining. *J Immunol* **187**, 654–663 (2011).
38. Shore, D. A., Issafras, H., Landais, E., Teyton, L. & Wilson, I. A. The crystal structure of CD8 in complex with YTS156.7.7 Fab and interaction with other CD8 antibodies define the binding mode of CD8 alphabeta to MHC class I. *J Mol Biol.* **384**, 1190–1202 (2008).
39. Skowera, A. *et al.* CTLs are targeted to kill beta cells in patients with type 1 diabetes through recognition of a glucose-regulated preproinsulin epitope. *J Clin Invest* **118**, 3390–3402 (2008).
40. Wooldridge, L. *et al.* A single autoimmune T cell receptor recognizes more than a million different peptides. *J Biol Chem.* **287**, 1168–1177 (2012).
41. Knight, R. R. *et al.* Human beta-cell killing by autoreactive preproinsulin-specific CD8 T cells is predominantly granule-mediated with the potency dependent upon T-cell receptor avidity. *Diabetes* **62**, 205–213 (2012).
42. Cole, D. K. *et al.* Hotspot autoimmune T cell receptor binding underlies pathogen and insulin peptide cross-reactivity. *J Clin Invest* **126**, 2191–2204 (2016).
43. Tynan, F. E. *et al.* T cell receptor recognition of a ‘super-bulged’ major histocompatibility complex class I-bound peptide. *Nat Immunol.* **6**, 1114–1122 (2005).
44. Tynan, F. E. *et al.* High resolution structures of highly bulged viral epitopes bound to major histocompatibility complex class I. Implications for T-cell receptor engagement and T-cell immunodominance. *J Biol Chem.* **280**, 23900–23909 (2005).
45. Friese, M. A. *et al.* Opposing effects of HLA class I molecules in tuning autoreactive CD8+ T cells in multiple sclerosis. *Nat Med.* **14**, 1227–1235 (2008).
46. Motozono, C. *et al.* Distortion of the major histocompatibility complex class I binding groove to accommodate an insulin-derived 10-mer peptide. *J Biol Chem.* **290**, 18924–18933 (2015).
47. Cole, D. K., Laugel, B., Clement, M., Price, D. A., Wooldridge, L. & Sewell, A. K. The molecular determinants of CD8 co-receptor function. *Immunology* **137**, 139–148 (2012).
48. Roep, B. O. & Peakman, M. Diabetogenic T lymphocytes in human type 1 diabetes. *Curr Opin Immunol.* **23**, 746–753 (2011).
49. Coppieters, K. T. *et al.* Demonstration of islet-autoreactive CD8 T cells in insulinitic lesions from recent onset and long-term type 1 diabetes patients. *J Exp Med.* **209**, 51–60 (2012).
50. Huseby, E. S., Huseby, P. G., Shah, S., Smith, R. & Stadinski, B. D. Pathogenic CD8 T cells in multiple sclerosis and its experimental models. *Front Immunol.* **3**, 64 (2012).
51. Prinz, J. C. The role of T cells in psoriasis. *J Eur Acad Dermatol Venereol* **17**, 257–270 (2003).
52. Zehn, D., Lee, S. Y. & Bevan, M. J. Complete but curtailed T-cell response to very low-affinity antigen. *Nature* **458**, 211–214 (2009).
53. Leishman, A. J. *et al.* T cell responses modulated through interaction between CD8alphalpha and the nonclassical MHC class I molecule, TL. *Science* **294**, 1936–1939 (2001).
54. Gangadharan, D. & Cheroutre, H. The CD8 isoform CD8alphalpha is not a functional homologue of the TCR co-receptor CD8alphabeta. *Curr Opin Immunol.* **16**, 264–270 (2004).
55. Dinarello, C. A. Anti-inflammatory agents: present and future. *Cell* **140**, 935–950 (2010).
56. Rammohan, K. W. & Shoemaker, J. Emerging multiple sclerosis oral therapies. *Neurology* **74** Suppl 1, S47–S53 (2010).
57. Steward-Tharp, S. M., Song, Y. J., Siegel, R. M. & O’Shea, J. J. New insights into T cell biology and T cell-directed therapy for autoimmunity, inflammation, and immunosuppression. *Ann N Y Acad Sci.* **1183**, 123–148 (2010).
58. Moysey, R. K. *et al.* High affinity soluble ILT2 receptor: a potent inhibitor of CD8(+) T cell activation. *Protein Cell* **1**, 1118–1127 (2011).
59. Kronenberg, D. *et al.* Circulating preproinsulin signal peptide-specific CD8 T cells restricted by the susceptibility molecule HLA-A24 are expanded at onset of type 1 diabetes and kill beta-cells. *Diabetes* **61**, 1752–1759 (2012).
60. Green, K. J., Miles, J. J., Tellam, J., van Zuylen, W. J., Connolly, G. & Burrows, S. R. Potent T cell response to a class I-binding 13-mer viral epitope and the influence of HLA micropolymorphism in controlling epitope length. *Eur J Immunol.* **34**, 2510–2519 (2004).
61. Wooldridge, L. *et al.* Enhanced immunogenicity of CTL antigens through mutation of the CD8 binding MHC class I invariant region. *Eur J Immunol.* **37**, 1323–1333 (2007).
62. Wong, F. S., Visintin, I., Wen, L., Flavell, R. A. & Janeway, C. A. Jr. CD8 T cell clones from young nonobese diabetic (NOD) islets can transfer rapid onset of diabetes in NOD mice in the absence of CD4 cells. *J Exp Med.* **183**, 67–76 (1996).
63. Wong, F. S. *et al.* Activation of insulin-reactive CD8 T-cells for development of autoimmune diabetes. *Diabetes* **58**, 1156–1164 (2009).
64. Wong, F. S. *et al.* Identification of an MHC class I-restricted autoantigen in type 1 diabetes by screening an organ-specific cDNA library. *Nat Med.* **5**, 1026–1031 (1999).
65. Ekeruche-Makinde, J. *et al.* Peptide length determines the outcome of TCR/peptide-MHCI engagement. *Blood* **121**, 1112–1123 (2013).
66. Huang, G. C. *et al.* The development of new density gradient media for purifying human islets and islet-quality assessments. *Transplantation* **77**, 143–145 (2004).
67. Blomberg, K., Hautala, R., Lovgren, J., Mukkala, V. M., Lindqvist, C. & Akerman, K. Time-resolved fluorometric assay for natural killer activity using target cells labelled with a fluorescence enhancing ligand. *J Immunol Methods* **193**, 199–206 (1996).
68. Gras, S. *et al.* The shaping of T cell receptor recognition by self-tolerance. *Immunity* **30**, 193–203 (2009).
69. Gras, S. *et al.* Structural bases for the affinity-driven selection of a public TCR against a dominant human cytomegalovirus epitope. *J Immunol.* **183**, 430–437 (2009).
70. Borg, N. A. *et al.* The CDR3 regions of an immunodominant T cell receptor dictate the ‘energetic landscape’ of peptide-MHC recognition. *Nat Immunol.* **6**, 171–180 (2005).
71. Sprent, P. & Smeeton, N. C. Applied nonparametric statistical methods. *Chapman & Hall/CRC, London* (2007).
72. Casella, G. & Berger, R. L. Statistical inference. *Duxbury Advanced Series, Cengage Learning* (2008).
73. Dolton, G. *et al.* Comparison of peptide-major histocompatibility complex tetramers and dextramers for the identification of antigen-specific T cells. *Clin Exp Immunol.* **177**, 47–63 (2014).

## Acknowledgements

We thank Dr. Ann Ager for helpful discussions and critical review of the manuscript, Prof Andrew Sewell for generous funding support, and Dr. Katherine Attfield and Prof Lars Fugger for providing BW58 cells expressing human CD8 $\alpha\beta$  and the 2D1 TCR. The 4C6 clone was kindly provided by D Kronenberg-Versteeg. This work was supported by a Wellcome Trust Institutional Strategic Support Fund Award and a National Multiple Sclerosis Society Pilot Project Award (5997425). J.A.P. was funded by a Diabetes UK PhD Studentship. A.S. and M.P. were supported by the National Institute for Health Research Biomedical Research Centre at Guy’s and St Thomas’ National Health Service Foundation Trust and King’s College London. S.G. is an Australian Research Council Future Fellow (FT120100416), J.R. is a National Health and Medical Research Council Australia Fellow (FT50) and D.A.P. is Wellcome Trust Senior Investigator (100326/Z/12/Z). L.W. was supported by a Wellcome Trust Intermediate Clinical Fellowship (WT079848MA).

### Author Contributions

M.C., J.A.P., S.G., H.A.v.d.B., A.L., S.L.-L., M.D.W., T.D., J.E.M., J.E.-M., E.G. and A.S. performed experiments and data analysis. N.P.R., J.R., S.R.B., D.A.P., F.S.W. and M.P. provided reagents and critical supervision. M.C., A.S. and L.W. designed experiments. M.C., D.A.P., A.S. and L.W. wrote the manuscript.

### Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Clement, M. *et al.* Targeted suppression of autoreactive CD8<sup>+</sup> T-cell activation using blocking anti-CD8 antibodies. *Sci. Rep.* **6**, 35332; doi: 10.1038/srep35332 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016