

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/98041/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Svenssoon, O., Sanderson Bellamy, Angelina, Van den Brink, P, Tedengren, M. and Gunnarsson, J. S. 2018. Assessing the ecological impact of banana farms on water quality using aquatic macroinvertebrate community composition. *Environmental Science and Pollution Research* 25 (14) , pp. 13373-13381. 10.1007/s11356-016-8248-y file

Publishers page: <http://dx.doi.org/10.1007/s11356-016-8248-y> <<http://dx.doi.org/10.1007/s11356-016-8248-y>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Assessing the ecological impact of banana farms on water quality using aquatic macroinvertebrate**  
2 **community composition**

3 **In: Environmental Science and Pollution Research**

4  
5 Ola Svensson<sup>1\*</sup>, Angelina Sanderson Bellamy<sup>1,2</sup>, Paul J. Van den Brink<sup>3,4</sup>,

6 Michael Tedengren<sup>1</sup>, Jonas S. Gunnarsson<sup>1</sup>

7  
8 <sup>1</sup> Department of Ecology, Environment and Plant sciences (DEEP), Stockholm University, SE-10961,  
9 Stockholm, Sweden.

10 <sup>2</sup> Sustainable Places Research Institute, Cardiff University, 33 Park Place, CF10 3BA United Kingdom

11 <sup>3</sup>Dept of Aquatic Ecology and Water Quality Management, Wageningen University, P.O. Box 47, 6700 AA  
12 Wageningen, The Netherlands

13 <sup>4</sup> Alterra, Wageningen University and Research, P.O. Box 47, 6700 AA Wageningen, The Netherlands

14 \* Corresponding author, Phone number: +46-8-16 40 13

15 Address:

16 Ola Svensson

17 Department of Ecology, Environment and Plant sciences (DEEP)

18 Stockholm University

19 SE-10691 Stockholm

20 Sweden

21 Email: [ola.svensson@su.se](mailto:ola.svensson@su.se)

22 Phone: +46-8-16 40 13

23  
24  
25  
26  
27  
28 **Abstract**

29 In Costa Rica considerable effort goes to conservation and protection of biodiversity, while at the same time  
30 agricultural pesticide use is among the highest in the world. Several protected areas, some being wetlands or  
31 marine reserves, are situated downstream large-scale banana farms, with an average of 57 pesticide applications  
32 per year. The banana industry is increasingly aware of the need to reduce their negative environmental impact,  
33 but few ecological field studies have been made to evaluate the efficiency of proposed mitigation strategies. This  
34 study compared the composition of benthic macroinvertebrate communities up- and downstream effluent water  
35 from banana farms in order to assess whether benthic invertebrate community structure can be used to detect  
36 environmental impact of banana farming, and thereby usable to assess improvements in management practices.  
37 Aquatic invertebrate samples were collected at 13 sites, using kick-net sampling, both up- and downstream  
38 banana farms in fast flowing streams in the Caribbean zone of Costa Rica. In total, 2888 invertebrate specimens  
39 were collected, belonging to 15 orders and 48 families or taxa. The change in community composition was  
40 analyzed using multivariate statistics. Additionally, a biodiversity index and the Biological Monitoring Working  
41 Party (BMWP) score system was applied along with a number of community composition descriptors.  
42 Multivariate analyses indicated that surface waters immediately up- and downstream large-scale banana farms  
43 have different macroinvertebrate community compositions with the most evident differences being higher  
44 dominance by single taxa and a much higher total abundance, mostly of that same taxon. Assessment of  
45 macroinvertebrate community composition thus appears to be a viable approach to detect negative impact from  
46 chemical-intensive agriculture and could become an effective means to monitor the efficacy of changes/proposed  
47 improvements in farming practices in Costa Rica and similar systems.

48  
49  
50 **Key words:** Costa Rica, banana production, benthic macroinvertebrates, water quality, monitoring, risk  
51 assessment.

1  
2  
3  
4  
5  
6  
7  
8  
9

### **Acknowledgements**

We would like to thank: Manuel Zumbado at INBio and Bert Kohlmann at the University of Agronomy EARTH, for help with taxonomic identification material; Miguel Rivera, Jose Moore and other farm owners for their assistance and for giving us access to the field sites and John Bellamy for illustrations. Funding was provided by a Swedish International Development Cooperation Agency planning grant and by the Swedish Research Council Formas, grant nr 2005-473-3035-21.

## 1 **1. Introduction**

2 Costa Rica is one of the richest countries in the world in terms of biodiversity and considerable effort goes to  
3 conservation and protection. Several protected areas, some being wetlands or marine reserves, are however,  
4 situated downstream agricultural areas, where the use of agrochemicals is very high (Schreinemachers &  
5 Tipraqsa 2012) and run-off into nearby surface waters is of particular concern (Castillo et al. 2006). A major  
6 contributor of agrochemicals to the surrounding environment is the large-scale banana production, which  
7 receives an average of 57.5 pesticide applications per year as well as 2775 kg/ha of synthetic fertilizers (Bellamy  
8 2013; Bravo et al. 2013). Several of the pesticides used in banana production have been detected in the aquatic  
9 environment downstream of banana production areas (Castillo et al. 2006), some in concentrations expected to  
10 have acute or chronic toxic effects on aquatic organisms according to toxicity values derived from laboratory  
11 toxicity tests (Diepens et al. 2014; Arias-Andres et al. 2016; Rämö et al. 2016).

12 Banana companies are today increasingly aware of the need to reduce their negative environmental impact, and  
13 several changes in management practices have resulted in some companies being certified according to one of  
14 several certification systems (e.g. Rainforest Alliance and ISO14000). Attempts to reduce environmental impact  
15 by farms include: sediment traps that are constructed to reduce erosion and capture/retain pesticides adhered to  
16 solids; riparian vegetation zones that are planted/left to intercept spray drift, prevent erosion and reduce surface  
17 flow and leaching of pesticides; manual chopping of weeds instead of using herbicides; manual injections of  
18 nematicides into the banana plant instead of applying soil granular nematicides; and post-harvest applications of  
19 fungicides using brushes instead of fumigation chambers, thereby reducing the amount of pesticides used.

20 Some of these practices may reduce the negative impact on the environment, but few ecological field studies  
21 have been done to evaluate the efficiency of mitigation strategies that aim to reduce negative environmental  
22 impact in Costa Rican rivers and in similar tropical aquatic systems. Monitoring changes of benthic  
23 macroinvertebrate community composition is commonly used in monitoring programs and ecological status  
24 assessments of freshwater and marine coastal systems around the world (e.g. van Hoey et al. 2010; von der Ohe  
25 & Goedkoop 2013). In this study we evaluate changes in benthic community composition up- and downstream  
26 from banana plantations as a means to evaluate ecological effects of current agricultural practices and the  
27 efficiency of proposed improvements, as well as a complement to chemical analysis of pesticide residues in  
28 environmental risk assessment.

29 Benthic macroinvertebrates are usually abundant in rivers, represent several trophic levels, participate in nutrient  
30 cycling and differ in sensitivity to pollution. Most of them have small home ranges, at least in aquatic stages, and  
31 usually have long life cycles and thus are good bioindicators as they provide information about the water quality  
32 integrated over a longer time period, compared to the values given by water samples taken at discrete points in  
33 time. Pesticide and nutrient levels in the aquatic environment can be expected to vary, with peaks after  
34 application and high rainfall events. Monitoring of pesticide levels thus requires a very frequent sampling to  
35 detect peak concentrations (Liess et al. 2003). Another concern is that toxic effects can result from exposure near  
36 or below the analytical detection limit for a given pesticide (Walter et al. 2002) or from a combination of  
37 pesticides and other stressors, e.g. temperature or high nutrient loads (Polidoro & Morra 2016). It is also  
38 important to consider the effect of chronic exposure to pesticides as well as the exposure to mixtures of several  
39 pesticides, which together can cause toxic effects through additive toxicity (Verbruggen & Van den Brink 2010).

40  
41 The possible additive or synergistic effects between stressors are a major concern in rivers, which receive  
42 irrigation and run-off water from banana farms. Large-scale banana farming relies on the use of fungicides,  
43 nematicides, insecticides and herbicides. Most often several different compounds of each type of pesticide are  
44 applied over the year in order to minimize risk of inducing resistance in pests. The extensive system of drainage  
45 canals in a typical banana farm causes increased stream flashiness and sedimentation and due to high  
46 precipitation a substantial amount of pesticides and nutrients end up in the aquatic environment. Non-target  
47 aquatic organisms further downstream will thus be subjected to a complex mixture of toxic substances, fertilizers  
48 and changes in stream flow. To assess cumulative effects of several physical and chemical stressors, responses  
49 thus have to be studied at the community or ecosystem level and using benthic macroinvertebrates has proved to  
50 be a cost-effective monitoring tool.

51 In the present study we evaluated whether the overall impact of banana farming affects aquatic benthic  
52 macroinvertebrate fauna in waters subjected to agricultural run-off to test if benthic macroinvertebrate  
53 community composition can be used as a bioindicator of ecological stress to these aquatic ecosystems. Our  
54 research hypothesis was that surface waters downstream of banana farms will have a different benthic  
55 macrofauna community composition with a lower diversity compared to upstream sites. The changes in

1 community composition were assessed at the family level, with the objective of testing a robust, ecologically  
2 relevant method to detect environmental impact of agricultural run-off.

3

## 4 **2. Material and Methods**

### 5 *2.1 Sites*

6 Aquatic benthic invertebrate samples were collected at 13 sites in the Caribbean low-lands of Costa Rica  
7 between March 8 and April 26, i.e. during the dry season, 2007 (Table 1). Sites were chosen both up- and  
8 downstream in rivers and watercourses receiving run-off from banana farms and at sites assumed not to be  
9 affected by banana farming (Fig 1). A high natural variability in community composition can be expected  
10 between and along streams. Both stream order and stream size influence taxa richness and community structure  
11 (Malmqvist and Hoffsten 2000; Vannote et al. 1980), as do local factors, such as riparian characteristics, water  
12 chemistry and in-stream habitat structure. Sampling sites were thus, when possible, chosen in pairs along the  
13 same watercourse, with one site situated upstream and the second downstream banana farms (Fig 1). By  
14 comparing sites in an upstream-downstream fashion the difficulty with interpretation associated with the natural  
15 inter-stream variation is greatly reduced. Spatial habitat heterogeneity, current velocity at base and high flows,  
16 and type of substrate also affect within-site diversity of stream invertebrates (Beisel et al. 2000). Sampling of  
17 highly similar habitats was therefore favoured to reduce this variability, with fast flowing streams, mostly  
18 cobbles for substrate in runs and riffles, and no or little macrophytes being the preference (see Table 1 for  
19 comparison between sites). Keeping to those prerequisites in combination with limited access, only 3 rivers were  
20 sampled in a true, replicated upstream- downstream fashion (see Table 2).

21 In the provinces of Sarapiquí and Siquirres, where conventional, large-scale banana farms are abundant, Río  
22 Sucio and Río Pacuare were sampled up- and downstream of single farms as well as downstream ‘banana  
23 districts’, with large-scale banana farms being the dominating land use (Fig 1, Table 1). In addition, samples  
24 were taken in two small streams, with one site (SSS1Nat) located within National Park Braulio Carrillo near the  
25 source of Río Sucio, and another site upstream from banana farms (SSS1up). In the province of Talamanca a  
26 small stream between Cahuita and Hone Creek was sampled up- and downstream a small-scale, certified organic  
27 farm. Finally, in the province of Guácimo a small, first order stream located adjacent to a low-input banana farm  
28 within EARTH University a few hundred meters west of Río Dos Novillos was sampled as was Río Parismina  
29 downstream a conventional banana farm. At each site the following data were recorded in a field protocol: GPS-  
30 position, date and time, rainfall within last 24 hours, present cloud conditions, measurements of temperature (air  
31 and water), assessment of river width, depth, current, velocity, turbidity, colour and amount of shade, assessment  
32 of substrate composition (amount boulders, stones, sand, silt, plant material etc.). Additionally, we measured  
33 approximate distances from the sample site to the source of the stream or river sampled.

### 34 *2.2 Sampling and identification of macroinvertebrates*

35 We used kick net sampling since Armitage (1978) and Pollard (1981) found the method to give consistent  
36 results. By disturbing the bottom, specimens are dislodged and drift into a net held immediately downstream.  
37 The net used was a D-framed 40 cm wide kick net with 0.5 mm mesh size. To obtain a representative composite  
38 sample an area equivalent to the area of the net was disturbed at six sub-sampling positions (randomly chosen  
39 within an area of about 25 m<sup>2</sup>). Each sampling position was approached either at a right angle to the flow  
40 direction or from downstream in order not to sample where the bottom had been inadvertently disturbed. The net  
41 was held as close as possible to the streambed. The substrate in front of the net was disturbed, either by kicking  
42 or by hand. The latter was favoured due to the substrate; in most cases rocks of a size that would not easily turn  
43 over by kicking. Animals and epiphytes were dislodged by brushing hands over rock surfaces and collected with  
44 the softer substrate into the net. All sweeping of substrate/disruption of bottom was directed toward the net to  
45 reduce loss of swimming specimens. One composite sample represents approximately 1 m<sup>2</sup> and up to three  
46 composite samples were collected at each site. Sites are designated by an abbreviation for actual watercourse, a  
47 numeral for relative position along the watercourse and whether it is an up- or downstream site.

48 Samples were transferred to labelled containers and preserved in 70% alcohol. Sorting and identification was  
49 done to family level (Oligochaeta, Acarina were only identified to order and Bivalvia to class) under stereoscope  
50 using relevant taxonomical keys (Thorp & Covich 1991; Roldán Pérez 1992). Reference specimens were  
51 deposited at Costa Rica's National Institute of Biodiversity (INBio).

### 52 *2.3 Analyses of benthic community composition*

1 Our research question was whether surface waters downstream banana farms have a different benthic  
 2 macroinvertebrate community composition with a lower diversity compared to surface waters upstream banana  
 3 farms. The effect of agricultural run-off on benthic macroinvertebrate community composition was studied with  
 4 multivariate statistics. Similarity between invertebrate communities up- and downstream plantations was  
 5 assessed using principal component analysis (PCA) (Van den Brink et al. 2003; van Wijngaarden et al. 1995)  
 6 using CANOCO (version 5) (Ter Braak and Smilauer, 2012). PCA was used since the invertebrate data set had a  
 7 short length of gradient (2.8 SD; Van Wijngaarden et al., 1995), while the abundance data were  $\ln(2x+1)$   
 8 transformed (see Van den Brink et al., 2000 for rationale). Analyses were performed on mean values per site  
 9 where more than one composite sample exists. PCA generates an ordination diagram, which allows comparison  
 10 of how closely the different sites are related to each other in terms of community composition and additionally  
 11 show how the taxa composition varies between sites, i.e. upstream or downstream banana farms. Sites that lie  
 12 close together on the PCA diagram share a more similar community composition than those sites that lie further  
 13 apart (Ter Braak 1995). Site characteristics were introduced as supplementary explanatory variables to assess the  
 14 correlations between taxa abundance values and the levels of the explanatory variables (Van den Brink et al.,  
 15 2003).

16 The Biological Monitoring Working Party (BMWP) score system (National Water Council 1981), originally  
 17 developed in Great Britain as a rapid and sensitive method to determine water quality using macroinvertebrate  
 18 sensitivity to organic pollution has also been adapted for use in tropical environments, and has proved to  
 19 correctly assess water quality in e.g. Thailand (Mustow 2002). The BMWP scores adapted to Costa Rican  
 20 conditions according to Springer et al. (2007) were used to rank the sites based on their indicated sensitivity to  
 21 organic pollution. In the BMWP system a score (from 1 for the most tolerant to 10 for the most sensitive) is  
 22 assigned to different taxa depending on their sensitivity to organic pollution and only requires identification to  
 23 the family level (Oligochaeta only to class). In order to make interpretation less sensitive to sampling effort,  
 24 Armitage et al. (1983) and others (e.g. Friedrich et al. 1996) suggest dividing the total sample score with the  
 25 number of contributing taxa, giving the result as Average Score Per Taxon (ASPT). These values as well as  
 26 number of families, individuals per taxon, Shannon-Wiener diversity index, EPT index (Lenat 1988) and percent  
 27 contribution of the most abundant taxon out of the total abundance were compared to determine if differences  
 28 between sites could be detected.

29

### 30 **3. Results**

31 In total, 2888 specimens were collected, belonging to 15 orders and 48 families or taxa. The PCA diagram  
 32 clearly shows the differences in community composition between the rivers and between up- and downstream  
 33 sites (Fig. 2). The horizontal and vertical axes (displaying 30% and 18% of the total variation, respectively)  
 34 show that most upstream sites and taxa are located in the upper, right part of the diagram, while most  
 35 downstream sites are located in the lower, left quadrant, where also no taxa are located. This shows that most  
 36 downstream sites have a poorer community composition compared to the upstream sites. Surprisingly, the  
 37 sample taken in the national park (Sarapiquí) clusters together with the downstream sites. The samples taken in  
 38 the Río Sucio and Río Pacuare just below the conventional farm are still relatively rich in taxa (i.e. located on the  
 39 right side of the diagram), while their more downstream located sites are located in the lower, left quadrant.

40 Fig. 2B shows how the community composition at each site relates to the explanatory variables. The sites located  
 41 in the left, lower part of the diagram, have a low number of taxa and are correlated with being downstream both  
 42 organic and conventional farms, high turbidity, a high % of still water, a high presence of sandy, silty  
 43 substratum, a high distance from source and a large width of the river. Upstream sites are correlated with a  
 44 higher amount of shade, an intermediate river width and a higher number of taxa.

45 The number of families per site ranged from 2.3 to 19 with a mean value of 11.6 taxa per site. Family/taxa  
 46 richness was higher in upstream samples as were BMWP-scores and Shannon-Wiener diversity values (Table 2).  
 47 In the Talamanca stream passing an organic farm there were fewer orders present in the upstream site (site  
 48 SST1up, 7 orders) compared to the downstream site (site SST2down, 11 orders). Only 6 orders were found at  
 49 site SSS1Nat within the national park, but 14 families, meaning higher within-order diversity (Table 2). With  
 50 regards to abundance, Chironomidae (i.e. midge larvae) was the most abundant family at several sites (RS1up,  
 51 RS2down, SST2down, SSS1Nat, RS4down and RPc3down) and were found at all sites. The range of  
 52 Chironomid abundance varied between 10% downstream a banana district along Río Sucio (site RS3down) to  
 53 91% downstream another banana district along Río Pacuare (site RPc3down). Within the order of Trichoptera  
 54 (caddisflies), Glossomatidae, Hydropsychidae and Philopotamidae all had higher abundances at downstream

1 sites despite high BMWP scores (score 8, 5, 7, respectively). Leptoceridae, on the other hand, were fewer  
 2 downstream (score 8; Table 2). EPT index values varied between 0,63 downstream a banana district and 9  
 3 upstream an organic farm, and were higher at upstream sites.

4 Invertebrate diversity at the order level varied between 3 and 11 orders (mean 7.1), and was higher in upstream  
 5 samples except for sites SST1up and SST2down (up- and downstream the organic farm). Diptera were found at  
 6 all sites (total 624 individuals), and were the dominant order at site RS4down and RPc3down, both downstream  
 7 banana districts. At site RS4down, one Diptera family contributed with 53% of total abundance while at site  
 8 RPc3down two families added up to 92%. This is in contrast to site SSS1Nat, in the national park, where Diptera  
 9 contributed 32% of the sample, but were represented by four families. Trichoptera contributed a total of 1185  
 10 individuals. Ephemeroptera (553 individuals), missing only at site RPc3down, were the dominant order at site  
 11 RS3down, both sites situated downstream large banana districts. Coleoptera were missing at sites RS3down,  
 12 RS4down, and RPc3down, i.e. the sites downstream large banana districts. Plecoptera were found only at site  
 13 SST1up and SSG1down, i.e. upstream the organic and downstream the low-input farm. Only a few taxa were  
 14 absent at most sites. None of those taxa present only at one single site were found downstream large-scale  
 15 conventional banana farms. Special note should be taken that oligochaetes (with the lowest sensitivity score, 1)  
 16 were only found at upstream sites.

17

#### 18 4. Discussion

19 The aim of the study was to assess if benthic macroinvertebrate community composition at the resolution of  
 20 family level could be used to evaluate improvements in banana farming practices. The PCA ordination plots  
 21 show differences in community composition between rivers but community composition also differed between  
 22 up- and downstream sites in the same river (Fig 2). The explanatory variable 'upstream' was positively  
 23 associated with a larger number of taxa (Fig 2), indicating a general trend that upstream sites are more species-  
 24 diverse. Higher river width was inversely correlated with number of taxa indicating that other factors than  
 25 oxygen stress are affecting these communities. The fact that Oligochaetes, normally very tolerant to low oxygen  
 26 levels, are only found at upstream sites further supports this conclusion (Fig 3). Oligochaetes have been found to  
 27 be relatively sensitive to fungicides (Cuppen et al. 2000), which have been found in surface waters downstream  
 28 banana farms in streams nearby (Castillo et al. 2000; Diepens et al. 2014; Arias-Andres et al. 2016; Echeverria-  
 29 Saenz et al. 2016; Rämö et al. 2016). Thus, pesticides used by banana farms may be influencing patterns shown  
 30 in the biplots presented here (Fig 2).

31 The BMWP scores, taxa richness and diversity indices were slightly lower downstream conventional farms than  
 32 upstream, interpreted as a response to water quality or habitat deterioration. It should be emphasized that  
 33 upstream 'reference' sites in some cases are affected by land use further upstream, i.e. not to be considered to  
 34 represent pristine conditions (Fig 1). This is e.g. the case for sites RS1up and RPc1up, the upstream conventional  
 35 farm sites in the pairwise comparison. Therefore, a reduction of taxa richness and loss of the most sensitive  
 36 species may have already occurred further upstream, possibly explaining the sometimes modest differences  
 37 found when sites up- and downstream banana farms were compared (Table 2). The small difference in taxa  
 38 richness is to some extent related to an increased richness of tolerant taxa at some of the downstream sites.  
 39 Environmental impact is therefore difficult to interpret from taxa richness alone. Diversity index figures are  
 40 likewise unaffected if one taxon replaces another in response to pollution, and, accordingly, differences in  
 41 diversity were moderate. A metric that showed a distinct difference was abundance, where, contrary to results by  
 42 Paaby et al. (1998), abundance was found to be higher at sites downstream conventional farms. In the streams  
 43 not affected by banana farms and the one passing a small-scale organic farm the number of individuals per taxon  
 44 was quite low, but the number of individuals per taxon almost doubles downstream large-scale banana farms,  
 45 despite the short distance. Higher abundance in these cases co-varies with increased dominance by one or two  
 46 taxa (Hydropsychidae and Chironomidae at site RS2down resp. Glossomatidae at site RPc2down), indicating  
 47 stress (Pearson & Rosenberg 1978) and evenness have been found to respond faster than species richness to  
 48 environmental stress (Chapin et al. 2000). Previous as well as recent studies in nearby rivers, which also receive  
 49 fertilizers and pesticides from banana plantations have shown similar effects on the invertebrate community  
 50 (Castillo et al. 2006; Pringle and Ramirez 1998; Echeverria- Saenz et al. 2016). The ASPT (Average Score Per  
 51 Taxon) score, contrary to expected, suggested that there were more sensitive taxa downstream banana farms  
 52 compared to sites upstream (Table 2). Oligochaetes, with a low score, were present only at the upstream sites  
 53 possibly due to fungicide exposure, lowering the ASPT scores compared to downstream sites and this highlights  
 54 the shortcomings of many water quality score systems. The BMWP score system is based on sensitivity of  
 55 families to oxygen depletion, which is used as an indicator of organically polluted surface waters. While this  
 56 generally correlates with agricultural runoff, particularly with regards to sediment and fertilizer run-off, the

1 BMWP score system does not reflect species' sensitivity to pesticide exposure. Rico and Van den Brink (2015)  
 2 propose a trait-based methodology using focal species that also incorporates landscape characteristics to improve  
 3 insecticide risk assessment based on invertebrate monitoring. However, at the community level, the response of  
 4 aquatic invertebrates to stress, e.g. lower diversity and higher abundance of some more tolerant species can be  
 5 expected to be similar regardless of the stressor (e.g. oxygen deficiency or contaminants such as metals, oil or  
 6 polycyclic aromatic hydrocarbons (Diaz 1992)).

7 Pringle and Ramirez (1998) found Diptera (e.g. Chironomidae) and Ephemeroptera to be dominant insect groups  
 8 at sites both in primary forest and in streams draining banana plantations. Sensitivity varies among Dipteran  
 9 families, but Chironomidae, according to the BMWP score system, are considered tolerant to organic pollution  
 10 and oxygen deficiency. In the present study Chironomidae were found at all sites, and were also to varying  
 11 degree the dominating taxon at six of the sites, four of which were downstream sites. At the site furthest  
 12 downstream Río Pacuare (site RPe3down), receiving run-off from several large-scale banana farms, the  
 13 abundance of Chironomidae was one order of magnitude greater than that of the second most abundant taxon,  
 14 and the extremely low total abundance at this site indicates very poor conditions. Ephemeroptera, generally  
 15 considered as relatively sensitive to organic pollution, were not represented at the aforementioned site, but  
 16 dominated at four other sites, two of which were situated downstream banana farms. However, the  
 17 Ephemeroptera families observed in these streams include several species (*Baetis spp.* and *Caenis spp.*) that are  
 18 commonly found in organically enriched streams (Barbour et al. 1999). Thus it is important to consider that  
 19 while scores are based on aggregate sensitivity of species within the order or family, there may be some species  
 20 more or less sensitive than the aggregate score given to a group. Other examples from the data are  
 21 Leptophlebiidae (Ephemeroptera) and Glossomatidae (Trichoptera), two families with high BMWP scores, which  
 22 were also present in high numbers or even dominant at downstream sites.

23 Sampling of rivers affected by small-scale organic farms proved to be difficult due to the lack of permanent  
 24 streams or due to streams being impacted by other land use upstream. The one sampled, though, presented an  
 25 upstream reference site within primary forest, and the banana farm being the only land use. The choice of  
 26 sampling sites was in general limited by access difficulties in combination with aforementioned requisites of  
 27 comparable habitats. Ideally more sites would have been sampled and preferably some with large-scale banana  
 28 farms being the only land use. However, with the upstream-downstream samples taken within a rather short  
 29 distance, and banana farming being the most significant land use and also the most chemical intense, one can  
 30 assume that most of the observed effects are due to production practices on banana farms.

31

## 32 5. Conclusions

33 Although it can be difficult to distinguish natural variations in e.g. diversity and community composition from  
 34 the effects of human impact, the consistent pattern with increasing dominance by a single or only a few taxa at  
 35 downstream sites indicates an impact from banana farming. The present study hints at a lower impact by organic  
 36 farming, but lacks replication to support it. The fact that differences, although small, were detected when  
 37 comparing up- and downstream single farms, implies that monitoring of macroinvertebrate community  
 38 composition is useful for assessing management practices and improvements proposed or introduced by the  
 39 banana industry aiming to produce bananas in a more sustainable way. As monitoring invertebrate community  
 40 composition is highly ecologically relevant, we recommend that it should be done in combination with chemical  
 41 analysis of pesticide residues in environmental monitoring programs in Costa Rica and in ecological risk  
 42 assessment of these rivers and similar aquatic systems.

43

## 44 References

- 45 Armitage PD (1978) Downstream changes in the composition, numbers and biomass of bottom fauna below  
 46 Cow Green Reservoir and in unregulated Maize Beck, in the first five years after impoundment.  
 47 *Hydrobiologia* 58:145–156
- 48 Arias-Andrés M, Rämö R, Mena Torres F, Ugalde R, Grandas L, Ruepert C, Castillo LE, van den Brink PJ,  
 49 Gunnarsson JS (2016) Lower tier toxicity risk assessment of agriculture pesticides detected on the Rio Madre  
 50 de Dios watershed, Costa Rica *Environ Sci Pollut Res* DOI 10.1007/s11356-016-7875-7 (this special issue)
- 51 Armitage PD, Moss D, Wright JF, Furse MY (1983) The performance of a new biological water quality score  
 52 system based on macroinvertebrates over a wide range of unpolluted running-water sites. *Water*  
 53 *Research* 17:333–347




- 1 Barbour MT, Gerritsen J, Snyder BD, Stribling JB (1999) Rapid Bioassessment Protocols for Use in Streams and  
 2 Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish. 2nd ed. EPA 841-B-99-002.  
 3 Washington, D.C.: U.S. Environmental Protection Agency, Office of Water
- 4 Beisel JN, Usseglio-Polatera P, Moreteau JC (2000) The spatial heterogeneity of a river bottom: a key factor  
 5 determining macroinvertebrate communities. *Hydrobiologia* 422/423:163–171
- 6 Bellamy AS (2013) Banana production systems: identification of alternative systems for more sustainable  
 7 production. *Ambio* 42:334–343
- 8 Bravo V, de la Cruz E, Herrera G, Ramírez F (2013) Agricultural pesticide use as tool for  
 9 monitoring health hazards. *Uniciencia* 27, 351-376
- 10 Bray RJ, Curtis JT (1957) An ordination of the upland forest communities of southern  
 11 Wisconsin. *Ecol Monogr* 27: 325-349
- 12 Castillo LE, Martínez E, Ruepert C, Savage C, Gilek M, Pinnock M, Solis E (2006) Water quality and  
 13 macroinvertebrate community response following pesticide applications in a banana plantation, Limon,  
 14 Costa Rica. *Science of the total environment* 367:418–432
- 15 Castillo LE, Ruepert C, Solis E (2000) Pesticide residues in the aquatic environment of banana plantation areas  
 16 in the north Atlantic zone of Costa Rica. *Environmental Toxicology and Chemistry*, Vol. 19, 8: 1942–1950
- 17 Chapin FS, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, Hooper DU, Lavorel S, Sala OE,  
 18 Hobbie SE, Mack MC, Díaz S (2000) Consequences of changing biodiversity. *Nature* 405: 234–242
- 19 Cuppen JGM, Van den Brink PJ, Camps E, Uil KF, Brock TCM (2000) Impact of the fungicide carbendazim in  
 20 freshwater microcosms. I. Water quality, breakdown of particulate organic matter and responses of  
 21 macroinvertebrates. *Aquatic Toxicology* 48:233–250
- 22 Diaz RJ (1992) Ecosystem assessment using estuarine and marine benthic community structure. In: Burton GAJ,  
 23 editor. *Sediment toxicity assessment*: Lewis Publishers, BocaRaton, FL pp 67–85
- 24 Diepens NJ, Pfennig S, Van den Brink PJ, Gunnarsson JS, Ruepert C, Castillo LE (2014) Effect of pesticides  
 25 used in banana and pineapple plantations on aquatic ecosystems in Costa Rica. *Journal of Environmental*  
 26 *Biology*, 35: 73-84
- 27 Echeverría-Sáenz S, Mena F, Arias-Andrés M, Vargas S, Ruepert C, van den Brink PJ, Castillo LE, Gunnarsson  
 28 (2016) In situ toxicity and ecological risk of agro-pesticide runoff in the Madre de Dios River in Costa Rica  
 29 JS. *Environ Sci Pollut Res*. DOI 10.1007/s11356-016-7817-4 (this special issue)
- 30 Friedrich G, Chapman D, Beim A (1996) The use of biological material. In: Chapman D, editor. *Water Quality*  
 31 *Assessments. A Guide to the Use of Biota, Sediments and Water in Environmental Monitoring*. 2nd ed.  
 32 Chapman & Hall, London
- 33 Lenat DR (1988) Water quality assessment using a qualitative collection method for benthic macroinvertebrates.  
 34 *J.N. Am. Benthological Soc.* 7: 222-233
- 35 Liess M, Brown C, Dohmen P, Duquesne S, Hart A, Heimbach F, Krueger J, Lagadic L, Maund S, Reinert W,  
 36 Streløke M, Tarazona JV. Editors (2003). *Effects of Pesticides in the field*. EU & SETAC Europe Workshop.  
 37 SETAC, Le Croisic, France; 136p
- 38 Malmqvist B, Hoffsten PO (2000) Macroinvertebrate taxonomic richness, community structure and nestedness  
 39 in Swedish streams. *Archiv für Hydrobiologie* 150:29–54
- 40 Mustow SE (2002) Biological monitoring of rivers in Thailand: use and adaptation of the BMWP score.  
 41 *Hydrobiologia* 479:191–229
- 42 National Water Council (1981) *River quality: the 1980 survey and future outlook*. National Water Council,  
 43 London 39 p
- 44 Paaby P, Ramírez A, Pringle CM (1998) The benthic macroinvertebrate community in Caribbean Costa Rican  
 45 streams and the effect of two sampling methods. *Revista de biología tropical* 46: 185-199
- 46 Pearson TH, Rosenberg R (1978) Macrobenthic succession in relation to organic enrichment and pollution of the  
 47 marine environment. *Oceanogr Mar Biol Annu Rev* 16:229-311
- 48 Pollard JE (1981) Investigator differences associated with a kicking method for sampling macroinvertebrates.  
 49 *Journal of Freshwater Ecology* 1:215–224
- 50 Polidoro BA, Morra MJ (2016) An ecological risk assessment of pesticides  
 51 and fish kills in the Sixaola watershed, Costa Rica. *Environ Science and Pollution Research International*
- 52 Pringle CM, Ramirez A (1998) Use of both benthic and drift sampling techniques to assess tropical stream  
 53 invertebrate communities along an altitudinal gradient, Costa Rica. *Freshwater Biology* 39:359–373
- 54 Rämö R, van den Brink PJ, Ruepert C, Castillo LE, Gunnarsson JS (2016) Environmental risk assessment of  
 55 pesticides in the River Madre de Dios, Costa Rica using PERPEST, SSD and msPAF models (2016) *Environ*  
 56 *Sci Pollut Res*. DOI 10.1007/s11356-016-7375-9 (this special issue)
- 57 Rico A, Van den Brink PJ (2015) Evaluating aquatic invertebrate vulnerability to insecticides based on intrinsic  
 58 sensitivity, biological traits and toxic mode-of-action. *Environmental Toxicology and Chemistry* 34: 1907–  
 59 1917

- 1 Roldán Pérez G (1992) Guía para el estudio de los macroinvertebrados acuáticos del departamento de Antioquia.  
2 Colombia, Colciencias, Universidad de Antioquia; Bogotá, Colombia 217 p
- 3 Schreinemachers P, Tipraqsa P (2012) Agricultural pesticides and land use intensification in high, middle and  
4 low income countries, *Food Policy*. 37: 616-626
- 5 Springer M, Vásquez D, Castro A, Kohlmann B (2007) Bioindicadores de la calidad del agua. Earth University,  
6 Guacimo, Costa Rica
- 7 Ter Braak CJF (1995) Ordination. In: Jongman RGH, Ter Braak CJF, van Tongeren OFR, editors. *Data Analysis*  
8 *in Community and Landscape Ecology*. Cambridge University Press, Cambridge, UK 91–173
- 9 Ter Braak CJF, Smilauer P (2012) *Canoco Reference Manual and User's Guide: Software for Ordination*  
10 (Version 5.0). Microcomputer Power, Ithaca, NY, USA, 496 pp
- 11 Thorp JH, Covich AP (1991) editors. *Ecology and classification of North American freshwater invertebrates*. 1st  
12 ed. Academic Press, San Diego, CA 911 p
- 13 Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980) The river continuum concept.  
14 *Canadian Journal of Fisheries and Aquatic Sciences* 37:130–137
- 15 Van den Brink PJ, Hattink J, Bransen F, Van Donk E, Brock TCM (2000) Impact of the fungicide carbendazim  
16 in freshwater microcosms. II. Zooplankton, primary producers and final conclusions. *Aquatic Toxicology* 48:  
17 251-264
- 18 Van den Brink PJ, van den Brink NW, Ter Braak CJF (2003) Multivariate analysis of ecotoxicological data  
19 using ordination: demonstrations of utility on the basis of various examples. *Australasian Journal of*  
20 *Ecotoxicology* 9:141–156
- 21 van Hoey G, Borja A, Birchenough S, Buhl-Mortensen L, Degraer S, Fleischer D, Kerckhof F, Magni P, Muxika  
22 I, Reiss H, Schröder A, Zettler ML (2010) The use of benthic indicators in Europe: From the Water  
23 Framework Directive to the Marine Strategy Framework Directive. *Mar Poll Bull.* 60 (12) 2187–2196
- 24 van Wijngaarden RPA, Van den Brink PJ, Oude Voshaar JH, Leeuwangh P (1995) Ordination techniques for  
25 analyzing response of biological communities to toxic stress in experimental ecosystems. *Ecotoxicology*  
26 4:61–77
- 27 Verbruggen EMJ, van den Brink PJ (2010) Review of recent literature concerning mixture toxicity of pesticides  
28 to aquatic organisms. RIVM report 601400001/2010. National Institute for Public Health and the  
29 Environment, Bilthoven, The Netherlands: 34 p
- 30 von der Ohe P C, Goedkoop W (2013) Distinguishing the effects of habitat degradation and pesticide stress on  
31 benthic invertebrates using stressor-specific metrics. *Science of The Total Environment*: 444: 480–490
- 32 Walter H, Consolaro F, Gramatica P, Scholze M, Altenburger R (2002) Mixture toxicity of priority pollutants at  
33 no observed effect concentrations (NOECs). *Ecotoxicology* 11:299–310
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48

1 **Figures and Tables with captions:**

2




3

4 **Fig 1:** Map of Costa Rica, showing the location of the 13 sampling sites and rivers sampled. White and black  
 5 squares denote upstream and downstream sites respectively. RS= Río Sucio, RPc= Río Pacuare, SST= Small  
 6 stream in Talamanca, SSS= Small stream in Sarapiquí, SSG= Small stream in Guácimo and Rpm= Río  
 7 Parismina. For GPS coordinates and description of sites and farming type at each site see Table1. For number of  
 8 composite samples and mean values for different community structure descriptors see Table 2

9

10

11



12

13 **Fig 2:** PCA biplots showing the variation in taxa composition between the sites (Fig 2A) and the correlation  
 14 between the taxa and the measured explanatory variables (Fig 2B) Of the variation in taxa composition, 30% is  
 15 displayed on the horizontal axis and another 18% on the vertical axis. Analyses were performed on mean values  
 16 where more than 1 composite sample was taken

17

1 **Table 1:** Sampling sites with GPS coordinates, a description of the site and some characteristics including the  
 2 distance from the source of the surface water, the type of substrate, the width of the river and the velocity of  
 3 water flow at the sampling site

Site	GPS coordinates	Water course	Location relative farms	Distance from river source (km)	Bottom substrate	River width (m)	Velocity (m/s)
RS1up	Sarapiquí, N 10° 20.320' W 83° 53.432'	Río Sucio	Upstream conventional farm	25	rocks,sand	11–20	0.2-0.4
RS2down	Sarapiquí, N 10° 21.600' W 83° 53.433'	Río Sucio	Downstream conventional farm	28	rocks,sand	6–10	0.2-0.4
RS3down	Sarapiquí, N 10° 20.819' W 83° 50.910'	Río Sucio	Downstream banana district	48	rocks,sand	11–20	0.5-0.8
RS4down	Sarapiquí, N 10° 29.151' W 84° 00.091'	Río Sucio accessed through Penjamo	Downstream banana district	85	sand/silt	>20	0.5-0.8
RPc1cup	Siquirres, N 10° 06.069' W 83° 28.967'	Río Pacuare	Upstream conventional farm	82.5	rocks,sand	>20	0.2-0.8
RPc2down	Siquirres, N 10° 07.166' W 83° 27.515'	Río Pacuare	Downstream conventional farm	86	rocks,sand	>20	0.2-0.4
RPc3down	Siquirres, N 10° 12.636' W 83° 19.328'	Río Pacuare	Downstream banana district	120	sand/silt	>20	0.5-0.8
SST1up	Talamanca, N 9° 41.800' W 82° 49.408'	Small stream between Cahuita and Hone Creek	Upstream small scale organic farm, Primary forest	0.5	rocks, leaf litter	1–2	0.2-0.4
SST2down	Talamanca, N 9° 42.105' W 82° 49.189'	Small stream between Cahuita and Hone Creek	Downstream small scale organic farm	1.25	rocks, leaf litter	<1	0.2-0.4
SSS1Nat	Sarapiquí, N 10° 12.919' W 83° 53.310'	Small stream east of Río Sucio, upstream bridge	Pristine, National park Braulio Carillo	1.75	rocks, sand, leaf litter	1–2	<0.2
SSS1up	Sarapiquí, N 10° 17.511' W 83° 53.310'	Small stream near Río San José	Upstream conventional farm	3	rocks, leaf litter	3–5	<0.2/0.2-0.4
SSG1down	Guácimo, N 10° 11.773' W 83° 35.657'	Small stream accessed through EARTH P4	Downstream less pesticide intensive farm (EARTH)	3	rocks, leaf litter	3–5	0.5-0.8
RPm1down	Guácimo, N 10° 13.519' W 83° 36.260'	Río Parismina accessed through EARTH P1	Downstream conventional farm (EARTH)	25	sand/silt	11–20	0.2-0.4

4

5

6 **Table 2:** Benthic community structure comparisons of upstream and downstream sites with mean values  
 7 (standard deviation within brackets). Each composite sample consists of 6 pooled kick-samples, corresponding to  
 8 approximately 1 square meter

Site	Water course/ Site relative banana farms	Composite samples	BMWP score	ASPT score	Taxa richness (S)	Abundance	Individuals/ Taxon	Shannon-Wiener (H')	Evenness (E)	% Dominating taxon	EPT Index
RS1up	Río Sucio	3	72.7	5.1	19	177	9.4	2.362	0.573	23.5	7.33
	Upstream conventional farm		(3.79)	(0.33)	(3)	(13.89)	(1.05)	(0.116)	(0.121)	(3.88)	(0.58)
RS2down	Río Sucio	3	55.0	5.2	13.3	257.6	19.2	1.796	0.461	33.3	6.68
	Downstream conventional farm		(5.00)	(0.32)	(0.58)	(139.23)	(10.26)	(0.228)	(0.112)	(9.35)	(0.58)
RS3down	Río Sucio	3	22.7	3.5	6	23.6	2.6	1.263	0.502	20.7	3.33
	Downstream banana district		(20.0)	(3.02)	(5.29)	(21.83)	(2.28)	(1.095)	(0.449)	(18.97)	(3.06)
RS4down	Río Sucio Downstream banana district	1	7	3.5	5	19	3.8	1.313	0.743	52.6	3
RPc1up	Río Pacuare	3	49	3.4	12.6	74.3	5.9	1.684	0.429	52.3	6.33
	Upstream conventional farm		(1.73)	(1.5)	(0.58)	(20.03)	(1.86)	(0.155)	(0.065)	(9.02)	(0.58)
RPc2down	Río Pacuare	3	44.3	5.1	11	236	21.5	1.666	0.487	41.2	5.33
	Downstream conventional farm		(4.73)	(0.23)	0	(120.40)	(10.95)	(0.191)	(0.087)	(12.03)	(1.15)
RPc3down	Río Pacuare	3	7.3	3.4	2.3	12.7	5.3	0.322	0.718	90.9	0.67
	Downstream banana district		(4.62)	(1.50)	(1.53)	(0.84)	(1.53)	(0.359)	(0.254)	(9.26)	(0.58)
SST1up	Small stream Upstream organic farm	1	84	6	18	80	4.4	2.294	0.551	26.2	9
SST2down	Small stream Downstream organic farm	1	43	4.8	16	57	3.62	2.261	0.600	27.6	4
SSS1Nat	Small stream National park Braulio Carillo	1	55	5.5	14	37	2.6	2.443	0.822	21.6	5
SSS1up	Small stream Upstream conventional farm	2	67	5.6	16.5	88	5.3	2.075	0.491	23.5	4
			(1.41)	(0.54)	(0.71)	(32.53)	(1.74)	(0.215)	(0.125)	(4.97)	(0)
SSG1down	Small stream Downstream low-input farm	1	35	5.8	10	32	3.2	2.018	0.753	31.2	3
RPm1down	Río Parismina Downstream large scale farm	2	28.5	4.7	7.5	72	9.4	1.366	0.530	47.5	2
			(7.78)	(0.18)	(0.71)	(39.80)	(4.39)	(0.151)	(0.129)	(9.26)	(0)

9