

Gamal A. El-Hiti*, Keith Smith, Mohammed B. Alshammari, Amany S. Hegazy and Benson M. Kariuki

Crystal structure of 1,1-dimethyl-3-(4-methoxyphenyl)urea, $C_{10}H_{14}N_2O_2$




Table 1: Data collection and handling.

Crystal:	Colorless block
Size:	0.29 × 0.13 × 0.09 mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	0.09 mm ⁻¹
Diffractometer, scan mode:	SuperNova, φ and ω
θ_{\max} , completeness:	29.6°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	5082, 2534, 0.021
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 1512
$N(\text{param})_{\text{refined}}$:	135
Programs:	CrysAlis ^{PRO} [17], SHELX [18], WinGX [19], CHEMDRAW Ultra [20].

DOI 10.1515/ncrs-2016-0238

Received August 2, 2016; accepted January 4, 2017; available online January 20, 2017

Abstract

$C_{10}H_{14}N_2O_2$, monoclinic, $P2_1/c$ (no. 14), $a = 14.9185(12)$ Å, $b = 7.7243(6)$ Å, $c = 9.2229(5)$ Å, $\beta = 91.032(6)^\circ$, $V = 1062.63(13)$ Å³, $Z = 4$, $T = 293(2)$ K.

CCDC no.: 1525571

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

To a stirred solution of triphosgene (1.0 mole equivalent) in dichloromethane (DCM), a solution of 4-methoxyaniline (2.5 mole equivalents) and triethylamine (5.5 mole equivalent)

lents) in DCM was slowly added over 30 min at 0 °C. The mixture was stirred at 0 °C for 2 h, after which a solution of dimethylamine (3.0 mole equivalents) in tetrahydrofuran was added. The reaction mixture was stirred at 0 °C for 1 h. The mixture was poured onto water and the DCM layer was separated, washed with water, and dried over magnesium sulfate and the solvent was then removed under reduced pressure. Crystallization of the obtained raw solid using a mixture of ethyl acetate and diethyl ether (1:3 by volume) gave the title compound, 1,1-dimethyl-3-(4-methoxyphenyl)urea (83%) as colorless crystals, Mp. 132–133 °C (lit. 131–132 °C [1]).

Experimental details

Non-hydrogen atoms were refined with anisotropic displacement parameters. The N-H proton was refined freely but all other hydrogen atoms were placed in calculated positions and refined using a riding model. C-H bonds of the methyl groups were fixed at 0.96 Å, with displacement parameters 1.5 times $U_{\text{eq}}(\text{C})$ for the hydrogen atoms, and were allowed to spin about the C-C bond. Aromatic C-H distances were set to 0.93 Å and their $U(\text{iso})$ values of the corresponding hydrogen atoms were set to 1.2 times the U_{eq} values of the atoms to which they are bonded.

Comment

Many urea derivatives are very important intermediates in organic syntheses and show a variety of biological activities

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Mohammed B. Alshammari: Chemistry Department, College of Sciences and Humanities, Prince Sattam bin Abdulaziz University, P.O. Box 83, Al-Kharij 11942, Saudi Arabia

© 2017 Gamal A. El-Hiti et al., published by De Gruyter.

This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 License.

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C1	0.24439(13)	0.3566(3)	0.26176(18)	0.0553(5)
C2	0.24067(15)	0.5161(3)	0.1958(2)	0.0665(6)
H2	0.2782	0.5391	0.1188	0.080*
C3	0.18216(16)	0.6430(3)	0.2417(2)	0.0710(6)
H3	0.1799	0.7496	0.1948	0.085*
C4	0.12752(14)	0.6108(3)	0.3569(2)	0.0670(6)
C5	0.13152(14)	0.4524(3)	0.4251(2)	0.0700(6)
H5	0.0953	0.4312	0.5042	0.084*
C6	0.18829(14)	0.3253(3)	0.3780(2)	0.0625(6)
H6	0.1893	0.2180	0.4237	0.075*
C7	0.35326(14)	0.1205(3)	0.29039(19)	0.0556(5)
C8	0.37929(19)	−0.0498(3)	0.0685(2)	0.0848(8)
H8A	0.3156	−0.0574	0.0529	0.127*
H8B	0.4063	−0.1596	0.0477	0.127*
H8C	0.4034	0.0370	0.0058	0.127*
C9	0.4621(2)	−0.1113(4)	0.2971(3)	0.1023(10)
H9A	0.4701	−0.0680	0.3940	0.153*
H9B	0.5185	−0.1095	0.2486	0.153*
H9C	0.4400	−0.2279	0.3005	0.153*
C10	0.0505(3)	0.8791(4)	0.3308(4)	0.1230(12)
H10A	0.1025	0.9520	0.3381	0.184*
H10B	−0.0002	0.9389	0.3695	0.184*
H10C	0.0387	0.8511	0.2309	0.184*
N1	0.30120(12)	0.2264(2)	0.20641(18)	0.0627(5)
N2	0.39827(13)	−0.0034(2)	0.21882(17)	0.0670(5)
O1	0.36005(11)	0.1371(2)	0.42292(13)	0.0739(5)
O2	0.06579(12)	0.7263(3)	0.40972(19)	0.0957(6)
H1	0.3151(16)	0.237(3)	0.117(3)	0.077(7)*
C1	0.24439(13)	0.3566(3)	0.26176(18)	0.0553(5)
C2	0.24067(15)	0.5161(3)	0.1958(2)	0.0665(6)
H2	0.2782	0.5391	0.1188	0.080*
C3	0.18216(16)	0.6430(3)	0.2417(2)	0.0710(6)
H3	0.1799	0.7496	0.1948	0.085*
C4	0.12752(14)	0.6108(3)	0.3569(2)	0.0670(6)
C5	0.13152(14)	0.4524(3)	0.4251(2)	0.0700(6)
H5	0.0953	0.4312	0.5042	0.084*

[2–5]. Various efficient procedures for the synthesis of aromatic ureas are known [6–11]. Aromatic ureas can be easily modified *via* treatment with lithium reagents followed by reactions with electrophiles, a methodology we have used in the last few years [12–16].

The asymmetric unit contains one molecule of C₁₀H₁₄N₂O₂. In the molecule, the methoxybenzene group is almost planar with a C3–C4–O2–C10 torsion angle of 75(4)°. The rest of the molecule is not planar with twist angles 46.90(9)° between the phenyl and the NC(O)N groups and 10.68(34)° between NC(O)N and N(Me₂) groups. In the crystal, N–H···O hydrogen bonding is observed with geometry: N1–H1···O1 = 158(2)°, N1···O1 = 2.966(2) Å. The hydrogen

bonds construct chains of molecules along [001]. All bond lengths and angles are in the expected ranges.

Acknowledgements: The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for its funding for this research through the research group project RGP-239 and to Cardiff University for the continued support.

References

- Hutchby, M.: *Novel Synthetic Chemistry of Ureas and Amides*. Springer Thesis, Springer, 2013.
- Chen, J.-N.; Wang, X.-F.; Li, T.; Wu, D.-W.; Fu, X.-B.; Zhang, G.-J.; Shen, X.-C.; Wang, H.-S.: Design, synthesis, and biological evaluation of novel quinazolinyl-diaryl urea derivatives as potential anticancer agents. *Eur. J. Med. Chem.* **107** (2016) 12–25.
- Kocuyigit-Kaymakcioglu, B.; Celen, A. O.; Tabanca, N.; Ali, A.; Khan, S. I.; Khan, I. A.; Wedge, D. E.: Synthesis and biological activity of substituted urea and thiourea derivatives containing 1,2,4-triazole moieties. *Molecules* **18** (2013) 3562–3576.
- Gennäs, G. B.; Mologni, L.; Ahmed, S.; Rajaratnam, M.; Marin, O.; Lindholm, N.; Viltadi, M.; Gambacorti-Passerini, C.; Scapozza, L.; Yli-Kauhaluoma, J.: Design, synthesis, and biological activity of urea derivatives as anaplastic lymphoma kinase inhibitors. *ChemMedChem* **6** (2011) 1680–1692.
- Wilson, A. A.; Garcia, A.; Houle, S.; Sadovski, O.; Vasdev, N.: Synthesis and application of isocyanates radiolabeled with carbon-11. *Chem. Eur. J.* **17** (2011) 259–264.
- Thalluri, K.; Manne, S. R.; Dev, D.; Mandal, B.: Ethyl 2-cyano-2-(4-nitrophenylsulfonyloxymino)acetate-mediated Lossen rearrangement: single-pot racemization-free synthesis of hydroxamic acids and ureas from carboxylic acids. *J. Org. Chem.* **79** (2014) 3765–3775.
- Carnaroglio, D.; Martina, K.; Palmisano, G.; Penoni, A.; Domini, C.; Cravotto, G.: One-pot sequential synthesis of isocyanates and urea derivatives via a microwave-assisted Staudinger–aza-Wittig reaction. *Beilstein J. Org. Chem.* **9** (2013) 2378–2386.
- Vinogradova, E. V.; Fors, B. P.; Buchwald, S. L.: Palladium-catalyzed cross-coupling of aryl chlorides and triflates with sodium cyanate: a practical synthesis of unsymmetrical ureas. *J. Am. Chem. Soc.* **134** (2012) 11132–11135.
- Wu, C.; Cheng, H.; Liu, R.; Wang, Q.; Hao, Y.; Yu, Y.; Zhao, F.: Synthesis of urea derivatives from amines and CO₂ in the absence of catalyst and solvent. *Green Chem.* **12** (2010) 1811–1816.
- Mizuno, T.; Nakai, T.; Mihara, M.: Synthesis of unsymmetrical ureas by sulfur-assisted carbonylation with carbon monoxide and oxidation with molecular oxygen under mild conditions. *Synthesis* **41** (2009) 2492–2496.
- Artuso, E.; Degani, I.; Fochi, R.; Magistris, C.: Preparation of mono-, di-, and trisubstituted ureas by carbonylation of aliphatic amines with *S,S*-dimethyl dithiocarbonate. *Synthesis* **39** (2007) 3497–3506.

12. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Directed lithiation of *N'*-(2-(4-methoxyphenyl)ethyl)-*N,N*-dimethylurea and *tert*-butyl (2-(4-methoxyphenyl)ethyl)carbamate. *Synthesis* **46** (2014) 394–402.
13. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Control of site of lithiation of 3-(aminomethyl)pyridine derivatives. *Synthesis* **45** (2013) 3426–3434.
14. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Lithiation and substitution of *N'*-(ω -phenylalkyl)-*N,N*-dimethylureas. *Synthesis* **44** (2012) 20139–2022.
15. Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Variation in the site of lithiation of 2-(2-methylphenyl)ethanamine derivatives. *J. Org. Chem.* **77** (2012) 11210–11215.
16. Smith, K.; El-Hiti, G. A.; Hegazy, A. S.: Lateral lithiation of *N'*-(2-methylbenzyl)-*N,N*-dimethylurea and *N*-(2-methylbenzyl)pivalamide: Synthesis of tetrahydroisoquinolines. *Synthesis* **42** (2010) 1371–1380.
17. Agilent. CrysAlis^{PRO}. Agilent Technologies, Yarnton, England, 2014.
18. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
19. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
20. Cambridge Soft. CHEMDRAW Ultra. Cambridge Soft Corporation, Cambridge, Massachusetts, USA, 2001.