

Pinching of the first eigenvalue for second order operators on hypersurfaces of the Euclidean space

Julien Roth · Julian Scheuer

Received: date / Accepted: date

Abstract We prove stability results associated with upper bounds for the first eigenvalue of certain second order differential operators of divergence-type on hypersurfaces of the Euclidean space. We deduce some applications to r -stability as well as to almost-Einstein hypersurfaces.

Keywords Reilly-type inequality · Eigenvalue pinching · Stable hypersurfaces · Almost-Einstein estimates

Mathematics Subject Classification (2000) 35P15 · 53C20 · 53C24 · 53C42 · 58C40

1 Introduction

There is a wide literature concerning estimates of the eigenvalues of the Laplacian (and more general divergence-type second order operators) on submanifolds of spaceforms. The first upper bound for the first eigenvalue of the Laplacian on compact hypersurfaces of \mathbb{R}^{n+1} was obtained by Bleeker and Weiner [10] who showed

$$\lambda_1 \leq \frac{1}{V(M)} \int_M |B|^2 dv_g, \quad (1)$$

where B is the second fundamental form of the hypersurface M and $V(M)$ its volume. After that, Reilly [27] improved this upper bound by getting the norm

J. Roth
Laboratoire d'Analyse et de Mathématiques Appliquées, UPEM-UPEC, CNRS, F-77454
Marne-la-Vallée, France
E-mail: julien.roth@u-pem.fr

J. Scheuer
Albert-Ludwigs-Universität, Mathematisches Institut, Eckerstr. 1, 79104 Freiburg, Germany
E-mail: julian.scheuer@math.uni-freiburg.de

of the mean curvature instead of the second fundamental form. Precisely, he proved the following estimate:

$$\lambda_1 \leq \frac{n}{V(M)} \int_M |H|^2 dv_g. \quad (2)$$

For these two inequalities the limiting case is attained if and only if the hypersurface is a hypersphere. Later on, Heintze [21] extended Reilly's upper bound to hypersurfaces of compact ambient spaces and, after a partial result due to Heintze, El Soufi and Ilias [14] proved an analogue in the hyperbolic space, also cf. [2]. In fact, in [27] Reilly proved a sequence of upper bounds involving higher order mean curvatures, for which (2) is just the particular case $r = 0$:

$$\lambda_1 \left(\int_M H_r dv_g \right)^2 \leq nV(M) \int_M H_{r+1}^2 dv_g. \quad (3)$$

As the previous ones the inequalities are sharp and the limiting cases are also characterised by geodesic hyperspheres. Note that the inequalities in (3) are not in general better than (2); they are not comparable to the other.

The Laplacian is not the only fundamental operator for which extrinsic estimates have been proved. In particular comparable upper bounds have been established for some divergence-type second order elliptic operators as the operators L_r associated with the higher order mean curvatures H_r and even for more general elliptic divergence-free operators, cf. [18]. Notations and basic facts about higher order mean curvatures will be given in Section 2.

The aim of the present article is first to study the stability of the limiting case for a sequence of optimal upper bounds for a larger class of second order operators including the Laplacian and the operators L_r . In [32] the first author proved the the following theorem.

Theorem 1 ([32]) *Let (M^n, g) be a connected, oriented and closed Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} . Assume that M is endowed with two symmetric and divergence-free $(1, 1)$ -tensors S and T . Assume in addition that T is positive definite. Then the first positive eigenvalue of the operator L_T satisfies*

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \leq \left(\int_M \operatorname{tr}(T) dv_g \right) \left(\int_M |H_S|^2 dv_g \right). \quad (4)$$

Moreover, if H_S does not vanish identically and equality occurs, then M is a hypersphere and $\operatorname{tr}(S)$ is constant.

Here the operator L_T is given by

$$L_T u := -\operatorname{div}(T \nabla u) \quad (5)$$

for functions $u \in C^2(M)$ and H_S is the generalised mean curvature vector

$$H_S = S^{ij} X_{ij}, \quad (6)$$

where X is the immersion vector, cf. Section 2 for details. Since the equality is achieved only for hyperspheres, it is natural to study the stability of the limiting case. We give the following quantitative result about the proximity to hyperspheres for hypersurfaces almost satisfying the equality case. Namely, under the condition

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \geq (1 - \varepsilon) \left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^2 V(M), \quad (7)$$

for $p > 1$, we prove the following result.

Theorem 2 *Let $n \geq 2$ and (M^n, g) be a connected, oriented and closed Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} by X . Assume that M is endowed with two symmetric and divergence-free $(1, 1)$ -tensors S and T . Assume in addition that T is positive definite, $S^{ij} B_{ij}$ is not identically zero and that for some $q > n$ there holds $V(M) \|B\|_q^n \leq A$. Let $p > 1$. Then there exists $\varepsilon_0 = \varepsilon_0(n, p, q, A) > 0$, such that if (7) holds with $\varepsilon < \varepsilon_0$, then M is ε^α -close, diffeomorphic and ε^α -almost isometric to a sphere $S(\bar{X}, r)$, i.e. there exists $r > 0$, a constant C depending on n, p, q and A , and $\alpha = \alpha(n, q)$, such that*

$$\| |X - \bar{X}| - r \|_\infty \leq Cr\varepsilon^\alpha, \quad (8)$$

where $\bar{X} = V(M)^{-1} \int_M X$ is the center of mass of $X(M)$, and for a natural diffeomorphism

$$F: (M, d_1) \rightarrow (S(r), d_2) \quad (9)$$

we have

$$|d_2(F(x_1), F(x_2)) - d_1(x_1, x_2)| \leq Cr\varepsilon^\alpha \quad \forall x_1, x_2 \in M. \quad (10)$$

Moreover, M is embedded and $X(M)$ is a starshaped hypersurface.

Remark 1 (i) In order to prove (8), it is not necessary that $V(M) \|B\|_q^n \leq A$, but only $V(M) \|H\|_q^n \leq A$.

(ii) The radius is given by

$$r = \frac{1}{V(M)} \frac{|\int_M \operatorname{tr}(S) dv_g|}{\|H_S\|_{2p}}. \quad (11)$$

In the sections 5 and 6 we derive some applications of Theorem 2. The first of those concerns the r -stability. Namely we prove that if a hypersurface with constant r -th mean curvature is almost stable in a suitable sense, then it is a geodesic sphere (see Theorem 4). Finally, the last section of the paper will be devoted to almost-Einstein hypersurfaces, where we improve previous closeness results of the first author [29] and Hu, Xu and Zhao [24].

Earlier results on the eigenvalue pinching problem can be found in [12] and [29] for the Euclidean case, [19] for other ambient spaces and [24] for an improvement of parameters compared to [29].

2 Preliminaries

2.1 Generalised Hsiung-Minkowski formula

Let (M^n, g) be a connected, oriented and closed Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} . We denote by X its vector position, by ν its unit normal vector and we consider a divergence-free symmetric $(1, 1)$ -tensor T on M . We associate to T the second order differential operator L_T defined by

$$L_T u := -\operatorname{div}(T\nabla u),$$

for any C^2 -function u on M . We also associate to T the following generalised mean curvature vectors:

$$H_T = T^{ij} X_{ij}, \quad (12)$$

where X_{ij} denotes the second covariant derivative of the immersion vector X . Then we have

$$L_T X = -H_T \quad (13)$$

and easily deduce the following identity:

$$\frac{1}{2} L_T |X|^2 = -\langle X, H_T \rangle - \operatorname{tr}(T). \quad (14)$$

After integrating this gives the so-called generalised Hsiung-Minkowski formula

$$\int_M \left(\langle X, H_T \rangle + \operatorname{tr}(T) \right) dv_g = 0. \quad (15)$$

The classical Hsiung-Minkowski formula is in fact this formula for the particular case where T is the identity and more generally, where T is the tensor T_r associated to the r -th mean curvature. In the next paragraph we will recall the definition of T_r and the associated operator L_r , which play a crucial role for the r -stability as we will see in Section 5.

2.2 Higher order mean curvatures

The higher order mean curvatures are extrinsic quantities defined from the second fundamental form and generalising the notion of mean curvature. Up to a normalisation constant the mean curvature H is the trace of the second fundamental form B :

$$H = \frac{1}{n} \operatorname{tr}(B), \quad (16)$$

where we use the convention

$$X_{ij} = -B_{ij}\nu. \quad (17)$$

In other words the mean curvature is

$$H = \frac{1}{n} S_1(\kappa_1, \dots, \kappa_n), \quad (18)$$

where S_1 is the first elementary symmetric polynomial and $\kappa_1, \dots, \kappa_n$ are the principal curvatures. Higher order mean curvatures are defined in a similar way for $r \in \{1, \dots, n\}$ by

$$H_r = \frac{1}{\binom{n}{r}} S_r(\kappa_1, \dots, \kappa_n), \quad (19)$$

where S_r is the r -th elementary symmetric polynomial, that is for any n -tuple (x_1, \dots, x_n) ,

$$S_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}. \quad (20)$$

By convention we set $H_0 = 1$ and $H_{n+1} = 0$. Finally, for convenience we also set $H_{-1} = -\langle X, \nu \rangle$.

To each H_r we associate a symmetric $(2, 0)$ -tensor, which is in coordinates given by

$$T_r = (T_r^{ij}) = \left(\frac{\partial S_{r+1}}{\partial B_{ij}} \right), \quad (21)$$

where S_{r+1} is now understood to depend on the second fundamental form and the metric. The relation between these two notions can be found in [16, Sec. 2.1] for example. These tensors T_r are divergence-free (see [17] for instance) and satisfy the following relations:

$$\text{tr}(T_r) = c(r)H_r \quad \text{and} \quad H_{T_r} = -c(r)H_{r+1}\nu, \quad (22)$$

where $c(r) = (n-r)\binom{n}{r}$ and H_{T_r} is given by the relation (12).

We finish this section by giving some classical inequalities between the H_r which are well-known. First, for any $r \in \{0, \dots, n-2\}$,

$$H_r H_{r+2} \leq H_{r+1}^2, \quad (23)$$

with equality at umbilical points, cf. [20, p. 104]. Moreover, cf. [7], if $H_{r+1} > 0$, then $H_s > 0$ for any $s \in \{0, \dots, r\}$ and

$$H_{r+1}^{\frac{1}{r+1}} \leq H_r^{\frac{1}{r}} \leq \dots \leq H_2^{\frac{1}{2}} \leq H. \quad (24)$$

Combining (23) and (24), we get that if $H_{r+1} > 0$, then

$$H_{r+2} \leq H H_{r+1}, \quad (25)$$

with equality at umbilical points.

2.3 Extrinsic Sobolev inequality

We finish this section of preliminaries by recalling the extrinsic Sobolev inequality proved by Michael and Simon [25] for submanifolds of the Euclidean space and by Hoffman and Spruck [23] for other ambient spaces. This inequality will play a crucial role in the iteration process to obtain L^∞ -estimates in the proof of Theorem 2. For any non-negative C^1 -function f on a hypersurface M of \mathbb{R}^{n+1} we have

$$\left(\int_M f^{\frac{n}{n-1}} dv_g \right)^{\frac{n-1}{n}} \leq K_n \int_M (|\nabla f| + |H|f) dv_g, \quad (26)$$

where K_n is a positive constant depending only on the dimension n and where H is the mean curvature of M . An immediate consequence of this inequality is that we can bound the volume of M in terms of the mean curvature from below. Indeed we have $V(M)^{\frac{n-1}{n}} \leq K_n \|H\|_\infty V(M)$, obtained by taking $f \equiv 1$ in (26), which gives the following lower bound for the volume:

$$V(M) \geq \frac{1}{(K_n \|H\|_\infty)^n}. \quad (27)$$

Finally we fix our convention for the L^p -norms. For $p \in [1, \infty)$ and a function f defined on M we set

$$\|f\|_p = \left(\frac{1}{V(M)} \int_M |f|^p dv_g \right)^{\frac{1}{p}}. \quad (28)$$

3 A general upper bound for $\lambda_1(L_T)$

We first recall the general upper bound for the first eigenvalue of the operator L_T in terms of a symmetric divergence-free $(1, 1)$ -tensor S .

Theorem 3 ([32]) *Let (M^n, g) be a connected, oriented and closed Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} . Assume that M is endowed with two symmetric and divergence-free $(1, 1)$ -tensors S and T . Assume in addition that T is positive definite. Then the first positive eigenvalue of the operator L_T satisfies*

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \leq \left(\int_M \operatorname{tr}(T) dv_g \right) \left(\int_M |H_S|^2 dv_g \right). \quad (29)$$

Moreover, if H_S does not vanish identically and equality occurs, then M is a hypersphere and $\operatorname{tr}(S)$ is constant.

Proof: First we recall that $L_T X = -H_T$. Without loss of generality let

$$\int_M X = 0. \quad (30)$$

Thus we may use the coordinates X^α as test functions in the Rayleigh quotient to obtain

$$\begin{aligned} \lambda_1(L_T) \int_M |X|^2 dv_g &\leq - \int_M \langle X, H_T \rangle dv_g \\ &= \int_M \operatorname{tr}(T) dv_g, \end{aligned} \quad (31)$$

where we have used the generalised Hsiung-Minkowski formula (15) for the tensor T . Hence, we deduce that

$$\begin{aligned} \left(\int_M \operatorname{tr}(T) dv_g \right) \left(\int_M |H_S|^2 dv_g \right) &\geq \lambda_1(L_T) \left(\int_M |X|^2 dv_g \right) \left(\int_M |H_S|^2 dv_g \right) \\ &\geq \lambda_1(L_T) \left(\int_M |X| |H_S| dv_g \right)^2 \\ &\geq \lambda_1(L_T) \left(\int_M \langle X, H_S \rangle dv_g \right)^2 \\ &= \lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2, \end{aligned} \quad (32)$$

where we have used the Cauchy-Schwarz inequality and the generalised Hsiung-Minkowski formula for the tensor S .

Let's study the limiting case. If H_S does not vanish identically, the integral of $\operatorname{tr}(S)$ is not zero in the equality case. Hence we have equality in the Cauchy-Schwarz inequality, that is, X and H_S are colinear. In particular the position vector is normal. We deduce easily that M is a sphere. Indeed we have

$$\frac{\partial}{\partial \xi^i} \langle X, X \rangle = 2 \langle X_i, X \rangle = 0. \quad (33)$$

Thus the norm of X is constant and so M is contained in a sphere. Since M has no boundary, M is the entire sphere. Moreover M is totally umbilic and so by definition of H_S we have $H_S = -k \operatorname{tr}(S) \nu$. Finally $\operatorname{tr}(S) = k^{-1} |H_S|$ is constant, since by equality in the Cauchy-Schwarz inequality H_S and X are proportional. \square

Remark 2 1. Note that for the equality case there is no equivalence. Indeed, if equality holds, then M is necessarily a geodesic sphere, but there is no reason that equality occurs if M is a geodesic sphere.
2. If T is the identity, that is L_T is the Laplacian, we have an alternative proof of the limiting case. If equality holds, then the coordinates are eigenvectors of the Laplacian and M is a geodesic sphere by Takahashi's Theorem [34].

The classical inequalities are particular cases of this general one. First of all, for $T = \operatorname{Id}$ we get the Reilly inequalities (2) and (3) cited above by taking $S = \operatorname{Id}$ or $S = T_r$. For $T = T_r$, $0 \leq r \leq n-1$, we recover the following

inequalities for the first eigenvalue of the operator L_r obtained by Alias and Malacarne [4]:

$$\lambda_1(L_r) \left(\int_M H_s dv_g \right)^2 \leq c(r) \int_M H_r dv_g \int_M H_{s+1}^2 dv_g, \quad (34)$$

with $S = T_s$. The special case $r = s$ gives a well known inequality proved by Alencar, do Carmo and Rosenberg [3]:

$$\lambda_1(L_r) \int_M H_r dv_g \leq c(r) \int_M H_{r+1}^2 dv_g. \quad (35)$$

Another particular case of our general inequality is the case $S = T$. We get the following Corollary.

Corollary 1 *Let (M^n, g) be a closed, connected and oriented Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} . Assume that M is endowed with a divergence-free and positive definite symmetric $(1, 1)$ -tensor T . Then, the first positive eigenvalue of the operator L_T satisfies*

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(T) dv_g \right) \leq \int_M |H_T|^2 dv_g. \quad (36)$$

Moreover, if equality holds, then M is a geodesic sphere and $\operatorname{tr}(T)$ is constant.

For $T = \operatorname{Id}$ we get the Reilly inequality (2) and for $T = T_r$ we have inequality (35).

4 A pinching result for $\lambda_1(L_T)$

In this section we consider the pinching problem associated with (29), which is the stability of its equality case. In other words, if equality is almost achieved, M is close to a sphere in a sense to be made precise. For technical reasons we will consider a less sharp but nevertheless optimal inequality which admits the same equality case:

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \leq \left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^2 V(M), \quad (37)$$

for $p > 1$, obtained from (29) by Hölder inequality. We introduce the following pinching condition for $0 < \varepsilon < 1$:

$$\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \geq (1 - \varepsilon) \left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^2 V(M). \quad (A_{p,\varepsilon})$$

4.1 L^2 -estimates

In this section we prove some lemmata giving proximity in an L^2 -sense between the hypersurface M and a geodesic sphere of appropriate radius under the assumptions of Theorem 2.

Lemma 1 *If $(A_{p,\varepsilon})$ is satisfied and $\bar{X} = 0$, then*

$$\frac{\int_M \operatorname{tr}(T) dv_g}{\lambda_1(L_T)V(M)}(1-\varepsilon)^2 \leq \|X\|_2^2 \leq \frac{\int_M \operatorname{tr}(T) dv_g}{\lambda_1(L_T)V(M)} \quad (38)$$

and

$$\frac{\left(\frac{1}{V(M)} \int_M \operatorname{tr}(S) dv_g\right)^2}{\|H_S\|_{2p}^2} (1-\varepsilon)^2 \leq \|X\|_2^2 \leq \frac{1}{1-\varepsilon} \frac{\left(\frac{1}{V(M)} \int_M \operatorname{tr}(S) dv_g\right)^2}{\|H_S\|_{2p}^2}. \quad (39)$$

Proof: We start from (31) obtained by injecting the coordinate functions into the Rayleigh quotient. This gives the upper bound for $\|X\|_2^2$ in (38). For the lower bound we start from (31) again. We have

$$\begin{aligned} & \lambda_1(L_T) \int_M |X|^2 dv_g \left(\int_M \operatorname{tr}(S) dv_g \right)^4 \\ & \leq \int_M \operatorname{tr}(T) dv_g \left(\int_M \operatorname{tr}(S) dv_g \right)^4 \\ & = \int_M \operatorname{tr}(T) dv_g \left(\int_M \langle H_S, X \rangle dv_g \right)^4 \\ & \leq \int_M \operatorname{tr}(T) dv_g \left(\int_M |H_S|^2 dv_g \right)^2 \left(\int_M |X|^2 dv_g \right)^2. \end{aligned} \quad (40)$$

Hence we get

$$\|X\|_2^2 \geq \frac{\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^4}{\left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^4 V(M)^3}. \quad (41)$$

Now we use $(A_{p,\varepsilon})$, which gives

$$\lambda_1(L_T)^2 \geq \frac{\left(\int_M \operatorname{tr}(T) dv_g \right)^2 \|H_S\|_{2p}^4 V(M)^2}{\left(\int_M \operatorname{tr}(S) dv_g \right)^4} (1-\varepsilon)^2 \quad (42)$$

and, together with (41), yields the desired result,

$$\frac{\int_M \operatorname{tr}(T) dv_g}{\lambda_1(L_T)V(M)}(1-\varepsilon)^2 \leq \|X\|_2^2. \quad (43)$$

The upper estimate of (39) follows from inserting $(A_{p,\varepsilon})$ into (38), the lower one from inserting (37) into (38). \square

Now we state a second lemma which gives an L^2 -estimate for the tangential part of the position vector under the pinching condition.

Lemma 2 *If $(A_{p,\varepsilon})$ holds and $\bar{X} = 0$, then*

$$\|X^T\|_2^2 \leq \varepsilon \|X\|_2^2. \quad (44)$$

Proof: As in the previous lemma we start from (31). We have

$$\begin{aligned} & \lambda_1(L_T) \int_M |X|^2 dv_g \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \\ & \leq \int_M \operatorname{tr}(T) dv_g \left(\int_M \operatorname{tr}(S) dv_g \right)^2 \\ & = \int_M \operatorname{tr}(T) dv_g \left(\int_M \langle H_S, X \rangle dv_g \right)^2 \\ & \leq \int_M \operatorname{tr}(T) dv_g \left(\int_M |H_S|^2 dv_g \right) \left(\int_M \langle X, \nu \rangle^2 dv_g \right) \\ & \leq \int_M \operatorname{tr}(T) dv_g \left(\int_M \langle X, \nu \rangle^2 dv_g \right) \|H_S\|_{2p}^2 V(M). \end{aligned} \quad (45)$$

From this we deduce that

$$\begin{aligned} \|X^T\|_2^2 &= \frac{1}{V(M)} \int_M (|X|^2 - \langle X, \nu \rangle^2) dv_g \\ &\leq \|X\|_2^2 - \frac{\lambda_1(L_T) \|X\|_2^2 \left(\int_M \operatorname{tr}(S) dv_g \right)^2}{\left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^2 V(M)} \\ &= \|X\|_2^2 \left(1 - \frac{\lambda_1(L_T) \left(\int_M \operatorname{tr}(S) dv_g \right)^2}{\left(\int_M \operatorname{tr}(T) dv_g \right) \|H_S\|_{2p}^2 V(M)} \right) \\ &\leq \varepsilon \|X\|_2^2, \end{aligned} \quad (46)$$

by using $(A_{p,\varepsilon})$ for the last line. \square

4.2 From L^2 to L^∞

Now we will give a sequence of lemmas, based on an iteration process, which allow us to control the L^∞ -norm of some functions by their L^2 -norm. Note that this iteration process does not depend on the pinching condition. We have the following lemma for the norm of the position vector. The proof can be found in [24, Lemma 5]. This lemma is an improvement of a similar lemma given in [19] and [31].

Lemma 3 [24, Lemma 5] *Let $q > n$ be a real number. There exists a constant $\Gamma(n, q) > 0$, so that for any isometrically immersed, compact submanifold M^n of \mathbb{R}^{n+1} we have*

$$\begin{aligned} & \| |X - \bar{X}| - \|X - \bar{X}\|_2 \|_\infty \\ & \leq \Gamma(V(M) \|H\|_q^n)^{\frac{\gamma}{n}} \|X - \bar{X}\|_2 \left(1 - \frac{\|X - \bar{X}\|_1}{\|X - \bar{X}\|_2} \right)^{\frac{1}{2(\gamma+1)}}, \end{aligned} \quad (47)$$

where $\gamma = \frac{nq}{2(q-n)}$.

The following L^∞ estimate of the tangential part is a straightforward adaptation of the proof of [24, Lemma 6]. Let us include it with the necessary modifications for completeness.

Lemma 4 *Let $q > n$ and $X: M \hookrightarrow \mathbb{R}^{n+1}$ be the immersion of a closed hypersurface. Then there exists a constant $C = C(n, q)$, such that*

$$\|X^T\|_\infty \leq C (V(M)\|B\|_q^n \|X\|_\infty)^{\frac{\gamma}{\gamma+1}} \|X^T\|_2^{\frac{1}{\gamma+1}}. \quad (48)$$

Proof: Set $\psi = |X^T|$. Due to

$$\psi^2 = |X|^2 - \langle X, \nu \rangle^2 \quad (49)$$

we obtain

$$|d\psi| \leq 1 + n\|B\| \|X\|_\infty. \quad (50)$$

Applying (26) with $f = \psi^{2\alpha}$, $\alpha \geq 1$, we obtain

$$\|\psi\|_{\frac{2\alpha n}{n-1}}^{2\alpha} \leq KV(M)^{\frac{1}{n}} (2\alpha(1+n\|B\|_q \|X\|_\infty) + \|H\|_q \|\psi\|_\infty) \|\psi\|_{\frac{(2\alpha-1)q}{q-1}}^{2\alpha-1}. \quad (51)$$

From here we follow the proof of [24, Lemma 5]. Setting

$$\mu = \frac{n(q-1)}{(n-1)q}, \quad a_0 = \frac{2q}{q-1}, \quad a_{p+1} = \mu a_p + \frac{n}{n-1}, \quad (52)$$

then we obtain, using

$$\alpha = a_p \frac{q-1}{2q} + \frac{1}{2}, \quad (53)$$

that

$$\begin{aligned} & \left(\frac{\|\psi\|_{a_{p+1}}}{\|\psi\|_\infty} \right)^{\frac{a_{p+1}}{\mu^{p+1}}} \\ & \leq \left(C(n)V(M)^{\frac{1}{n}} \left(a_{p+1} \frac{1+n\|B\|_q \|X\|_\infty}{\|\psi\|_\infty} + \|H\|_q \right) \right)^{\frac{n}{\mu^{p+1}(n-1)}} \left(\frac{\|\psi\|_{a_p}}{\|\psi\|_\infty} \right)^{\frac{a_p}{\mu^p}} \\ & \leq \left(C(n)V(M)^{\frac{1}{n}} a_{p+1} \frac{\|B\|_q \|X\|_\infty}{\|\psi\|_\infty} \right)^{\frac{n}{\mu^{p+1}(n-1)}} \left(\frac{\|\psi\|_{a_p}}{\|\psi\|_\infty} \right)^{\frac{a_p}{\mu^p}}, \end{aligned} \quad (54)$$

where we also used

$$1 \leq \|X\|_2 \|H\|_2 \leq C(n) \|X\|_\infty \|B\|_q, \quad \|\psi\|_\infty \leq \|X\|_\infty. \quad (55)$$

As in the proof of [24, Lemma 5] we obtain

$$\|\psi\|_\infty \leq C(q, n) \left(V(M)^{\frac{1}{n}} \frac{\|B\|_q \|X\|_\infty}{\|\psi\|_\infty} \right)^\gamma \|\psi\|_2, \quad (56)$$

where

$$\gamma = \frac{nq}{2(q-n)}. \quad (57)$$

Rearranging terms gives the desired estimate. \square

4.3 Proof of Theorem 2

In order to prove Theorem 2, we examine the consequences from the L^∞ -estimates under the pinching condition $(A_{p,\varepsilon})$. The proofs closely follow the argumentation in [24, Sec. 3].

Lemma 5 *Under the condition $(A_{p,\varepsilon})$ the immersion X satisfies*

$$1 - \frac{\|X - \bar{X}\|_1}{\|X - \bar{X}\|_2} \leq C(p)\varepsilon. \quad (58)$$

Proof: Assume by translation that $\bar{X} = 0$ and use Lemma 1 and the Hsiung-Minkowski formula to deduce

$$\begin{aligned} \|H_S\|_{2p}\|X\|_2 &\leq \frac{\int_M |\langle X, H_S \rangle|}{V(M)\sqrt{1-\varepsilon}} \\ &\leq \frac{1}{\sqrt{1-\varepsilon}} \|H_S\|_{2p} \|X\|_{\frac{2p}{2p-1}} \\ &\leq \frac{1}{\sqrt{1-\varepsilon}} \|H_S\|_{2p} \|X\|_1^{1-\frac{1}{p}} \|X\|_2^{\frac{1}{p}}. \end{aligned} \quad (59)$$

We obtain

$$\|X\|_2 \leq (1-\varepsilon)^{-\frac{p}{2(p-1)}} \|X\|_1 \quad (60)$$

and hence

$$1 - \frac{\|X\|_1}{\|X\|_2} \leq 1 - (1-\varepsilon)^{\frac{p}{2(p-1)}} \leq C(p)\varepsilon \quad (61)$$

with a constant $C = C(p)$. \square

Let us combine these results to get final pinching estimates for $|X|$ and $|X^T|$. First we prove (8).

Corollary 2 *Let $q > n$ and suppose $V(M)\|H\|_q^n \leq A$. Define*

$$r = \frac{\frac{1}{V(M)} \left| \int_M \operatorname{tr}(S) dv_g \right|}{\|H_S\|_{2p}} \quad (62)$$

and let S and T satisfy the conditions of Theorem 2. Then under the condition $(A_{p,\varepsilon})$ with $p > 1$ and $\varepsilon < \frac{1}{2}$ there exists a constant $C = C(n, p, q, A)$, such that

$$\| |X - \bar{X}| - r \|_\infty \leq Cr\varepsilon^{\frac{1}{2(\gamma+1)}}. \quad (63)$$

Proof: Again let $\bar{X} = 0$. Then we obtain

$$\| |X| - r \|_\infty \leq \| |X| - \|X\|_2 \|_\infty + \| \|X\|_2 - r \|. \quad (64)$$

The lemmata 1, 3 and 5 give the result. \square

Corollary 3 *Let $q > n$, $V(M)\|B\|_q^n \leq A$ and $\bar{X} = 0$. Let S and T satisfy the conditions of Theorem 2. Then under the condition $(A_{p,\varepsilon})$ with $p > 1$ and $\varepsilon < \frac{1}{2}$ there exists a constant $C = C(n, p, q, A)$, such that*

$$\|X^T\|_\infty \leq Cr\varepsilon^{\frac{1}{2(\gamma+1)}}. \quad (65)$$

Proof: Due to Lemma 2 and Lemma 4 we have

$$\|X^T\|_\infty \leq C\|X\|_\infty^{\frac{\gamma}{\gamma+1}}\|X^T\|_2^{\frac{1}{\gamma+1}} \leq C\|X\|_\infty^{\frac{\gamma}{\gamma+1}}\|X\|_2^{\frac{1}{\gamma+1}}\varepsilon^{\frac{1}{2(\gamma+1)}}. \quad (66)$$

Since

$$\|X\|_\infty \leq \| |X| - r \|_\infty + r \quad (67)$$

and

$$\|X\|_2 \leq Cr, \quad (68)$$

we obtain the result from Corollary 2. \square

Remark 3 Now the proof of the rest of Theorem 2 proceeds literally as the corresponding proof of [24, Thm. 2]. The starting points are the equations (63) and (65), which correspond to the equations (10) and (12) in [24]. Our radius r corresponds to the quantity $\frac{\|H_{k-1}\|_1}{\|H_k\|_p}$ in [24]. Note that this is where we have to restrict to some small $\varepsilon_0 > 0$ as claimed in Theorem 2.

5 Application to r -stability

In this section, we are interested in the stability of constant mean curvature hypersurfaces and, more generally, in r -stability. For this we introduce the r -area functionals

$$\mathcal{A}_r = \left(\int_M S_r dv_g \right), \quad r \in \{0, \dots, n-1\}. \quad (69)$$

Now we consider a variation of the immersion X_0 . Precisely, let $\delta > 0$ and

$$X: (-\delta, \delta) \times M \rightarrow \mathbb{R}^{n+1}, \quad (70)$$

such that for all $t \in (-\delta, \delta)$, $X_t := X(t, \cdot)$ is an immersion of M into \mathbb{R}^{n+1} and $X(0, \cdot) = X_0$. The precise assumptions on M will be specified in Theorem 4 below. We denote by $S_r(t)$ the corresponding curvature functions, by $\mathcal{A}_r(t)$ the r -area of X_t and finally we set

$$f_t = \left\langle \frac{dX}{dt}, \nu_t \right\rangle, \quad (71)$$

where ν_t is the unit normal to M induced by X_t and g_t is the induced metric on M .

Note that

$$\frac{d}{dt} \left(\int_M S_r(t) dv_{g_t} \right) = -(r+1) \int_M f S_{r+1} dv_{g_t}, \quad (72)$$

cf. [26, Thm. B]. We also consider the volume functional

$$V(t) = \int_{[0,t] \times M} X^* dv. \quad (73)$$

It is easy to see, cf. [9, Lemma 2.1], that V satisfies

$$V'(t) = \int_M f_t dv_{g_t} \quad (74)$$

and so X preserves the volume if and only if $\int_M f_t dv_{g_t} = 0$ for all t . Moreover, according to [9, Lemma 2.2], for any function $f_0 : M \rightarrow \mathbb{R}$ such that $\int_M f_0 dv_g = 0$, there exists a variation of X_0 preserving the volume and with normal part given by f_0 . Hence, in order to study variations with constant volume, we can equivalently consider normal parts with vanishing integral. This has the advantage that they can be used as test functions in the Rayleigh quotient for the first eigenvalue of the Laplacian or, more generally, for the operator L_r . As we will see, these operators play a crucial role in the study of r -stable hypersurfaces. Indeed, a standard argument as in [8, Prop. 2.7] shows that X_0 is a critical point for the functional \mathcal{A}_r for variations with constant volume if and only if H_{r+1} is constant. For such a critical point, Reilly [26] has computed the second variation of \mathcal{A}_r , also compare [3, equ. (2.2)]:

$$\begin{aligned} \mathcal{A}_r''(0) &= (r+1) \int_M f_0 L_r f_0 + \left[c(r+1) H_{r+2} - n \frac{c(r)}{r+1} H H_{r+1} \right] \int_M f_0^2 dv_g \\ &=: J_r f_0, \end{aligned} \quad (75)$$

where f_0 is the normal part of a variation with constant volume at $t = 0$ and

$$c(r) = (n-r) \binom{n}{r}. \quad (76)$$

We say that a hypersurface M with constant H_{r+1} is r -stable if $J_r f$ is non-negative for all functions f of vanishing integral. A classical result of Barbosa and do Carmo [8] for $r = 0$, Alencar, do Carmo and Colares [1] for $r = 1$ and Alencar, do Carmo and Rosenberg [3] for all r says that a compact hypersurface of the Euclidean space with constant H_{r+1} is r -stable if and only if it is a geodesic sphere. This result has been generalised later by Barbosa and Colares [7] for hypersurfaces of the hyperbolic space and the half-sphere.

Here, using the pinching result for the first eigenvalue of L_r proven above, we will show that this characterisation of spheres of the Euclidean space remains true if we only assume that $J_r f$ is almost positive, which means

$$J_r f \geq -\varepsilon \int_M f^2 H_{r+1}^{\frac{r+2}{r+1}} \quad \forall f \in C^\infty(M): \int_M f = 0, \quad (77)$$

where ε is a sufficiently small positive constant. Note that the term $H_{r+1}^{\frac{r+2}{r+1}}$ is present in order to have a condition which is invariant under any homothety of \mathbb{R}^{n+1} . Precisely, we prove the following result.

Theorem 4 *Let (M^n, g) be a connected, oriented and closed Riemannian manifold isometrically immersed into \mathbb{R}^{n+1} with constant $H_{r+1} > 0$. Let $A > 0$, $q > n$ and assume that $V(M)\|B\|_q^n \leq A$. Then there exists $\varepsilon_0 > 0$ depending on n , q and A such that if M is almost r -stable in the sense of (77) with $\varepsilon < \varepsilon_0$, then M is a sphere of radius $H_{r+1}^{-\frac{1}{r+1}}$.*

Proof: Taking f as an eigenfunction of L_r associated with the first positive eigenvalue, we get from the almost-stability that

$$\begin{aligned}
0 &\leq \int_M \left((r+1)\lambda_1 + (r+1)c(r+1)H_{r+2} - nc(r)H_1H_{r+1} + \varepsilon H_{r+1}^{\frac{r+2}{r+1}} \right) f^2 \\
&\leq \int_M \left((r+1)\lambda_1 + ((r+1)c(r+1) - nc(r))H_1H_{r+1} + \varepsilon H_{r+1}^{\frac{r+2}{r+1}} \right) f^2 \\
&\leq \int_M \left((r+1)\lambda_1 - ((r+1)c(r) - \varepsilon)H_{r+1}^{\frac{r+2}{r+1}} \right) f^2 \\
&= (r+1) \int_M \left(\lambda_1 - c(r) \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) H_{r+1}^{\frac{r+2}{r+1}} \right) f^2,
\end{aligned} \tag{78}$$

where we used (24), (25) and also

$$c(r+1) - n \frac{c(r)}{r+1} = -c(r). \tag{79}$$

Hence we find

$$\lambda_1 - c(r) \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) H_{r+1}^{\frac{r+2}{r+1}} \geq 0. \tag{80}$$

Up to a minor change this is the pinching condition of Theorem 2. Indeed, for $p > 1$ we have

$$\begin{aligned}
\lambda_1(L_r) &\geq \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) c(r) H_{r+1}^{\frac{r+2}{r+1}} \\
&= \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) c(r) \frac{H_{r+1}^2}{H_{r+1}^{\frac{r}{r+1}}} \\
&= \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) c(r) \frac{\|H_{r+1}\|_{2p}^2 V(M)}{\int_M H_{r+1}^{\frac{r}{r+1}}} \\
&\geq \left(1 - \frac{\varepsilon}{(r+1)c(r)} \right) c(r) \frac{\|H_{r+1}\|_{2p}^2 V(M)}{\int_M H_r},
\end{aligned} \tag{81}$$

where we used the fact that H_{r+1} is constant to make integrals appear and (24) for the last line. Now we have exactly the pinching condition for $\lambda_1(L_r)$ with $S = T = T_r$ and we can apply Theorem 2 to conclude that in particular M is embedded. In this argument also note that T_r is elliptic since H_{r+1} is positive. By the Alexandrov theorem for H_{r+1} proved by Ros, cf. [28], we get that M is a sphere. \square

6 An almost-Einstein type estimate

We finish this paper by applying the eigenvalue pinching result to deduce an explicit spherical closeness estimate of almost-Einstein type for hypersurfaces, namely an estimate of the form

$$d_{\mathcal{H}}(M, S(\bar{X}, r)) \leq \frac{cr}{\bar{R}^\alpha} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_p^\alpha, \quad (82)$$

whenever the right-hand side is small. We assume that the average of the scalar curvature R over M is positive, $\bar{R} > 0$, and that $R \geq 0$. We will prove this estimate for $p > \max(2, \frac{n}{2})$ and c and α will be suitable constants depending on n , p and an integral bound on H . In [24, Thm. 3] a similar application of the eigenvalue pinching was given, however here the authors derived the almost-isometry to the sphere with the help of the pinching

$$\|\text{Ric} - \lambda g\|_\infty < \varepsilon. \quad (83)$$

Here we want to improve this application, which is by the way also possible with the help of their eigenvalue pinching result: we want to relax the norm of the pinching quantity to L^p instead of L^∞ . Besides the improvement of the estimate itself this has the advantage that we also obtain an almost-Schur type estimate right away, due to the well-known almost-Schur estimate by De Lellis and Topping, cf. [13]. For manifolds of nonnegative Ricci curvature and $n \geq 3$ they provide the estimate

$$\int_M \left| \text{Ric} - \frac{\bar{R}}{n}g \right|^2 \leq \frac{n^2}{(n-2)^2} \int_M \left| \text{Ric} - \frac{R}{n}g \right|^2. \quad (84)$$

Similar estimates, relaxing the assumption on nonnegative Ricci curvature in different directions, were obtained by Cheng, cf. [11, Thm. 1.2] and by Ge and Wang in [15]. Instead of L^2 we need an L^p -estimate, but as was pointed out in [13], their proof easily adapts provided one has a Calderon-Zygmund type inequality. For the sake of completeness, we will provide the proof of this L^p -estimate. For a class of manifolds which allow for a Calderon-Zygmund estimate see the paper [22].

To achieve the relaxation from the L^∞ - to an L^p -pinching condition, we imitate the proof of a similar result by the second author concerning an almost-umbilical type estimate, compare the proof of [33, Thm. 4.1]. A similar result due to the first author under slightly different assumptions was achieved in [30].

Let us first recall De Lellis' and Topping's argument from [13] how to obtain the L^p -version of their almost-Schur lemma.

Lemma 6 *Let $n \geq 3$ and (M^n, g) be a closed Riemannian manifold, which admits an estimate of the form*

$$\|\nabla^2 u\|_q \leq C \|\Delta u\|_q, \quad q > 1, \quad (85)$$

for all smooth functions u , where C is a constant only depending on n , q and possibly some fixed geometric quantities of M . Then for $p \geq 2$ there holds

$$\left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_p \leq c \left\| \text{Ric} - \frac{R}{n}g \right\|_p, \quad (86)$$

where $c = c(C, n, p)$.

Proof: Compare [13, Sec. 2]. Let $f \in C^{2,p-2}(M)$ be the solution of

$$\begin{cases} \Delta f = |R - \bar{R}|^{p-2}(R - \bar{R}) \\ \int_M f = 0. \end{cases} \quad (87)$$

Then, with exactly the same computation as in [13, equ. (2.3)], we obtain

$$\begin{aligned} \frac{1}{V(M)} \int_M |R - \bar{R}|^p &= \frac{1}{V(M)} \int_M (R - \bar{R}) \Delta f \\ &\leq \frac{2n}{n-2} \|\mathring{\text{Ric}}\|_p \|\nabla^2 f\|_{\frac{p}{p-1}}, \end{aligned} \quad (88)$$

and hence, due to the Calderon-Zygmund estimate,

$$\frac{1}{V(M)} \int_M |R - \bar{R}|^p \leq c \|\mathring{\text{Ric}}\|_p \| |R - \bar{R}|^{p-1} \|_{\frac{p}{p-1}}. \quad (89)$$

This yields the desired estimate, also using

$$\left| \text{Ric} - \frac{\bar{R}}{n}g \right|^2 = |\mathring{\text{Ric}}|^2 + \frac{1}{n}(R - \bar{R})^2. \quad (90)$$

□

Let us come to the proof of (82). Here we will need to estimate the first Laplace eigenvalue in terms of an L^p -Ricci bound. We have the following estimate due to Aubry, originally proved in [5, Prop. 1.5], maybe more accessible in the version of [6, Thm. 1.6]. It says that for $p > n/2$, a complete Riemannian manifold (M^n, g) with

$$\frac{1}{V(M)} \int_M (\underline{\text{Ric}} - (n-1))_-^p < \frac{1}{C(p, n)} \quad (91)$$

is compact and satisfies

$$\lambda_1 \geq n \left(1 - C(n, p) \left(\frac{1}{V(M)} \int_M (\underline{\text{Ric}} - (n-1))_-^p \right)^{\frac{1}{p}} \right), \quad (92)$$

where $\underline{\text{Ric}}$ denotes the smallest eigenvalue of the Ricci tensor and $x_- = \max(0, -x)$.

Theorem 5 *Let $n \geq 2$, $q > n$, $p > \max(2, \frac{n}{2})$ and (M^n, g) be a closed, connected and oriented Riemannian manifold with $\bar{R} > 0$ and $R \geq 0$, isometrically immersed into \mathbb{R}^{n+1} . Suppose $V(M)\|H\|_q^n \leq A$. Then there exists a constant $\varepsilon_0(n, p, q, A) > 0$, such that whenever there holds*

$$\left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_p \leq \varepsilon_0 \bar{R}, \quad (93)$$

then we also have

$$d_{\mathcal{H}}(M, S(\bar{X}, r)) \leq \frac{cr}{\bar{R}^\alpha} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_p^\alpha, \quad (94)$$

where $\alpha = \alpha(n, q)$ and $c = c(n, p, q, A)$. If furthermore $V(M)\|B\|_q^n \leq A$, then M is embedded as a starshaped hypersurface and ε^α -almost-isometric to a sphere in the sense of Theorem 2.

Proof: We want to apply Theorem 2. We apply this theorem with the tensors $T = \text{Id}$ and $S = T_1$. Hence we have to provide the estimate

$$\lambda_1 \|H\|_1^2 \geq (1 - \varepsilon)n \|H_2\|_{2s}^2, \quad (95)$$

where λ_1 now is the first eigenvalue of $L_T = -\Delta$ and $s = \frac{p}{2}$. Using a simple rescaling argument we obtain a scaled version of Aubry's result [6, Thm. 1.6], namely that

$$\left(\frac{1}{V(M)} \int_M \left(\frac{\text{Ric} - \bar{R}}{n} \right)_-^{2s} \right)^{\frac{1}{2s}} < \frac{\bar{R}}{C(n, s)} \quad (96)$$

implies

$$\begin{aligned} \lambda_1 &\geq \frac{\bar{R}}{n-1} \left(1 - \frac{C(s, n)}{\bar{R}} \left(\frac{1}{V(M)} \int_M \left(\frac{\text{Ric} - \bar{R}}{n} \right)_-^{2s} \right)^{\frac{1}{2s}} \right) \\ &\geq \frac{\bar{R}}{n-1} - C(s, n) \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s}. \end{aligned} \quad (97)$$

Let us estimate $\|H\|_1^2$.

$$\begin{aligned}
\|H\|_1^2 &\geq \left(\frac{1}{V(M)} \int_M H_2^{\frac{1}{2}} \right)^2 \\
&= \frac{1}{n(n-1)} \left(\frac{1}{V(M)} \int_M R^{\frac{1}{2}} \right)^2 \\
&= \frac{\bar{R}}{n(n-1)} + \frac{1}{n(n-1)} \left(\|R^{\frac{1}{2}}\|_1^2 - \|\bar{R}^{\frac{1}{2}}\|_1^2 \right) \\
&\geq \frac{\bar{R}}{n(n-1)} - \frac{1}{n(n-1)} (\|R^{\frac{1}{2}}\|_1 + \bar{R}^{\frac{1}{2}}) \|R^{\frac{1}{2}} - \bar{R}^{\frac{1}{2}}\|_1 \\
&\geq \frac{\bar{R}}{n(n-1)} - \frac{1}{n(n-1)\sqrt{\bar{R}}} \|R - \bar{R}\|_1 (\|R^{\frac{1}{2}}\|_1 + \bar{R}^{\frac{1}{2}}) \\
&\geq \frac{\bar{R}}{n(n-1)} - c_n \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_1.
\end{aligned} \tag{98}$$

We also have to estimate $\|H_2\|_{2s}^2$.

$$\begin{aligned}
\|H_2\|_{2s}^2 &= \frac{1}{n^2(n-1)^2} \left(\frac{1}{V(M)} \int_M R^{2s} \right)^{\frac{1}{s}} \\
&\leq \frac{\bar{R}^2}{n^2(n-1)^2} + \frac{1}{n^2(n-1)^2} \| \|R\|_{2s}^2 - \|\bar{R}\|_{2s}^2 \| \\
&\leq \frac{\bar{R}^2}{n^2(n-1)^2} + \frac{\|R\|_{2s} + \bar{R}}{n^2(n-1)^2} \|R - \bar{R}\|_{2s} \\
&\leq \frac{\bar{R}^2}{n^2(n-1)^2} + c_n (\|R\|_{2s} + \bar{R}) \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s}.
\end{aligned} \tag{99}$$

Bringing together (97), (98) and (99), we obtain, also noting that the right-hand side of (97) is still positive when $\varepsilon_0 > 0$ is small,

$$\begin{aligned}
\lambda_1 \|H\|_1^2 &\geq \frac{\bar{R}^2}{n(n-1)^2} - c_n \bar{R} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_1 - c(n, s) \bar{R} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s} \\
&\geq n \|H_2\|_{2s}^2 - c (\|R\|_{2s} + \bar{R}) \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s} \\
&= \left(1 - c \frac{\|R\|_{2s} + \bar{R}}{\|R\|_{2s}^2} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s} \right) n \|H_2\|_{2s}^2 \\
&\geq \left(1 - \frac{c}{\bar{R}} \left\| \text{Ric} - \frac{\bar{R}}{n}g \right\|_{2s} \right) n \|H_2\|_{2s}^2.
\end{aligned} \tag{100}$$

Thus when ε_0 is chosen small enough, we may apply Theorem 2 to conclude the result. \square

Remark 4 Let us note again, as already mentioned in the introduction to this section, that Theorem 5 enables us to obtain an almost-Schur type estimate, whenever we obtain an estimate in the sense of De Lellis and Topping [13].

References

1. Alencar, H., Do Carmo, M., Colares, A.: Stable hypersurfaces with constant scalar curvature. *Math. Z.* **213**(1), 117–131 (1993)
2. Alencar, H., Do Carmo, M., Marques, F.C.: Upper bounds for the first eigenvalue of the operator l_r and some applications. *Illinois J. Math.* **45**(3), 851–863 (2001)
3. Alencar, H., Do Carmo, M., Rosenberg, H.: On the first eigenvalue of the linearized operator of the r -th mean curvature of a hypersurface. *Ann. Glob. Anal. Geom.* **11**(4), 387–395 (1993)
4. Alias, L., Malacarne, M.: On the first eigenvalue of the linearized operator of the higher order mean curvature for closed hypersurfaces in space forms. *Illinois J. Math.* **48**(1), 219–240 (2004)
5. Aubry, E.: Finiteness of π_1 and geometric inequalities in almost positive Ricci curvature. *Ann. Sci. Éc. Norm. Supér. (4)* **40**(4), 675–695 (2007)
6. Aubry, E.: Diameter pinching in almost positive Ricci curvature. *Comment. Math. Helv.* **84**(2), 223–233 (2009)
7. Barbosa, J., Colares, A.: Stability of hypersurfaces with constant r -mean curvature. *Ann. Glob. Anal. Geom.* **15**(3), 277–297 (1997)
8. Barbosa, J., Do Carmo, M.: Stability of hypersurfaces with constant mean curvature. *Math. Z.* **185**(3), 339–353 (1984)
9. Barbosa, J., Do Carmo, M., Eschenburg, J.H.: Stability of hypersurfaces of constant mean curvature in Riemannian manifolds. *Math. Z.* **197**(1), 123–138 (1988)
10. Bleecker, D., Weiner, J.: Extrinsic bound on λ_1 of δ on a compact manifold. *Comment. Math. Helv.* **51**, 601–609 (1976)
11. Cheng, X.: A generalization of almost-Schur lemma for closed Riemannian manifolds. *Ann. Glob. Anal. Geom.* **43**, 153–160 (2013)
12. Colbois, B., Grosjean, J.F.c.: A pinching theorem for the first eigenvalue of the Laplacian on hypersurfaces of the Euclidean space. *Comment. Math. Helv.* **82**(1), 175–195 (2007)
13. De Lellis, C., Topping, P.: Almost-Schur lemma. *Calc. Var. Partial Differ. Equ.* **43**(3–4), 347–354 (2012)
14. El Soufi, A., Ilias, S.: Une inégalité du type "Reilly" pour les sous-variétés de l'espace hyperbolique. *Comment. Math. Helv.* **67**, 167–181 (1992)
15. Ge, Y., Wang, G.: An almost Schur theorem on 4-dimensional manifolds. *Proc. Amer. Math. Soc.* **140**(3), 1041–1044 (2012)
16. Gerhardt, C.: Curvature problems, *Series in Geometry and Topology*, vol. 39. International Press of Boston Inc. (2006)
17. Grosjean, J.F.c.: Upper bounds for the first eigenvalue of the Laplacian on compact submanifolds. *Pac. J. Math.* **206**(1), 93–112 (2002)
18. Grosjean, J.F.c.: Extrinsic upper bounds for the first eigenvalue of elliptic operators. *Hokkaido Math. J.* **33**(2), 319–339 (2004)
19. Grosjean, J.F.c., Roth, J.: Eigenvalue pinching and application to the stability and the almost umbilicity of hypersurfaces. *Math. Z.* **271**(1–2), 469–488 (2012)
20. Hardy, G., Littlewood, J., Polya, G.: *Inequalities*. Cambridge University Press (1952)
21. Heintze, E.: Extrinsic upper bounds for λ_1 . *Math. Ann.* **280**(3), 389–402 (1988)
22. Hiroshima, T.: Construction of the Green function on Riemannian manifold using harmonic coordinates. *J. Math. Tokyo Univ.* **36**(1), 1–30 (1996)
23. Hoffmann, D., Spruck, J.: Sobolev and isoperimetric inequalities for Riemannian submanifolds. *Comm. Pure Appl. Math.* **27**(6), 715–727 (1974)
24. Hu, Y., Xu, H., Zhao, E.: First eigenvalue pinching for Euclidean hypersurfaces via k -th mean curvatures. *Ann. Glob. Anal. Geom.* **48**(1), 23–35 (2015)
25. Michael, J., Simon, L.: Sobolev and mean-value inequalities on generalized submanifolds of \mathbb{R}^n . *Comm. Pure Appl. Math.* **26**(3), 361–379 (1973)
26. Reilly, R.: Variational properties of functions of the mean curvatures for hypersurfaces in space forms. *J. Differ. Geom.* **8**(3), 465–477 (1973)
27. Reilly, R.: On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space. *Comment. Math. Helv.* **52**(4), 525–533 (1977)
28. Ros, A.: Compact hypersurfaces with constant higher order mean curvatures. *Rev. Mat. Iberoam.* **3**(3–4), 447–453 (1987)

-
29. Roth, J.: Pinching of the first eigenvalue of the Laplacian and almost-Einstein hypersurfaces of the Euclidean space. *Ann. Glob. Anal. Geom.* **33**(3), 293–306 (2008)
 30. Roth, J.: A remark on almost umbilical hypersurfaces. *Arch. Math.* **49**(1), 1–7 (2013)
 31. Roth, J.: Upper bounds for the first eigenvalue of the Laplacian of hypersurfaces in terms of anisotropic mean curvatures. *Result. Math.* **64**(3-4), 383–403 (2013)
 32. Roth, J.: General Reilly-type inequalities for submanifolds of weighted Euclidean spaces. *Colloq. Math.* **144**(1), 127–136 (2016)
 33. Scheuer, J.: Quantitative oscillation estimates for almost-umbilical closed hypersurfaces in Euclidean space. *Bull. Aust. Math. Soc.* **92**(1), 133–144 (2015)
 34. Takahashi, T.: Minimal immersions of Riemannian manifolds. *J. Math. Soc. Japan* **18**(4), 380–385 (1966)