

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/140621/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Casale, Marisa, Cluver, Lucie, Boyes, Mark, Toska, Elona, Gulaid, Laurie, Armstrong, Alice, Shenderovich, Yulia, Rudgard, William E., Zhou, Siyanai and Langwenya, Nontokozo 2021. Bullying and ART nonadherence among South African ALHIV: effects, risks, and protective factors. *Journal of Acquired Immune Deficiency Syndromes* 86 (4), pp. 436-444. 10.1097/QAI.0000000000002574

Publishers page: <http://doi.org/10.1097/QAI.0000000000002574>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Accepted manuscript

Bullying and ART non-adherence among South African adolescents living with HIV: effects, risk and protective factors

Running head: Bullying and non-adherence among ALHIV

Marisa Casale PhD^{a, b, §}, Lucie Cluver PhD^{a, c}, Mark Boyes PhD^d, Elona Toska PhD^{a, e, f}, Laurie Gulaid PhD^g, Alice Armstrong PhD^g, Yulia Shenderovich PhD^a, William E. Rudgard PhD^a, Siyanai Zhou MPhil^e, Nontokozo Langwenya MPH^{e, h}

^aDepartment of Social Policy and Intervention, University of Oxford, Oxford, United Kingdom

^bSchool of Public Health, University of the Western Cape, Cape Town, South Africa

^cDepartment of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

^dSchool of Psychology, Faculty of Health Sciences, Curtin University, Australia

^eCentre for Social Science Research, University of Cape Town, Cape Town, South Africa.

^fDepartment of Sociology, University of Cape Town, Cape Town, South Africa

^gUNICEF Eastern and Southern Africa Regional Office

^hOxford Research South Africa, East London, South Africa

§Corresponding author: Marisa Casale, School of Public Health, University of the Western Cape, Robert Sobukwe Road, Bellville 7535, South Africa; phone: +27 (82) 9090276; email: maj.casale@icloud.com; mcasale@uwc.ac.za.

Conflicts of Interest and Source of Funding:

The Mzantsi Wakho study was jointly funded the Nuffield Foundation [CPF/41513], however the views expressed are those of the authors and not necessarily those of the Foundation; the Evidence for HIV Prevention in Southern Africa (EHPSA), a UK aid programme managed by Mott MacDonald; Janssen Pharmaceutica N.V., part of the

Janssen Pharmaceutical Companies of Johnson & Johnson; the Regional Inter-Agency Task Team for Children Affected by AIDS - Eastern and Southern Africa (RIATT-ESA); UNICEF Eastern and Southern Africa Office (UNICEF-ESARO); the International AIDS Society through the CIPHER grant [155-Hod; 2018/625-TOS], the views expressed in written materials or publications do not necessarily reflect the official policies of the International AIDS society. Further funding provided by: Oxford University Clarendon-Green Templeton College Scholarship (ET); Claude Leon Foundation [F08 559/C]; Oak Foundation [R46194/AA001]; the Leverhulme Trust [PLP-2014-095].

Additional funding was provided for the research team by: the John Fell Fund [103/757; 161/033]; the University of Oxford's ESRC Impact Acceleration Account [IAA-MT13-003; 1602-KEA-189; K1311-KEA-004]; the UKRI GCRF Accelerating Achievement for Africa's Adolescents (Accelerate) Hub (Grant Ref: ES/S008101/1); the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement, and by the Department of Health Social Care (DHSC) through its National Institutes of Health Research (NIHR) [MR/R022372/1]; the Fogarty International Center, National Institute on Mental Health, National Institutes of Health under Award Number K43TW011434, the content is solely the responsibility of the authors and does not represent the official views of the National Institutes of Health; the European Research Council (ERC) under the European Union's Seventh Framework Programme [FP7/2007-2013]/ ERC grant agreement n°313421 and the European Union's Horizon 2020 research and innovation programme (grant agreement n°737476 and n°771468).

The authors have no conflicts of interest to disclose.

Author contributions

MC: led conceptualization, analysis and writing. LC: conceptualization, analysis and writing. MB: conceptualization, analysis and revision. ET: conceptualization, data management, analysis and revision. LG, AA: conceptualization and revision. YS, WR, SZ and NL: data collection and management, analysis and revision.

Abstract (240 words)

Background: Identifying risk and protective factors for adolescent antiretroviral therapy (ART) adherence is a public health priority, given high HIV-related mortality in this population. An area that merits further investigation is the relationship between bullying victimization, mental health problems, and ART non-adherence among adolescents living with HIV (ALHIV). However, no known studies assess effects of bullying on adolescent non-adherence, or risk and protective factors that could moderate this relationship.

Setting: This study investigates a) the direct longitudinal relationship between bullying exposure and ART non-adherence, and the indirect relationship via psychological distress, and b) potential risk and modifiable protective factors moderating these pathways, among vertically and horizontally infected ALHIV who initiated treatment across 53 public healthcare facilities in a South African health district.

Methods: Survey data was collected at two-time points, between 2014 and 2017, with 1046 ALHIV (94% retention). Various mediation and moderated mediation models were run as part of a staged analysis approach.

Results: A significant longitudinal relationship was found between bullying victimization and non-adherence, operating indirectly via psychological distress ($B=0.07$; 95%CI: [0.03, 0.13]). Moderation analyses indicated that older adolescents exposed to bullying are more at risk of non-adherence ($B=0.52$; 95%CI: [0.07, 0.97] $p<0.05$), and parental monitoring is a potential protective factor buffering indirect effects of bullying on non-adherence ($B=-0.22$; 95%CI: [-0.42, -0.02] $p<0.05$).

Conclusions: These findings underscore the importance of interventions that address bullying and psychological distress, and strengthen parental monitoring, particularly among older ALHIV.

Keywords: bullying victimization, ART non-adherence, adolescents, mental health, parental monitoring

INTRODUCTION

Adolescents and youth represent a growing proportion of people living with HIV worldwide ¹⁻². They are also a high-risk population for HIV treatment non-adherence and AIDS-related mortality ³⁻⁵. This is of particular public health concern in low- and middle- income countries (LMICs) facing high HIV prevalence rates among youth and challenges in the provision of adequate health care services. Identifying risk and protective factors for adolescent ART adherence is key to developing more effective interventions and reaching the most vulnerable adolescents ⁶.

An area that merits further investigation is the relationship between bullying victimization, mental health problems, and ART non-adherence. Bullying remains a widespread phenomenon among adolescents worldwide, estimated to affect over 30% of school-going youth globally, and around 50% in Eastern and Southern Africa ^{7,8}. It has been associated with various negative health outcomes that can manifest during adolescence and extend into adulthood. These include sleep disorders, psychosomatic complaints, poor self-reported physical health, and more difficult social relations ⁹⁻¹¹. However, evidence of causal effects of bullying victimization is strongest for mental health problems, such as depression, anxiety, and suicidal ideation ¹²⁻¹⁴.

Understanding the adverse effects of bullying, and how to prevent or mitigate these, may be particularly salient for adolescents living with HIV (ALHIV). First, these adolescents are already at risk of exposure to HIV-related stigma ^{15,16} and may be at elevated risk of mental health disorders compared to the general population and other high-risk groups ¹⁷⁻²⁰. Research with ALHIV in Southern Africa, for example, has

shown bullying victimization to be associated with worse mental health ^{14,21,22}. Second, poor mental health has been associated with ART non-adherence among ALHIV ²³⁻²⁵. However, there are no known studies assessing direct or indirect effects of bullying on adolescent ART non-adherence. Moreover, there is no evidence of risk or protective factors that could moderate these effects.

This study seeks to address the following three research questions: i) Are ALHIV who experience bullying more likely to be non-adherent to ARV treatment over time and is this relationship mediated by greater psychological distress? ii) Is the risk of worse adherence greater for specific subgroups of ALHIV who experience bullying? iii) Are there modifiable protective factors at the clinic, school, or family level that can buffer the potential effects of being bullied on worse mental health or ART non-adherence?

In order to address the question of particularly vulnerable subgroups of adolescents (ii), age, gender, mode of HIV transmission and urban versus rural location are tested as moderators; previous analyses with South African ALHIV have shown (older) age, (female) gender, living in an urban versus rural location, and horizontal infection to be associated with worse mental health ^{22,26}. To address the question of potentially modifiable protective factors (iii) we assess moderation of factors within adolescents' family, clinic, and school environments, which have been shown to play a protective role for better adolescent mental health or treatment outcomes more broadly. These environments are key structures within adolescents' microsystems and can shape their beliefs and health behaviors ²⁷.

Family-, clinic- and school- level potential protective factors

At the family level, we test for moderating effects of positive parenting and parental monitoring. Both have respectively been associated with a lower likelihood of mental health problems among children and adolescents ²⁸⁻³¹, although not specifically with ALHIV. Parental monitoring has also been found to buffer effects of bullying on suicidal behavior ³² and antisocial behavior ^{28,33}. Supportive parenting styles have been linked to better treatment adherence ^{34,35}, while a lack of parental monitoring has been associated with worse treatment adherence ^{36,37} among youth living with (other) chronic illness.

At the clinic level, we test for moderating effects of support groups and treatment buddies. Support group interventions have been found to improve mental health and psychosocial functioning in ALHIV and HIV-affected youth ^{38,22}. Evaluation research from Southern Africa indicates that HIV support groups, within or linked to public health facilities, may have a positive impact on adherence and virological outcomes among children, adolescents, and adults ^{39,40, 41,42}. Treatment buddies are individuals in the patient's network who provide support with ART adherence; support may include transport to health facilities, observing ingestion of medication, assisting with management of adverse effects, reminding participants of drug pickup, and providing encouragement. The majority of studies conducted in Southern Africa show treatment buddies to be positively associated with patient outcomes such as increased ART adherence, better clinic attendance ^{43,44}, better medium-term survival rates, and virological and immunological responses ^{40,45}.

Lastly, at the school level, we test for moderating effects of school enrolment. Although there is limited evidence on the role of school-level variables as protective factors for mental health and treatment adherence in LMICs, associations have been found between school absenteeism or dropout and poor mental health in Lebanon, North America, Europe, and Asia ^{46,47,48}.

METHODS

Data collection

The longitudinal [*identifying information omitted*] study was conducted with 1046 ALHIV (10-19) in one of South Africa's highest HIV-prevalence health districts in the Eastern Cape Province. The province is characterized by high morbidity, low human development and poor infrastructure. Led by the University of [*identifying information omitted*] and University of [*identifying information omitted*], this study was a collaboration with the South African Departments of Health and Basic Education, UNICEF, PEPFAR-USAID, and regional and local NGOs. Ethical clearance was obtained from [*identifying information of universities omitted*], and provincial South African Departments of Health, Basic Education and Social Development.

Adolescents were interviewed at two-time points between 2014 and 2017, approximately 12-18 months apart. This paper analyses data pertaining to the 1046 adolescents living with HIV, who had initiated ARV treatment at one of the district's public healthcare facilities (n=53) at baseline. Both vertically and horizontally infected adolescents were included in this sample. Eligible ALHIV were identified through

clinic patient records and traced back to their communities, to ensure inclusion of adolescents no longer engaged in care. Voluntary informed consent was obtained from all adolescents and their primary caregivers at both time points. Questionnaires were developed with input from adolescent advisory groups, pre-piloted and administered in the adolescents' language of choice, including Xhosa and English. The tablet-assisted 60–90-minute individual interviews were conducted in clinics or communities by researchers trained in working with vulnerable adolescents. Confidentiality was maintained, except where participants disclosed serious risk of harm to themselves or others, in which case safeguarding procedures were followed, in conjunction with the relevant government services. The rate of baseline uptake was 90% (1046) and 94% of these adolescents (979) were retained at follow up (with 2.4% mortality).

Outcome variable: non-adherence at follow up

Self-reported ART adherence over the past week was measured at baseline and follow-up via items adapted from the standardized Patient Medication Adherence Questionnaire ⁴⁹, combined with measures developed in Botswana ⁵⁰. This adherence measure was validated against undetectable viral load (<50 copies/ml) for adolescent subsamples for which patient file data was available, controlling for key socio-demographic factors and health status (see Table S1). In this analysis a binary variable was used to denote full past-week self-reported adherence (0) or non-adherence (1).

Independent variable: Exposure to bullying at baseline

Bullying victimization was measured using the 9-item 'Social and Health Assessment Peer Victimization Scale' ⁵¹ previously used in research with vulnerable children in

South Africa ⁵² ($\alpha = 0.78$ for this sample). Items include: being called names, being hit or threatened, and having possessions broken or stolen. Response options range from 1/ 'not at all' to 4/ '4 or more times.' For this analysis, the scale variable was dichotomized so that any reported experience of bullying was recorded as '1', and no experience of bullying as '0' (i.e. when the adolescent answered 1/ 'not at all' to all items).

Mediator: psychological distress at follow up

A composite psychological distress variable was computed, by adding the standardized scores of the respective depression and anxiety scales at follow up. Depression was measured with the Child Depression Inventory – Short Form ⁵³ while anxiety was measured using an abbreviated version of the Revised Children's Manifest Anxiety Scale ⁵⁴, previously validated in South Africa with AIDS-affected children ⁵⁵. Internal consistency was good ($\alpha = 0.8$) for the combined scale in this sample.

Moderators: family-, clinic- and school-related

Both positive parenting and parental monitoring or supervision were measured using the short form of the Alabama Parenting Questionnaire ⁵⁶. Positive parenting included 7 items on praise, positive reinforcement, and support from the caregiver ($\alpha = 0.9$ at baseline and 0.92 at follow up). Good parental supervision included 10 items focused on monitoring of adolescent social activities and home rule-setting ($\alpha = 0.93$ at baseline and 0.89 at follow up). Response options were on a 5-point scale ranging from 0/ 'never' to 4/ 'always', based on the child-report of frequency of parental behaviors. For greater

consistency with the other moderators tested and ease of interpretation, these scale variables were dichotomized into ‘high’ (1) and low (0) levels of positive parenting and parental monitoring respectively. A code of ‘1’ was allocated if adolescents answered ‘always’ or ‘often’ to all positive parenting or good parental monitoring items for the respective scales. Binary variables were also used to denote clinic support group attendance, as in previous analyses¹⁷, having a treatment buddy and being enrolled in a school.

Socio-demographic and HIV-related moderators and covariates

All analyses conducted in this paper controlled for six binary covariates measured at baseline, where these were not included as moderators. These were recorded using adolescent self-report, in some cases combined with clinical records: (1) age (dichotomized to denote younger adolescents aged 10 to 14 and older adolescents aged 15 to 19); (2) male or female gender; (3) rural versus urban residential location; (4) vertical versus horizontal HIV transmission, assessed following a literature-informed algorithm used for existing Sub-Saharan African pediatric HIV cohorts⁵⁷; (5) length of time on ART, measured via self-report and clinic records to indicate more versus less than one year on treatment; (6) informal (versus formal) housing, as a proxy measure of socio-economic-status.

Statistical Analysis

Statistical analyses were conducted using SPSS 25 and the PROCESS macro⁵⁸. Descriptive statistics and frequencies were run for all variables used in the analysis, to

provide an overview of participant characteristics for the sample. The main analysis was conducted in 4 sequential stages. First, PROCESS model 4 was run to test for a direct relationship between being bullied at baseline and non-adherent at follow up, and/or an indirect relationship mediated by greater psychological distress, controlling for baseline non-adherence. PROCESS models use bootstrapping, a non-parametric sampling procedure, to simultaneously test for direct and indirect effects of the mediator⁵⁹; for 5000 bootstrap samples, results are statistically significant where 95% confidence intervals do not overlap zero.

The second stage of the analysis aimed to address the question of whether specific subgroups of adolescents were at greater risk of bullying leading to poor adherence. It tested for moderation of the direct and indirect pathways between bullying and non-adherence, by running a moderated mediation model in PROCESS (model 59) for each of four potential socio-demographic or HIV-related moderators of interest: age, gender, mode of HIV transmission and urban versus rural location.

The third stage consisted of testing for moderation of the five clinic, school, and parenting-related potential protective factors, to determine whether any of these could buffer direct or indirect effects of bullying on adherence. Each potential moderator was tested in a separate moderated mediation model, controlling for baseline non-adherence; baseline values of the moderator variables were tested for the bullied - psychological distress association and the (direct) bullied - non-adherence association, while follow-up values of the moderator variables were tested for the psychological distress - non-adherence association, using PROCESS model 22.

The fourth stage of the analysis consisted of combining all previously tested moderating variables with interaction terms significant at the $p < 0.05$ level into one final moderated mediation model. In the final model, moderators were tested for the specific association(s) for which they had shown significance during stages 2 and 3 of the analysis. Significant interactions were then probed and illustrated⁶⁰ to further explore moderation effects.

RESULTS

At baseline, the majority of ALHIV were between 10 and 14 years of age (63%), female (55%), and based in an urban area (73%). 78.5% were vertically infected with HIV and 82.5% had been on treatment for a year or more. The rates of non-adherence were 34.2% at baseline and 35.5% at follow-up. While more than half the sample reported having a treatment buddy at both data collection points, less than 15% were attending a clinic support group, and the large majority were enrolled in school (89% at follow up). Approximately half the sample reported high positive parenting at both time points, while just over 40% reported high parental monitoring. Additional descriptive statistics for this sample, disaggregated by mode of HIV transmission and gender respectively, are also provided in the supplementary material (see Table S2).

[Table 1: Approx. here]

Results of mediation analysis: effects of bullying on ART non-adherence

As indicated in Table 2 and illustrated in Figure S1, mediation analysis showed no direct association between being bullied at baseline and non-adherence at follow up. However, findings indicated an indirect effect mediated through greater psychological distress ($B= 0.07$; 95%CI: [0.03, 0.13]). Both the association between having experienced bullying at baseline and more psychological distress at follow up ($B = 0.37$, 95%CI: [0.16, 0.59] $p<0.001$), and the association between more psychological distress and higher odds of non-adherence at follow up ($B=0.20$; 95%CI: [0.11, 0.28] $p<0.001$) were significant.

[Table 2: Approx. here]

Results of moderated mediation analyses: testing for risk and protective factors

Each of the three associations constituting the direct and indirect relationship between bullying and non-adherence⁶¹ were evaluated for moderation by socio-demographic or HIV-related factors (age, gender, mode of transmission, and rural versus urban location). Two interaction terms were significant for the psychological distress outcome: older age x bullied ($B=0.57$, $p<0.05$) and female x bullied ($B=0.43$, $p< 0.05$). The direction of the interaction term coefficients suggested that older (15-19) adolescents exposed to bullying victimization were at higher risk than younger adolescents, and that female adolescents exposed to bullying were more at risk than male adolescents of having greater psychological distress one year later. Instead the non-significant interaction terms for mode of transmission and participant location indicated that these factors were not moderating the direct or indirect relationship between bullying victimization and non-adherence.

Moderated mediation models were then run to test both the direct and indirect pathways for moderation of each of the clinic-, school- and family-related variables. Only the better parental monitoring x psychological distress at follow up ($B=-0.23$, $p<0.05$) interaction term was significant. The negative coefficient indicated that this was in the expected direction, suggesting that parental monitoring could be buffering the risk of high psychological distress leading to non-adherence among these adolescents.

Based on the results of the first three stages of this analysis, described above, a final moderated mediation model was developed and tested, using PROCESS model 23. The three moderators for which interaction terms were significant at the 0.05 level in the individual models were included in this final model: 1) older age for the association between being bullied at baseline and higher psychological distress at follow up; 2) gender (female) for the association between being bullied at baseline and higher psychological distress at follow up; 3) high parental monitoring at follow up for the association between higher psychological distress and non-adherence at follow up.

Results of this final moderated mediation model are illustrated in Table 3 and Figure 1 below. They showed no significant direct effect between being bullied at baseline and non-adherent at follow up, as expected. When all three potential moderators were included in the same model, the interaction for female gender was not significant but the interaction terms for age ($B=0.52$; 95% CI: [0.07, 0.97] $p=0.024$) and high parental monitoring ($B=-0.22$; 95% CI: [-0.42, -0.02] $p=0.029$) maintained significance. These significant interaction terms suggest that age and parental monitoring respectively moderate the indirect relationship between bullying victimization and HIV treatment

non-adherence. Specifically, older adolescents and adolescents with low parental monitoring appear to be more at risk of not adhering to treatment if they experience bullying. Interaction plots support these moderation findings (see Figures S2 and S3). There was no significant direct association between high parental monitoring and non-adherence.

[Table 3: Approx. here]

[Figure 1: Approx. here]

Table 4 below indicates whether the indirect relationship between being bullied at baseline and non-adherent at follow up holds at different combinations of the three moderators. For adolescents reporting high parental monitoring at follow up, there is no evidence of a significant indirect relationship between being bullied at baseline and worse non-adherence at follow up (regardless of age or gender). However, for older adolescents who report low parental monitoring this indirect relationship is significant (for older males: Effect = 0.13; 95%CI: [0.02, 0.28] and for older females: Effect= 0.22; 95%CI: [0.1, 0.39]).

[Table 4: Approx. here]

DISCUSSION

Results of this analysis indicate that adolescents living with HIV who experience bullying victimization are at greater risk of non-adherence to HIV treatment one year later, and that this relationship operates indirectly via psychological distress. This

highlights the importance of preventing, identifying, and addressing exposure to bullying and psychological distress. Findings also indicate that older adolescents who are bullied are more at risk for psychological distress and treatment non-adherence. Particular attention should be paid to this group of older (15-19) adolescents, also considering that trends in HIV-associated deaths in this group are not following the decline observed among younger (10-14) ALHIV⁴.

Evaluations of school-based anti-bullying interventions in high-income countries suggest that complex multi-component interventions are most successful, particularly where they include families of students, and lead to increased adult supervision in school and community locations where aggression is likely to occur ^{62,63}. These approaches recognize that bullying is a complex phenomenon that involves not only victims, but also other actors in an adolescent's life, including bullies, teachers and caregivers ⁶³. Despite promising findings of recent school-based behavioral interventions focusing on teachers and peers ^{64,65}, to date evaluations of anti-bullying interventions in LMICs have been scarce and the evidence not overall conclusive ⁶⁶. There is clearly a need to invest in developing and testing further interventions in LMICS in Southern Africa and beyond.

This study also exposed parental monitoring as a potential protective factor against the effects of bullying and psychological distress on treatment non-adherence among ALHIV. Positive parenting and peer support, through support groups and treatment buddies, instead did not emerge as similarly protective in this study. However, these factors have been found to have direct positive effects on mental health and ART adherence in various populations and settings ^{22,39,67,68} and should be considered as

protective resources for health and retention in care more broadly. For this study, caregiver rule-setting and monitoring of adolescents' social activities and treatment were found to be important for treatment adherence among ALHIV already experiencing bullying and psychological distress. This finding reinforces the importance of developing and expanding access to family-based interventions, such as those that work with both adolescents and caregivers to strengthen parenting skills and improve caregiver-adolescent interaction⁶⁹⁻⁷¹. It also highlights the need to involve caregivers, where possible, in interventions not primarily designed as 'family interventions'; these include school-based initiatives aimed at addressing violence or boosting positive health behaviors^{62,63}.

Moreover, there is recent evidence of the acceptability and efficacy of certain family- and community- based interventions in improving adolescent mental health in LMICs, particularly those that incorporate cognitive-behavioral approaches^{70,72-74}. It would be useful to further explore the effectiveness of community-based interventions with ALHIVs that seek to both strengthen caregiver-adolescent interaction and address adolescent psychological distress. Another key area for intervention is increasing the awareness of bullying and mental health among school and clinic staff, community health workers, and ART treatment supporters. It may be possible to use or adapt existing tools, such as the World Health Organization mhGAP training or the HEADSSS tool, to help these providers better identify and support adolescents experiencing bullying and distress⁷⁵⁻⁷⁷.

This study has a number of limitations. The adherence variable, and other indicators used in this study, are based primarily on self-report data and therefore may reflect

some level of bias in perception and recall of events. However, adolescent self-report was considered the most reliable option for non-adherence in this context, given limited availability and reliability of patient medical data in this context of a very overburdened health system. Moreover, a short-term measure of non-adherence was chosen to minimize recall bias and tested against viral load indicators for the subsample for which clinic data was available (as described above). Also, despite the strong empirical and theoretical bases for our hypotheses, causality cannot be inferred; it is possible, for example, that consequences of non-adherence could affect psychological distress, or that the relationship between bullying victimization and psychological distress could be bidirectional¹³. Testing for mediation would ideally require data at 3 time-points. We note the limitations and bias related to conducting mediation analysis with two waves of data, where the mediator variable (psychological distress) and the outcome variable (non-adherence) were measured at the same time point. In addition, our choice of potential protective factors to test as moderators was limited by the variables included in the dataset; for example, we were not able to test peer support⁸ or strong school performance⁷⁸. It was also not possible to determine the extent to which reported bullying victimization was linked to participants' HIV status.

Despite these limitations, this remains the first longitudinal analysis to investigate the direct and indirect relationship between bullying victimization and non-adherence among ALHIV, and to identify factors potentially moderating these pathways. It supports the case for investing in evaluation research and interventions that can address bullying exposure and poor mental health among ALHIV in low-resourced settings, as well as programs that strengthen parenting and caregiver-adolescent interaction. Improving adolescent HIV treatment adherence should be a key public health priority

in Southern Africa and globally if we are to protect adolescent health and move closer to achieving the Sustainable Development Goals' health targets ⁷⁹.

REFERENCES

1. UNICEF. *For every child, end AIDS: seventh stocktaking report*. New York: UNICEF;2016.
2. UNICEF. Turning the tide against AIDS will require more concentrated focus on adolescents and young people. 2018; <https://data.unicef.org/topic/adolescents/hiv-aids/>. Accessed 5th August, 2018.
3. Adejumo O, Malee K, Ryscavage P, Hunter S, Taiwo B. Contemporary issues on the epidemiology and antiretroviral adherence of HIV-infected adolescents in sub-Saharan Africa: a narrative review. *Journal of the International AIDS Society*. 2015;18:20049.
4. Slogrove AL, Sohn AH. The global epidemiology of adolescents living with HIV: time for more granular data to improve adolescent health outcomes. *Curr Opin HIV AIDS*. 2018;13(3):170-178.
5. Kim MH, Mazenga AC, Yu X, Ahmed S, Paul ME, Kazembe PN, Abrams EJ. High self-reported non-adherence to antiretroviral therapy amongst adolescents living with HIV in Malawi: barriers and associated factors. *JIAS*. 2017; 20(1): 21437.
6. *[reference with identifying information omitted]*
7. Richardson D, Fen Hiu C. *Developing a Global Indicator on Bullying of School-aged Children*. Italy: UNICEF;2018.
8. Biswas T, Scott J, Munir K, et al. Global variation in the prevalence of bullying victimisation amongst adolescents: Role of peer and parental supports. *EClinicalMedicine*. 2020;<https://doi.org/10.1016/j.eclinm.2020.100276>.

9. Takizawa R, Maughan B, Arseneault L. Adult health outcomes of childhood bullying victimization: evidence from a five-decade longitudinal British birth cohort. *Am J Psychiatry*. 2014;17(1):777-784.
10. Hansson E, Garmy P, Vilhjálmsdóttir R, Kristjánssdóttir G. Bullying, health complaints, and self-rated health among school-aged children and adolescents. *J Int Med Res*. 2020;48(2):0300060519895355.
11. Richter M, Bowles D, Melzer W, Hurrelmann K. Bullying, psychosocial health and risk behaviour in adolescence. *Gesundheitswesen*. 2007;69(8-9):475-482.
12. Arseneault L, Bowes L, Shakoor S. Bullying victimization in youths and mental health problems: 'much ado about nothing'? *Psychological Medicine*. 2010;40(5):717-729.
13. Boyes M, Bowes L, Cluver L, Ward C, Badcock N. Bullying victimisation, internalising symptoms, and conduct problems in South African children and adolescents: a longitudinal investigation. *Journal of Abnormal Child Psychology*. 2014;42(8):1313-1324.
14. Moore S, Norman R, Suetani S, Thomas H, Sly P, Scott J. Consequences of bullying victimization in childhood and adolescence: A systematic review and meta-analysis. *World J Psychiatry*. 2017;7(1):60–76.
15. Human Sciences Research Council. *The People Living With HIV Stigma Index: South Africa*. Pretoria: Human Sciences Research Council;2014.
16. Pantelic M, Boyes M, Cluver L, Meinck F. HIV, violence, blame and shame: pathways of risk to internalized HIV stigma among South African adolescents living with HIV. *J Int AIDS Soc*. 2017;20(1):21771.
17. [reference with identifying information omitted]

18. Hoare J, Phillips N, Brittain K, Myer L, Zar H, Stein D. Mental Health and Functional Competence in the Cape Town Adolescent Antiretroviral Cohort. *J Acquir Immune Defic Syndr*. 2019;81(4):e109-e116.
19. Buckley J, Otjombe K, Joyce C, et al. Mental Health of Adolescents in the Era of Antiretroviral Therapy: Is There a Difference Between HIV-Infected and Uninfected Youth in South Africa? *The Journal of adolescent health*. 2020;S1054-139X(20)30037-9 [Epub ahead of print].
20. Mellins C, Malee K. Understanding the mental health of youth living with perinatal HIV infection: lessons learned and current challenges. *J Int AIDS Soc*. 2013;16(18593).
21. Ashaba S, Cooper-Vince C, Maling S, Rukundo G, Akena D, Tsai A. Internalized HIV stigma, bullying, major depressive disorder, and high-risk suicidality among HIV-positive adolescents in rural Uganda. *Global Mental Health*. 2018;5:e22.
22. [reference with identifying information omitted]
23. Dow DE, Turner EL, Shayo AM, Mmbaga B, Cunningham CK, O'Donnell K. Evaluating mental health difficulties and associated outcomes among HIV-positive adolescents in Tanzania. *AIDS Care*. 2016;28(7):825-833.
24. Mutumba M, Musiime V, Lepkwoski JM, et al. Examining the relationship between psychological distress and adherence to anti-retroviral therapy among Ugandan adolescents living with HIV. *AIDS Care*. 2016;28(7):807-815.
25. Smith Fawzi MC, Ng L, Kanyanganzi F, et al. Mental Health and Antiretroviral Adherence Among Youth Living With HIV in Rwanda. *Pediatrics*. 2016;138(4).

26. Sherr L, Cluver LD, Toska E, He E. Differing psychological vulnerabilities among behaviourally and perinatally HIV infected adolescents in South Africa - implications for targeted health service provision. *AIDS Care*. 2018;1-10.
27. Bronfenbrenner U. *The ecology of human development*. Cambridge, MA: Harvard University Press; 1979.
28. Bacchini D, Concetta Miranda M, Affuso G. Effects of Parental Monitoring and Exposure to Community Violence on Antisocial Behavior and Anxiety/Depression Among Adolescents. *Journal of Interpersonal Violence*. 2011;26(2):269-292.
29. Nguyen H, Nakamura K, Seino K, Al-Sobaihi S. Impact of parent-adolescent bonding on school bullying and mental health in Vietnamese cultural setting: evidence from the global school-based health survey. *BMC Psychology*. 2019;7(1):16.
30. Washington T, Rose T, Coard S, et al. Family-Level Factors, Depression, and Anxiety Among African American Children: A Systematic Review. *Child & Youth Care Forum*. 2017;46(1):137-156.
31. Paterson J, Tautolo E-S, Lusitini L, Sisk R. Cultural, individual, and familial risk and protective factors associated with depressive symptoms in Pacific youth living in New Zealand. *Social Work in Mental Health*. 2018;16(6):728-745.
32. Jantzer V, Haffner J, Parzer P, Resch F, Kaess M, Saag M. Does parental monitoring moderate the relationship between bullying and adolescent nonsuicidal self-injury and suicidal behavior? A community-based self-report study of adolescents in Germany. *BMC Public Health*. 2015;15(1):1-8.

33. Vassallo S, Edwards B, Renda J, Olsson C. Bullying in Early Adolescence and Antisocial Behavior and Depression Six Years Later: What Are the Protective Factors? *Journal of School Violence*. 2014;13(1):100-124.
34. Murphy C, Miller V. Concurrent and longitudinal associations among parenting style, responsibility, and adherence in youth with cystic fibrosis. *Children's Health Care*. 2019; <https://doi.org/10.1080/02739615.2019.1616295>.
35. Manne S, Jacobsen P, Gorfinkle K, Gerstein F, Redd W. Treatment adherence difficulties among children with cancer: the role of parenting style. *Journal of pediatric psychology*. 1993;18(1):47-62.
36. Al-Kloub M, Abed M, Al khawaldeh O, Al Tawarah Y, Froelicher E. Predictors of Non-Adherence to Follow-up Visits and Deferasirox Chelation Therapy Among Jordanian Adolescents with Thalassemia Major. *Pediatric Hematology & Oncology*. 2014;31(7):624-637.
37. Robinson E, Weaver P, Rusan C, Streisand R, Holmes C. A model of parental distress and factors that mediate its link with parental monitoring of youth diabetes care, adherence, and glycemic control. *Health Psychology*. 2016;35(12):1373-1382.
38. Smith Fawzi M, Eustache E, Oswald C, et al. Psychosocial Support Intervention for HIV-affected Families in Haiti: Implications for Programs and Policies for Orphans and Vulnerable Children. *Social Science & Medicine*. 2012;74(10):1494-1503.
39. Sithole Z, Mbizvo E, Chonzi P, et al. Virological failure among adolescents on ART, Harare City, 2017- a case-control study. *BMC infectious diseases*. 2018;18(1):469.

40. Wouters E, Van Damme W, Van Loona F, van Rensburg D, Meulemansa H. Public-sector ART in the Free State Province, South Africa: Community support as an important determinant of outcome. *Social Science & Medicine*. 2009;69(8):1177–1185.
41. Willis N, Napei T, Armstrong A, et al. Zvandiri--Bringing a Differentiated Service Delivery Program to Scale for Children, Adolescents, and Young People in Zimbabwe. *Journal of Acquired Immune Deficiency Syndromes*. 2018;78(2018 Supplement):S115-S123.
42. Vu L, Burnett-Zieman B, Banura C, et al. Increasing uptake of HIV, sexually transmitted infection, and family planning services, and reducing HIV-related risk behaviors among youth living with HIV in Uganda. *Journal of Adolescent Health*. 2017;60(2):S22-S28.
43. Kibaara C, Blat C, Lewis-Kulzer J, et al. Treatment Buddies Improve Clinic Attendance among Women but Not Men on Antiretroviral Therapy in the Nyanza Region of Kenya. *AIDS Research and Treatment*. 2016; Article ID 9124541.
44. Kunutsor S, Walley J, Katabira E, et al. Clinic Attendance for Medication Refills and Medication Adherence amongst an Antiretroviral Treatment Cohort in Uganda: A Prospective Study. *AIDS Research and Treatment*. 2010; Article ID 872396.
45. Nachega J, Chaisson, RE , Goliath R, et al. Randomized controlled trial of trained patient-nominated treatment supporters providing partial directly observed antiretroviral therapy. *AIDS*. 2010;24(9):1273–1280.

46. Finning K, Ukoumunne O, Ford T, et al. The association between child and adolescent depression and poor attendance at school: A systematic review and meta-analysis. *Journal of Affective Disorders*. 2019;245:928-938.
47. Hjorth C, Bilgrav L, Frandsen L, et al. Mental health and school dropout across educational levels and genders: a 4.8-year follow-up study. *BMC Public Health*. 2016;16(1):1-12.
48. Maalouf F, Ghandour L, Halabi F, et al. Psychiatric disorders among adolescents from Lebanon: prevalence, correlates, and treatment gap. *Social Psychiatry & Psychiatric Epidemiology*. 2016;51(8):1105-1116.
49. Duong M, Piroth L, Grappin M, et al. Evaluation of the Patient Medication Adherence Questionnaire as a tool for self-reported adherence assessment in HIV-infected patients on antiretroviral regimens. *HIV clinical trials*. 2001;2(2):128-135.
50. Lowenthal ED, Jibril HB, Sechele ML, Mathba K, Tshume O, Anabwani GM. Disclosure of HIV status to HIV-infected children in a large African treatment center: Lessons learned in Botswana. *Children and youth services review*. 2014;45:143-149
51. Ruchkin V, Schwab-Stone M, Vermeiren R. Social and Health Assessment (SAHA). *Psychometric Development Summary* 2004; <https://www.scienceopen.com/document?vid=25ea2912-527c-41fc-bfc2-a5d2558694cc>. Accessed 5 November, 2013.
52. Ward CL, Martin E, Theron C, Distiller GB. Factors affecting resilience in children exposed to violence. *South African Journal of Psychology*. 2007;37(1):165-187.

53. Kovacs M. *Childrens' Depression Inventory*. Niagara Falls, NY: Multi-Health Systems; 1992.
54. Reynolds CR, Richmond BO. What I think and feel: A revised measure of children's manifest anxiety. *Journal of Abnormal Child Psychology*. 1978;6(2):271-280.
55. [reference with identifying information omitted]
56. Elgar FJ, Waschbusch DA, Dadds MR, Sigvaldason N. Development and validation of a short form of the Alabama Parenting Questionnaire. *Journal of Child and Family Studies*. 2007;16(2):243-259.
57. [reference with identifying information omitted]
58. Hayes A. Beyond Baron and Kenny: Statistical Mediation Analysis in the New Millennium. *Commun Monogr [Internet]*. 2009;76(4):408–420.
59. Preacher K, Hayes A. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods* 2008;40:879-891.
60. Dawson JF, Richter AW. Probing three-way interactions in moderated multiple regression: Development and application of a slope difference test. *Journal of Applied Psychology*. 2006;91 (4):917-926.
61. Andersson U, Cuervo-Cazurra A, Nielsen BB. From the Editors: Explaining interaction effects within and across levels of analysis. *Journal of International Business Studies* 2014;45:1063-1071.
62. Vreeman R, Carroll A. A systematic review of school-based interventions to prevent bullying. *Arch Pediatr Adolesc Med*. 2007;161(1):78-88.

63. Silva J, Oliveira W, Mello F, Andrade L, Bazon M, Silva M. Anti-bullying interventions in schools: a systematic literature review. *Ciencia & saude coletiva*. 2017;22(7):2329-2340.
64. Devries K, Knight L, Child J, et al. The Good School Toolkit for reducing physical violence from school staff to primary school students: a cluster-randomised controlled trial in Uganda. *The Lancet Global health*. 2015;385:e378–386.
65. Arënliu A, Strohmeier D, Konjufca J, Yanagida T, Burger C. Empowering the Peer Group to Prevent School Bullying in Kosovo: Effectiveness of a Short and Ultra-Short Version of the ViSC Social Competence Program. *International Journal of Bullying Prevention*. 2020;2:65-78.
66. Sivaramana B, Nyea E, Bowes L. School-based anti-bullying interventions for adolescents in low- and middle-income countries: A systematic review. *Aggression and Violent Behavior*. 2019;45:154-162.
67. Bowes L, Maughan B, Caspi A, Moffitt T, Arseneault L. Families promote emotional and behavioural resilience to bullying: evidence of an environmental effect. *Journal of Child Psychology and Psychiatry*. 2010;51(7):809-817.
68. Mellins C, Brackis-Cott E, Dolezal C, Abrams E. The role of psychosocial and family factors in adherence to antiretroviral treatment in human immunodeficiency virus-infected children. *The Pediatric Infectious Disease Journal*. 2004;23(11):1035-1041.
69. Cluver L, Meinck F, Steinert J, et al. Parenting for Lifelong Health: a pragmatic cluster randomised controlled trial of a non-commercialised parenting programme for adolescents and their families in South Africa *BMJ Global Health*. 2018;3(1):3:e000539.

70. Kuo C, Mathews C, Giovenco D, et al. Acceptability, Feasibility, and Preliminary Efficacy of a Resilience-Oriented Family Intervention to Prevent Adolescent HIV and Depression: A Pilot Randomized Controlled Trial. *AIDS Education and Prevention*. 2020;32(1):67-81.
71. Annan J, Kazembe P, Bundervoet T, Seban J, Costigan J. *A Randomized Impact Evaluation of Village Savings and Loans Associations and Family-Based Interventions in Burundi. Final Evaluation. March 2013*. New York: International Rescue Committee;2013.
72. Barry M, Clarke A, Jenkins R, Patel V. A systematic review of the effectiveness of mental health promotion interventions for young people in low and middle income countries. *BMC Public Health*. 2013;11(13):835.
73. Davaasambuu S, Hauwadhanasuk T, Matsuo H, Szatmari P. Effects of interventions to reduce adolescent depression in low- and middle-income countries: A systematic review and meta-analysis. *Journal of psychiatric research*. 2020;123:201-215.
74. Rose-Clarke K, Bentley A, Marston C, Prost A. Peer-facilitated community-based interventions for adolescent health in low- and middle-income countries: A systematic review. *PLoS One*. 2019;14(1):e0210468.
75. World Health Organization. Global accelerated action for the health of adolescents (AA-HA!). 2017; <http://apps.who.int/iris/bitstream/handle/10665/255415/9789241512343eng.pdf;jsessionid=B2626A5AB85675D96B840C78270E4955?sequence=1>. Accessed 4 July 2020.
76. Akol A, Makumbi F, Babirye J, Nalugya J, Nshemereirwe S, Engebretsen I. Does mhGAP training of primary health care providers improve the

- identification of child-and adolescent mental, neurological or substance use disorders? Results from a randomized controlled trial in Uganda. *Global Mental Health*. 2018;5:e29.
77. Gaitho D, Kumar M, Wamalwa D, Nduku Wambua G, Nduati R. Understanding mental health difficulties and associated psychosocial outcomes in adolescents in the HIV clinic at Kenyatta National Hospital, Kenya. *Ann Gen Psychiatry*. 2018;17(29).
78. Ttofi M, Bowes L, Farrington D, Lösel F. Protective Factors Interrupting the Continuity From School Bullying to Later Internalizing and Externalizing Problems: A Systematic Review of Prospective Longitudinal Studies. *Journal of School Violence*. 2014;3(1):5-38.
79. United Nations. Transforming our world: The 2030 agenda for sustainable development. 2015.

Figure Legends

Figure 1: Results of the final moderated mediation analysis (PROCESS Model 23; n=969)




Table 1: Descriptive statistics and frequencies for the sample (n=979)

Variables		N (% sample)	
<i>Socio-demographic covariates</i>		Baseline	Follow-up
Age	10-14 years	617 (63%)	
	15-19 years	362 (37%)	
Gender	Female	539 (55.1%)	
	Male	440 (44.9%)	
Location	Rural	261 (26.7%)	
	Urban	716 (73.1%)	
	Missing	2 (0.2%)	
Housing (SES status)	Informal housing	183 (18.7%)	
	Formal housing	795 (81.2%)	
	Missing	1 (0.1%)	
<i>HIV-related covariates</i>			
Mode of transmission	Horizontal	203 (20.7%)	
	Vertical	769 (78.5%)	
	Missing	7 (0.7%)	
Time on Treatment	More than a year	808 (82.5%)	
	Less than a year	171 (17.5%)	
<i>Independent variable</i>			
Bullying victimization	Experienced bullying	580 (59.2%)	
	Did not experience bullying	399 (40.8%)	
<i>Dependent variable</i>			
Past week non-adherence	Non-adherent	335 (34.2%)	348 (35.5%)
	Adherent	644 (65.8%)	631 (64.5%)
<i>Mediator variable</i>			
Depression and anxiety symptoms scale (mean, SD)			1.85 (3.17)
<i>Potential moderating variables</i>			
Clinic support group attendance	Yes	142 (14.5%)	113 (11.5%)
	No	837 (85.5%)	866 (88.5%)
Treatment buddy	Yes	702 (71.2%)	565 (57.7%)
	No	277 (28.3%)	414 (42.3%)
School Enrolment	Yes	922 (94.2%)	867 (88.6%)
	No	57 (5.8%)	112 (11.4%)
Positive parenting	High	495 (50.6%)	469 (47.9%)
	Low	484 (49.4%)	510 (52.1%)
Parental monitoring	High	405 (41.4%)	433 (44.2%)
	Low	574 (58.6%)	546 (55.8%)

Table 2: Results of PROCESS Model 4 regressions respectively predicting psychological distress and ART treatment non-adherence at follow up (n=969ⁱ)

	Dependent variable: Psychological distress		Dependent variable: ART Non-adherence	
	<i>B</i> (unstandardized)	95 % CIs	<i>B</i> (unstandardized)	95 % CIs
<i>Socio-demographic and HIV-related covariates</i>				
Age (older)	0.447***	(0.204, 0.690)	-0.309	(-0.628, 0.010)
Female	0.202	(-0.016, 0.419)	-0.224	(-0.503, 0.056)
Rural location	-0.137	(-0.381, 0.107)	-0.286	(-0.604, 0.033)
Informal housing	-0.076	(-0.352, 0.199)	0.122	(-0.229, 0.473)
Horizontal infection	0.112	(-0.202, 0.426)	0.525**	(0.130, 0.920)
Time on treatment > 1 year	-0.108	(-0.415, 0.199)	-0.100	(-0.487, 0.287)
Non-adherence at baseline	0.143	(-0.085, 0.371)	0.449**	(0.161, 0.737)
<i>Independent variable</i>				
Bullied (baseline)	0.374***	(0.158, 0.589)	0.009	(-0.270, 0.288)
<i>Mediator</i>				
Psychological distress (follow up)	-	-	0.198***	(0.113, 0.283)
Model summary: $R=0.210$; $R-sq.=0.044$; $F = 5.553$; $p<0.001$			Model summary: Nagelkrk= 0.067 ; CoxSnell= 0.049 ; $p<0.001$	

*** $p<0.001$; ** $p<0.01$; * $p<0.05$

ⁱ 10 cases had a missing value for one or more variables and were excluded from the regression analyses

Table 3: Results of final PROCESS model 23 moderated mediation regressions, respectively predicting higher psychological distress and ART non-adherence at follow up (n=969)

	Dependent variable: Psychological distress		Dependent variable: ART Non-adherence	
	<i>B</i> (unstandardized)	95 % CIs	<i>B</i> (unstandardized)	95 % CIs
<i>Socio-demographic and HIV-related covariates</i>				
Age (older)	0.143	(-0.217, 0.503)	-	-
Female	-0.008	(-0.345, 0.329)	-	-
Rural location	-0.155	(-0.398, 0.088)	-0.307	(-0.628, 0.013)
Informal housing	-0.069	(-0.344, 0.206)	0.141	(-0.208, 0.491)
Horizontal infection	0.126	(-0.187, 0.439)	0.352	(-0.011, 0.714)
Time on treatment > 1 year	-0.102	(-0.408, 0.204)	-0.065	(-0.451, 0.322)
Non-adherence at baseline	0.147	(-0.081, 0.374)	0.383**	(0.095, 0.672)
<i>Independent Variable</i>				
Bullied (baseline)	-0.017	(-0.362, 0.329)	0.005	(-0.274, 0.283)
<i>Mediator</i>				
Psychological distress (follow up)	-	-	0.249***	(0.142, 0.356)
<i>Moderators</i>				
Older age x Bullied	0.517*	(0.068, 0.966)		
Female x Bullied	0.362	(-0.073, 0.797)	-	-
Better parental monitoring x Psychological distress	-	-	-0.220*	(-0.418, -0.023)
		Model summary: $R=0.231$; $R-sq.=0.531$; $F = 5.888$; $p<0.001$	Model summary: Nagelkrk= 0.065 ; CoxSnell= 0.047 ; $p<0.001$	

Table 4: Conditional indirect effect(s) of Bullying at baseline on Non-Adherence at follow up, at different values of the moderator(s):

Age (10-14 vs 15-19)	Gender	Parental monitoring	Effect	95 % CIs	Bullying – non-adherence relationship significant
Younger	Male	Low	- 0.004	(- 0.080, 0.070)	No
Younger	Male	High	- 0.001	(- 0.037, 0.023)	No
Younger	Female	Low	0.086	(0.011, 0.185)	Yes
Younger	Female	High	0.010	(-0.047, 0.099)	No
Older	Male	Low	0.125	(0.016, 0.279)	Yes
Older	Male	High	0.014	(-0.063, 0.143)	No
Older	Female	Low	0.215	(0.095, 0.385)	Yes
Older	Female	High	0.025	(-0.118, 0.204)	No